

Deakin Research Online

This is the published version:

Fairclough, Stuart J., Boddy, Lynne M., Ridgers, Nicola D. and Stratton, Gareth 2012, Weight status associations with physical activity intensity and physical self-perceptions in 10-to 11-year-old children, *Pediatric exercise science*, vol. 24, no. 1, pp. 100-112.

Available from Deakin Research Online:

<http://hdl.handle.net/10536/DRO/DU:30046178>

Reproduced with the kind permission of the copyright owner.

Copyright : 2012, Human Kinetics, Inc.

Weight Status Associations With Physical Activity Intensity and Physical Self-Perceptions in 10- to 11-Year-Old Children

Stuart J. Fairclough and Lynne M. Boddy

Liverpool John Moores University

Nicola D. Ridgers

Deakin University

Gareth Stratton

Liverpool John Moores University

The study examined associations between children's weight status, physical activity intensity, and physical self-perceptions. Data were obtained from 409 children (224 girls) aged 10–11 years categorized as normal-weight or overweight/obese. Physical activity was assessed using accelerometry, and children completed the Physical Self-Perception Profile. After controlling for the effects of age, maturation, and socioeconomic status vigorous physical activity was significantly associated with normal-weight status among boys (OR = 1.13, $p = .01$) and girls (OR = 1.13, $p = .03$). Normal-weight status was significantly associated with perceived Physical Condition (Boys: OR = 5.05, $p = .008$; Girls: OR = 2.50, $p = .08$), and Body Attractiveness (Boys: OR = 4.44, $p = .007$; Girls: OR = 2.56, $p = .02$). Weight status of 10–11 year old children was significantly associated with time spent in vigorous physical activity and self-perceptions of Body Attractiveness and Physical Condition.

Overweight and obesity are global public health issues implicated in the onset and development of a number of degenerative conditions through the life course (5,7). The causes of overweight and obesity are complex but the basic physiological premise is that positive energy balance over time leads to weight gain as more energy is taken in than expended (48). On this basis, increased energy expenditure through physical activity should help maintain energy balance and therefore a stable body weight. For these reasons physical activity is implicated in the cause and prevention of pediatric overweight and obesity (10). Although

Fairclough, Boddy, and Stratton are with the REACH Group and the Research Institute for Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK. Ridgers is with the REACH Group and the Centre for Physical Activity and Nutrition Research, Deakin University, Australia.

it is suggested that obesity prevalence among young people may have ‘leveled off’ in the last decade (30), the existing high incidence of obesity has coincided with low levels of habitual physical activity in most developed countries (37,39). Despite the intuitive logic that declining physical activity levels influence the increased prevalence of overweight and obesity, a recent review of children’s objectively assessed physical activity and changes in adiposity reported weak and inconsistent associations between the variables of interest (48). One factor implicated in the equivocal conclusions reached was a lack of studies assessing specific intensities of physical activity (48). In particular, the influences of moderate (MPA), and vigorous intensity physical activity (VPA) on adiposity may differ, with more intense activity often reported to be of greater benefit (24). Self-perceptions contribute to a person’s self-esteem, which reflects their evaluation of the good or worth inherent in their self-description (45). With regard to health-related changes to lifestyle, self-esteem and motivation are interlinked and can be viewed as ‘common concepts’ (46). Competence motivation theory (16) suggests that people will be motivated to engage in activities where they feel competent, and on this basis self-perceptions are important predictors of physical activity and other volitional behaviors (44). Physical self-perceptions such as perceived physical competence are recognized as positive correlates of youth physical activity (42), and specific components of the physical self are thought to influence motivated behavior (11).

Studies have shown significant associations between physical self-perceptions and physical activity in youth, with boys typically reporting stronger self-perceptions than girls (9). Furthermore, associations exist between overweight/obesity and physical activity (1), and evidence suggests that overweight and obese children may have lower self-perceptions than normal-weight counterparts (23). Moreover, excess adiposity impacts on selected domains of self-perception, which often relate to physical activity participation and physical appearance (12). The relationships between specific components of children’s physical self-perceptions, physical activity, and weight status are under researched but improved understanding of these factors may help inform future childhood obesity interventions. Thus, the study purpose was to examine associations between weight status, physical activity intensities, and physical self-perceptions among English primary school children.

Methods

Participants

Eight schools situated in urban and suburban areas of a large north-west England town were recruited to the study during the autumn school terms of 2008 and 2009. In these schools all children in Year 6 (aged 10–11 years; $N = 602$; 307 in 2008, 295 in 2009) received project and consent information and were invited into the study. Five hundred children consented to participate (230 in 2008 (116 girls), 270 in 2009 (139 girls)) resulting in an 83.1% participation rate. The ethnic origin of all the children was white British, which roughly reflects the ethnic demographic of the town’s school-age population which is 96% white British (47). During 2008 and 2009 data were collected in one school per week between October and December. Ethical approval was obtained from the University Ethics Committee.

Instruments and Procedures

Socioeconomic Status Socioeconomic status was calculated using the 2007 Indices of Multiple Deprivation (IMD; 6). IMD scores were derived from the children's home postcodes using the National Statistics Postcode Directory database (2).

Anthropometry Stature and sitting height were measured to the nearest 0.1 cm using a portable stadiometer (Leicester Height Measure, Seca, Birmingham, UK). Leg length was calculated by subtracting sitting height from stature. Body mass was measured to the nearest 0.1 kg using calibrated scales (Seca, Birmingham, UK). All measurements were taken by trained research staff with the children in light clothing and barefooted. The International Obesity Task Force age and sex-specific BMI cut-points were used to classify children as either normal-weight, overweight, or obese (3). Overweight and obese children's data were then collapsed to create an overweight/obese grouping category.

Maturity Status Somatic maturity status was estimated by determining years from attainment of peak height velocity. Attainment of peak height velocity reflects the age at maximum growth rate in stature during adolescence. Maturity offset (i.e., years from attainment of peak height velocity) for each child was predicted using sex-specific regression equations that included stature, sitting height, leg length, chronological age and their interactions (19). This noninvasive method has demonstrated acceptable agreement when correlated against skeletal age ($r = .83$; 19) and is commonly used in physical activity research (9,32)

Physical Self-Perceptions Physical self-perceptions were assessed using the Children and Youth version of the Physical Self-Perception Profile (CY-PSPP; 45). The CY-PSPP follows a hierarchical structure with global self-esteem at the apex and Physical Self-Worth positioned at the domain level. Subordinate to Physical Self-Worth are four subdomains of Sport Competence, Physical Condition, Body Attractiveness, and Physical Strength. Each domain is measured on a 1–4 scale by six items that use a structured alternative format to reduce socially desirable responses. The CY-PSPP was administered in the children's classrooms by research staff who provided verbal and visual examples of how and where to respond to items on the profile.

Physical Activity Physical activity was objectively measured every 5 s using ActiGraph accelerometers (GT1M, ActiGraph LLC, Pensacola, FL). The children were instructed to wear the ActiGraph over the right hip using a waist mounted nylon belt, from waking in the morning until bedtime. In 2008 children were asked to wear the accelerometers for 5 consecutive days, including 2 weekend days. In 2009 increased availability of ActiGraphs meant that the wear time protocol increased to 7 consecutive days, including 2 weekend days. At the end of the data collection period ActiGraphs were downloaded using Actlife software (ActiGraph LLC, Pensacola, FL). Downloaded files were initially checked for compliance to the monitoring protocol using customized software (MAHUFFe; www.mrc-epid.cam.ac.uk). Sustained 20 min periods of zero counts indicated that the ActiGraph had been removed, and total 'missing' counts for those periods represented the duration that monitors were not worn. Children were included in the data analysis if they wore the monitors for at least 600 min each day for a minimum of 3 days

(18). Published cut-points for different intensity levels in children vary substantially [e.g., between 906 (41) and 3200 counts \cdot min⁻¹ for MPA (26)], and in the absence of individual calibration data there is no consensus as to the most appropriate cut-points to use. Thus, we chose the values reported by Ekelund and colleagues (2007) in the European Youth Heart Study (8) because they fell between the most extreme values reported (26,41) and were appropriate to the age group of interest. The ActiGraph count cut-points for minutes spent in sedentary activity, MPA and VPA were <500, 2000, and 4000 counts \cdot min⁻¹, respectively (8).

Data Analysis Descriptive statistics were calculated for all measured variables and weight status differences were examined by independent *t* tests. Sex-specific binomial logistic regression was used to establish the association between normal weight status, and physical activity and physical self-perceptions. Analyses were adjusted for age, maturity status, and socioeconomic status. Specific predictor variables were minutes spent in sedentary time, MPA, and VPA, and perceived Sport Competence, Physical Condition, Body Attractiveness, Physical Strength, and Physical Self-Worth. Effect sizes were computed as adjusted odds ratios (ORs) and 95% confidence intervals. All analyses were conducted using SPSS v. 15 (SPSS Inc; Chicago, IL) and statistical significance was set at $p < .05$.

Results

Preliminary analyses confirmed that the data were normally distributed, and that the assumptions of linearity of logit and collinearity (Tolerance and VIF statistics) were met. Strong internal consistencies were demonstrated for each of the physical self-perception subdomains. Cronbach's alpha coefficients for boys and girls ranged from 0.75–0.76 (Sport Competence), 0.78–0.79 (Physical Condition), 0.83–0.87 (Body Attractiveness), 0.76–0.83 (Physical Strength), and 0.74–0.84 (Physical Self-Worth). In 2008 and 2009, 47 (20.4%) and 44 (16.3%) of children, respectively did not meet the minimum ActiGraph wear time criteria and so were excluded from the data set. The main reason for exclusion was noncompliance, but one ActiGraph was lost in 2008 and six malfunctioned in 2009. Overall, 409 children were included in the analyses (224 girls) giving an overall compliance rate of 81.8%. There were significant differences in time spent sedentary and in MPA between included and excluded children (sedentary: excluded = 339.9 min vs. included = 578.5 min, $p < .0001$; MPA: excluded = 31.2 min vs. included = 40.0 min, $p < .0001$). Nineteen percent of children provided data on 7 valid measurement days, 15.4% on 6 days, 33.4% on 5 days, 19.5% on 4 days, and 12.7% on 3 days. Physical activity levels did not differ between children with 7, 6, 5, 4, or 3 days of valid data. Mean duration of daily monitoring was 11.8 hr for boys and girls.

Descriptive Analyses

There were no significant differences in age, stature, body mass, BMI, or maturation between the 2008 and 2008 cohorts ($p < .05$). Overweight/obese boys and girls were significantly taller, heavier, and somatically more mature than normal weight peers (Table 1). The majority of children were categorized as normal-weight (Boys: 78.5%; Girls: 76.2%), with 18.3% and 18.8% of boys and girls, respectively overweight, and 3.2% (boys) and 4.9% (girls) categorized as obese. These values

are broadly representative of the town's school-age population when the IOTF BMI cut-points are applied (36). Stature and body mass were similar for boys and girls although girls' maturity offset values (-1.26 years) were more advanced than boys (-2.97 years).

Sedentary time was similar between sex and weight status groups (Table 2). Normal-weight and overweight/obese boys' MPA differed by around 0.7 min, compared with 1.3 min for girls. Significant differences in VPA were observed among normal-weight and overweight/obese boys ($t_{185} = 3.93, p = .001$) and girls ($t_{221} = 2.68, p = .008$). Sixty six percent of normal-weight boys achieved the recommended minimum of 60 min moderate-to-vigorous physical activity (MVPA) per day, compared with 47.5% of overweight/obese boys ($\chi^2_1 = 4.56, p = 0.03$). Sixty minutes of MVPA was achieved by 30% and 22.6% of normal-weight and overweight/obese girls, respectively ($\chi^2_1 = 1.08, p = 0.30$). Significant differences in normal-weight and overweight/obese boys' and girls' physical self-perceptions were evident in all subdomains, with the exception of Sport Competence, and Physical Strength (Table 2). The greatest differences in normal-weight and overweight/obese boys' and girls' physical self-perceptions were in perceived Physical Condition and Body Attractiveness.

Main Analyses

After controlling for the effects of boys' age, maturation, and socioeconomic status, normal-weight status was significantly associated with VPA but not with MPA or sedentary time (OR = 1.13 (1.03, 1.23), $p = .01$; Table 3). Furthermore, normal-weight status was significantly associated with boys' self-perceptions of Physical Condition (OR = 5.05 (1.52, 16.75), $p = .008$) and Body Attractiveness (OR = 4.44 (1.51, 13.06), $p = .007$). Following adjustment for girls' age, maturation and socioeconomic status, analysis revealed a significant association between normal-weight status and VPA (OR = 1.13 (1.02, 1.25), $p = .03$, but not MPA or sedentary time (Table 3). Moreover, among girls, normal-weight status was significantly associated with self-perceptions of Physical Condition (OR = 2.50 (0.90, 6.98), $p = .08$) and Body Attractiveness (OR = 2.56 (1.14, 5.71), $p = .02$).

Discussion

This study investigated associations between primary school children's weight status, physical activity levels and physical self-perceptions. VPA was significantly different between normal-weight and overweight/obese children and VPA was significantly and positively associated with normal-weight status, although the strength of the effect sizes was modest. Normal-weight children accumulated 5.8 min (boys) and 3.04 min (girls) more VPA per day than their overweight/obese peers. Differences in VPA between normal-weight and overweight and obese children have previously been reported in the range of 6.4 min for boys and girls combined (40), 2 min for boys (35), and ~1 min for girls (38). Comparisons with the current study are difficult due to the different accelerometer protocols, analyses, and cut-points used to classify VPA. Two studies (35,38) used higher VPA cut-points than in our investigation, which may explain in part why smaller differences in VPA were observed. Although Trost and colleagues used a similar VPA cut-point to the

Table 1 Descriptive Characteristics of Normal-Weight and Overweight/Obese Boys and Girls (*M* ± *SD*)

	Boys		Girls		<i>p</i>
	Normal-weight (<i>n</i> = 146)	Overweight/obese (<i>n</i> = 40)	Normal-weight (<i>n</i> = 170)	Overweight/obese (<i>n</i> = 53)	
Age (years)	10.66 (0.33)	10.74 (0.30)	10.65 (0.33)	10.66 (0.32)	0.80
Stature (cm)	142.72 (7.37)	146.44 (5.03)	143.75 (6.97)	146.88 (6.80)	0.005
Body mass (kg)	34.43 (5.88)	49.02 (6.38)	34.98 (5.97)	49.42 (6.60)	<0.0001
BMI (<i>m</i> • <i>kg</i> ⁻²)	16.80 (1.77)	22.80 (2.17)	16.83 (1.93)	22.95 (3.31)	<0.0001
Maturity offset (years)	-3.07 (0.48)	-2.59 (0.38)	-1.38 (0.50)	-0.87 (0.49)	< .0001
^a SES (^b IMD score)	18.60 (11.63)	17.82 (10.69)	17.22 (10.43)	20.10 (13.18)	0.11

^aSES = socioeconomic status; ^bIMD = indices of multiple deprivation

Table 2 Objectively Assessed Physical Activity and Physical Self-Perceptions of Normal-Weight and Overweight/Obese Boys and Girls ($M \pm SD$)

	Boys			Girls		
	Normal-weight (<i>n</i> = 146)	Overweight/obese (<i>n</i> = 40)	<i>p</i>	Normal-weight (<i>n</i> = 170)	Overweight/obese (<i>n</i> = 53)	<i>p</i>
Physical activity (min/day)						
Sedentary	572.33 (74.02)	582.64 (78.90)	0.44	590.75 (68.25)	581.31 (77.90)	0.40
^a MPA	43.23 (11.16)	42.46 (11.58)	0.70	36.60 (8.25)	35.33 (8.25)	0.37
^b VPA	23.79 (10.12)	17.99 (7.26)	0.001	17.92 (7.53)	14.88 (5.93)	0.008
Physical self-perceptions						
Sport Competence	3.11 (0.64)	2.91 (0.60)	0.09	2.93 (0.56)	2.73 (0.54)	0.03
Physical Condition	3.18 (0.64)	2.69 (0.51)	<0.0001	3.02 (0.58)	2.75 (0.59)	0.004
Attractive Body	2.91 (0.68)	2.38 (0.59)	<0.0001	2.84 (0.67)	2.31 (0.54)	<0.0001
Physical Strength	2.91 (0.67)	2.86 (0.50)	0.60	2.65 (0.60)	2.68 (0.54)	0.79
Physical Self-Worth	3.07 (0.64)	2.82 (0.56)	0.03	3.02 (0.61)	2.77 (0.60)	0.01

^aMPA = moderate intensity physical activity; ^bVPA = vigorous intensity physical activity

Table 3 Associations Between Boys' and Girls' Physical Activity, Physical Self-Perceptions and Normal Weight Status

Predictor	Boys			Girls		
	B (SE)	p	OR 95% CI	B (SE)	p	OR 95% CI
Constant	-12.16 (11.45)			-34.31 (10.21)		
Physical activity (min)						
Sedentary	0.004 (0.004)	0.22	1.0 1.0, 1.01	0.002 (0.003)	0.48	1.0 1.0, 1.01
^a MPA	-0.06 (0.03)	0.05	0.89, 1.0	-0.04 (0.03)	0.19	0.96 0.90, 1.02
^b VPA	0.12 (0.05)	0.01	1.03, 1.23	0.12 (0.05)	0.03	1.13 1.02, 1.25
Physical self-perceptions						
Sport competence	-0.39 (0.67)	0.56	0.18, 2.53	-0.54 (0.59)	0.36	0.58 0.18, 1.85
Physical condition	1.62 (0.61)	0.008	5.05 1.52, 16.75	0.92 (0.51)	0.08	2.50 0.90, 6.98
Attractive body	1.49 (0.55)	0.007	4.44 1.51, 13.06	0.94 (0.41)	0.02	2.56 1.14, 5.71
Physical strength	-0.67 (0.52)	0.20	0.51 0.19, 1.41	-0.69 (0.51)	0.17	0.50 0.19, 1.35
Physical self-worth	-1.03 (0.68)	0.13	0.36 0.09, 1.36	-0.61 (0.60)	0.31	0.54 0.17, 1.77

^aMPA = moderate intensity physical activity; ^bVPA = vigorous intensity physical activity

one used in our study, boys' and girls' VPA was combined which precludes comparisons by gender. These observed differences in VPA may not have accurately reflected energy expenditure of the normal-weight and overweight/obese groups. It is reported that the energy cost of participating in higher intensity physical activity can be greater for overweight/obese or low fit youth than for leaner or fitter peers (33). Using age-related reference values for children's physical activity energy expenditure (15) we estimated that 5.8 min VPA (i.e., walking at $5.6 \text{ km} \cdot \text{h}^{-1}$) for the normal-weight boys would yield an energy cost of 24.6 kcal, but the same duration of MPA (i.e., walking at $4 \text{ km} \cdot \text{h}^{-1}$) for the overweight/obese boys would require a greater energy cost at 26.2 kcal. Thus, it is possible that the overweight/obese children were doing more VPA than indicated by the accelerometer counts in terms of energy expended, but this was misclassified as MPA because accelerometers are unable to assess the physiological strain of physical activity. This issue may have partially influenced the magnitude of the associations between VPA and weight status.

MPA was not associated with weight status, which concurs with previous studies (49), although others have reported moderate correlations between MPA and body fat (31). Our results suggest that regardless of weight status most children in the sample engaged in health-enhancing physical activity at the minimal, moderate intensity level. Sedentary time did not differ by weight status which supports a recent investigation into associations between children's objectively assessed sedentary time and body composition (34). Although significant relationships between self-reported time spent TV viewing and video gaming, and body fat have been observed, the reported effect sizes were small, possibly due to the use of single, self-report, discrete markers of sedentary behavior, rather than objective measures (17). Differences in the measurement and classification of sedentary activity and weight status, as well as the affect of confounders such as socioeconomic status, sleep duration, and food intake may be contributory factors explaining the lack of consistency between studies investigating sedentary behaviors and weight status (21).

Normal-weight children reported significantly stronger physical self-perceptions than overweight/obese peers. Furthermore, overweight/obese girls scored lower in all domains than overweight/obese boys. Similar weight status and gender-related differences in self-perceptions have been observed elsewhere, particularly in relation to perceived physical competence and appearance (14). Body dissatisfaction is common among overweight/obese youth and is related to comparisons with societal body shape ideals and resultant social stigmatization (13). Recently it was reported how body dissatisfaction mediated associations between different aspects of physical self-perception and BMI in boys but not in girls (20). The authors of this study suggested that improved physical self-perceptions achieved through physical activity experiences could reduce body dissatisfaction of overweight and obese boys, but in girls body dissatisfaction might be linked more to aesthetic appearance than physical activity-related competence, hence the stronger mediating effects of physical self-perceptions among boys (20). Considering the influence of body dissatisfaction the significant associations between perceived Body Attractiveness and weight status were anticipated, although the magnitude of these effects (Boys: OR = 4.44; Girls: OR = 2.54) revealed differences in the relative influence of Body Attractiveness between the sexes. These effect size differences reflect the higher ratings of Body Attractiveness among boys, which were consistent

with other studies in similarly aged children (4,9). Although girls typically attach more importance to Body Attractiveness (46) their perceptions are often unable to match their expectations, possibly because pressure to conform to a societal and cultural stereotype body shape is greater on girls than boys, and girls experience more obvious physical changes during puberty than boys (22).

Perceived Physical Condition was also significantly associated with normal-weight status, with these associations strongest in boys (OR = 5.05 vs. 2.5 in girls). The strength of the associations observed for the normal-weight boys and girls reflects the relative differences in their ratings of Physical Condition, which concur with those described previously (9). In the current sample perceptions of Physical Condition, conceptualized in the CY-PSPP as fitness and exercise (45) were lowest among children with the highest BMI values. Previous studies have reported moderate correlations between perceived Physical Condition and cardiorespiratory fitness (44), while the inverse association between cardiorespiratory fitness and adiposity/weight status is well established (8). Thus, it is likely that the children's perceived Physical Condition was based in part on their actual levels of cardiorespiratory fitness (44). These perceptions of Physical Condition may have been manifested through class-based norm-referenced assessments (e.g., multistage fitness test) and/or direct comparisons with peers during play and physical activity (e.g., effort and exertion during sports or vigorous games; 43). Although cardiorespiratory fitness was not assessed in this study the strong associations between perceived Physical Condition and weight status may reflect differing levels of cardiorespiratory fitness between the normal-weight and overweight/obese children, which have been observed previously (28). Cardiorespiratory fitness is inversely and independently associated with clustered cardiovascular risk, although fatness may partly mediate this association (29). Though physical activity engagement is beneficial to both cardiorespiratory fitness and clustered cardiovascular risk, actual engagement in activity may be compromised among overweight and obese children without a reduction in abdominal fat (29). Nonetheless, physical activity interventions can be effective among children with unfavorable cardiovascular risk profiles, when such protocols are of sufficient duration, have appropriate volumes of activity, and are implemented by expert personnel (27).

Study Limitations and Strengths

This study had several limitations which need acknowledging. The cross-sectional nature of the study precludes any claims about causality and direction in relation to the variables of interest. The selection of accelerometer cut-points had an influence on the extent of the associations with weight status. As there is no consensus about the most appropriate accelerometer cut-points to use with pediatric populations we employed previously published values (8) that have been used to address similar research questions to ours with comparable populations (34). Moreover, accelerometers assess movement and are unable to assess the physiological strain of physical activity. For this reason the energy cost of the normal-weight and overweight/obese children's physical engagement was unknown. Data were collected in the autumn and winter when temperatures averaged 8.5 °C and daylight hours were limited to between 10 and 7.8 hr. For these reasons physical activity levels may have been lower than during the spring and summer. Although BMI has

limitations as a measure of adiposity, it is strongly associated with body fat among youth (25) and so is a most appropriate measure for school-based studies such as this. Strengths of this study were the use of a large sample of children, objective measures of physical activity to assess physical activity at different intensities, and analyses which were adjusted for common confounders.

Conclusions

This novel investigation studied associations between children's weight status, different intensities of physical activity, and physical self-perceptions. Significant modest associations were observed between normal-weight status and time spent in VPA. Time spent sedentary and in MPA were similar regardless of weight status. Normal-weight children had more positive physical self-perceptions than overweight/obese counterparts, particularly with regards to perceptions of Body Attractiveness and Physical Condition. Moreover, the magnitude of these associations was greatest among boys. Future research needs to confirm causal relationships between the correlates investigated in this study through longitudinal and interventional type investigations.

References

1. Antonogeorgos, G., A. Papadimitriou, D.B. Panagiotakos, K.N. Priftis, and P. Nikolaidou. Physical activity patterns and obesity status among 10- to 12- year old adolescents living in Athens, Greece. *J Phys Act Health*. 7:633–640, 2010.
2. Census.ac.uk. *National Statistics Postcode Directory*. Available from: http://census.ac.uk/guides/Lookup_tables.aspx, 2008.
3. Cole, T.J., M.C. Bellizzi, K.M. Flegal, and W.H. Dietz. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 320:1240–1244, 2000.
4. Crocker, P.R.E., R.C. Ekelund, and K.C. Kowalski. Children's physical activity and physical self-perceptions. *J Sports Sci*. 18:383–394, 2000.
5. Deckelbaum, R.J., and C.L. Williams. Childhood obesity: the health issue. *Obes. Res. Suppl*. 9(4):239–243, 2001.
6. Department for Communities and Local Government. *The English Indices of Deprivation 2007*. Wetherby: Communities and Local Government Publications, 2008.
7. Dietz, W.H. Health consequences of obesity in youth: childhood predictors of adult disease. *Pediatrics*. 101:518–525, 1998.
8. Ekelund, U., S.A. Anderssen, K. Froberg, L.B. Sardinha, L.B. Andersen, and S. Brage. Independent associations of physical activity and cardiorespiratory fitness with metabolic risk factors in children: the European Youth Heart Study. *Diabetologia*. 50:1832–1840, 2007.
9. Fairclough, S.J., and N.D. Ridgers. Relationships between maturity status, physical activity, and physical self-perceptions in primary school children. *J Sports Sci*. 28:1–9, 2010.
10. Fox, K.R. Tackling obesity in children through physical activity: a perspective from the United Kingdom. *Quest*. 56:28–40, 2004.
11. Fox, K.R., and C.B. Corbin. The physical self-perception profile: Development and preliminary validation. *J Sport Exer Psychol*. 11:408–430, 1989.
12. Franklin, J., G. Denyer, K.S. Steinbeck, I.D. Caterson, and A.J. Hill. Obesity and Risk of Low Self-esteem: A Statewide Survey of Australian Children. *Pediatrics*. 118:2481–2487, 2006.

13. Friedman, M.A., and K.D. Brownell. Psychological correlates of obesity: moving to the next research generation. *Psychol. Bull.* 117:3–20, 1995.
14. Griffiths, L.J., T.J. Parsons, and A.J. Hill. Self-esteem and quality of life in obese children and adolescents: A systematic review. *Int. J. Pediatr. Obes.* 5:282–304, 2010.
15. Harrell, J.S., R.G. McMurray, C.D. Baggett, M.L. Pennell, P.F. Pearce, and S.I. Bangdiwala. Energy costs of physical activities in children and adolescents. *Med. Sci. Sports Exerc.* 37:329–336, 2005.
16. Harter, S. The Perceived Competence Scale for Children. *Child Dev.* 53:87–97, 1982.
17. Marshall, S.J., S.J.H. Biddle, T. Gorely, N. Cameron, and I. Murdey. Relationships between media use, body fatness and physical activity in children and youth: A meta-analysis. *Int. J. Obes.* 28:1238–1246, 2004.
18. Mattocks, C., A.R. Ness, S.D. Leary, et al. Use of accelerometers in a large field-based study of children: Protocols, design issues, and effects on precision. *J Phys Act Health.* 5:S98–S111, 2008.
19. Mirwald, R.L., A.D.G. Baxter-Jones, D.A. Bailey, and G.P. Beunen. An assessment of maturity from anthropometric measurements. *Med. Sci. Sports Exerc.* 34:689–694, 2002.
20. Morano, M., D. Colella, C. Robazza, L. Bortoli, and L. Capranica. Physical self-perception and motor performance in normal-weight, overweight and obese children. *Scan. J. Med. Sci. Sports.* 2010.
21. Must, A., and D.J. Tybor. Physical activity and sedentary behavior: a review of longitudinal studies of weight and adiposity in youth. *Int. J. Obes.* 29(Suppl. 2):S84–S96, 2005.
22. Niven, A.G., S.G. Fawcner, A.M. Knowles, and C. Stephenson. Maturational differences in physical self-perceptions and relationships with physical activity in early adolescent girls. *Pediatr. Exerc. Sci.* 19:472–480, 2007.
23. O’Dea, J.A. Self-concept, self-esteem and body weight in adolescent females. *J. Health Psychol.* 11:599–611, 2006.
24. Parikh, T., and G. Stratton. Influence of intensity of physical activity on adiposity and cardiorespiratory fitness in 5-18 year olds. *Sports Med.* 41:477–488, 2011.
25. Pietrobelli, A., M.S. Faith, D.B. Allison, D. Gallagher, G. Chiumello, and S.B. Heymsfield. Body mass index as a measure of adiposity among children and adolescents: A validation study. *J. Pediatr.* 132:204–210, 1998.
26. Puyau, M., A. Adolph, F. Vohra, and N. Butte. Validation and calibration of physical activity monitors in children. *Obes. Res.* 10:150–157, 2002.
27. Resaland, G.K., S.A. Anderssen, I.M. Holme, A. Mamen, and L.B. Andersen. Effects of a 2-year school-based daily physical activity intervention on cardiovascular disease risk factors: The Sogndal school-intervention study. *Scand J Med Sci Sports.* 2010.
28. Resaland, G.K., A. Mamen, S.A. Anderssen, and L.B. Andersen. Cardiorespiratory fitness and body mass index values in 9-year-old Norwegian children. *Acta Paediatr.* 98:687–692, 2008.
29. Resaland, G.K., A. Mamen, C. Boreham, S.A. Anderssen, and L.B. Andersen. Cardiovascular risk factor clustering and its association with fitness in nine-year-old rural Norwegian children. *Scand J Med Sci Sports.* 20:e112–e120, 2010.
30. Rokholm, B., J.L. Baker, and T.I. Sorensen. The levelling off of the obesity epidemic since the year 1999 - a review of evidence and perspectives. *Obes. Rev.* 11:835–846, 2010.
31. Rowlands, A.V., D.K. Ingledeu, and R.G. Eston. The relationship between body fatness and habitual physical activity in children: A meta-analysis. *Pediatr. Exerc. Sci.* 11:263, 1999.
32. Sherar, L.B., D.W. Esliger, A.D.G. Baxter-Jones, and M.S. Trembaly. Age and gender differences in youth physical activity: Does physical maturity matter? *Med. Sci. Sports Exerc.* 39:830–835, 2007.

33. Spadano, J.L., A. Must, L.G. Bandini, G.E. Dallal, and W.H. Dietz. Energy cost of physical activities in 12 y old girls: MET values and the influence of body weight. *Int. J. Obes.* 27:1528–1533, 2003.
34. Steele, R.M., E.M. van Sluijs, A. Cassidy, S.J. Griffin, and U. Ekelund. Targeting sedentary time or moderate- and vigorous-intensity activity: independent relations with adiposity in a population-based sample of 10-y-old British children. *Am. J. Clin. Nutr.* 90:1185–1192, 2009.
35. Stone, M.R., A.V. Rowlands, and R.G. Eston. Characteristics of the activity pattern in normal weight and overweight boys. *Prev. Med.* 49:205–208, 2009.
36. The NHS Information Centre. *National Child Measurement Programme: England 2009/10 school year*. London: NHS, 2010.
37. The NHS Information Centre. *Statistics on Obesity, Physical Activity and Diet: England, 2010*. London: The Health and Social Care Information Centre, 2010.
38. Treuth, M.S., D.J. Catellier, K.H. Schmitz, et al. Weekend and weekday patterns of physical activity in overweight and normal-weight adolescent girls. *Obes.* 15:1782–1788, 2007.
39. Troiano, R.P., D. Berrigan, K.W. Dodd, L.C. Masse, T. Tilert, and M. McDowell. Physical activity in the United States measured by accelerometer. *Med. Sci. Sports Exerc.* 40:181–188, 2008.
40. Trost, S.G., L. Kerr, and R.R. Pate. Physical activity and determinants of physical activity in obese and non-obese children. *Int. J. Obes.* 25:822–829, 2001.
41. Trost, S.G., R.R. Pate, J.F. Sallis, et al. Age and gender differences in objectively measured physical activity in youth. *Med. Sci. Sports Exerc.* 34:350–355, 2002.
42. Van der Horst, K., M.J.C.A. Paw, J.W.R. Twisk, and W. Van Mechelen. A brief review on correlates of physical activity and sedentariness in youth. *Med. Sci. Sports Exerc.* 39:1241–1250, 2007.
43. Weiss, M.R., V. Ebbeck, and T.S. Horn. Children's self-perceptions and sources of physical competence information: A cluster analysis. *J Sport Exer Psychol.* 19:52–70, 1997.
44. Welk, G.J., and B. Eklund. Validation of the children and youth physical self perceptions profile for young children. *Psychol. Sport Exerc.* 6:51–65, 2005.
45. Whitehead, J.R. A study of children's physical self-perceptions using an adapted physical self-perception profile questionnaire. *Pediatr. Exerc. Sci.* 7:132–151, 1995.
46. Whitehead, J.R., and C.B. Corbin. Self-esteem in children and youth: The role of sport and physical education. In: *The Physical Self: From Motivation to Well-Being*, K.R. Fox (Ed.). Champaign, IL: Human Kinetics, 1997, pp. 175–203.
47. WiganCouncil. *Equality and Diversity - Welcome*. Available from: <http://www.wigan.gov.uk/Services/CouncilDemocracy/PoliciesPlans/EqualOpportunities/EqualityAnd-DiversityWelcome.htm>, 2011.
48. Wilks, D.C., H. Besson, A.K. Lindroos, and U. Ekelund. Objectively measured physical activity and obesity prevention in children, adolescents and adults: a systematic review of prospective studies. *Obes. Rev.* 2010.
49. Wittmeier, K.D., R.C. Mollard, and D.J. Kriellaars. Physical activity intensity and risk of overweight and adiposity in children. *Obes.* 16:415–420, 2008.