

Deakin Research Online

This is the published version:

Lubans, David R., Morgan, Philip J., Cliff, Dylan P., Barnett, Lisa M. and Okely, Anthony D. 2010, Fundamental movement skills in children and adolescents: review of associated health benefits, *Sports medicine*, vol. 40, no. 12, pp. 1019-1035.

Available from Deakin Research Online:

<http://hdl.handle.net/10536/DRO/DU:30031139>

Reproduced with the kind permission of the copyright owner.

Copyright: 2010, ADIS Press Ltd.

Fundamental Movement Skills in Children and Adolescents

Review of Associated Health Benefits

David R. Lubans,¹ Philip J. Morgan,¹ Dylan P. Cliff,² Lisa M. Barnett³ and Anthony D. Okely²

1 School of Education, University of Newcastle, Callaghan Campus, Newcastle, New South Wales, Australia

2 Interdisciplinary Educational Research Institute, University of Wollongong, Wollongong, New South Wales, Australia

3 Centre for Physical Activity and Nutrition, Deakin University, Melbourne, Victoria, Australia

Contents

Abstract	1019
1. Methods	1021
1.1 Identification of Studies	1021
1.2 Criteria for Inclusion/Exclusion	1021
1.3 Criteria for Assessment of Study Quality	1021
1.4 Categorization of Variables and Level of Evidence	1022
2. Results	1022
2.1 Overview of Studies	1022
2.2 Overview of Study Quality	1022
2.3 Psychological Benefits	1023
2.4 Physiological Benefits	1030
2.5 Behavioural Benefits	1030
3. Discussion	1031
3.1 Overview of Findings	1031
3.2 Strengths and Limitations	1032
4. Conclusions	1032

Abstract

The mastery of fundamental movement skills (FMS) has been purported as contributing to children's physical, cognitive and social development and is thought to provide the foundation for an active lifestyle. Commonly developed in childhood and subsequently refined into context- and sport-specific skills, they include locomotor (e.g. running and hopping), manipulative or object control (e.g. catching and throwing) and stability (e.g. balancing and twisting) skills. The rationale for promoting the development of FMS in childhood relies on the existence of evidence on the current or future benefits associated with the acquisition of FMS proficiency. The objective of this systematic review was to examine the relationship between FMS competency and potential health benefits in children and adolescents. Benefits were defined in terms of psychological, physiological and behavioural outcomes that

can impact public health. A systematic search of six electronic databases (EMBASE, OVID MEDLINE, PsycINFO, PubMed, Scopus and Sport-Discus®) was conducted on 22 June 2009. Included studies were cross-sectional, longitudinal or experimental studies involving healthy children or adolescents (aged 3–18 years) that quantitatively analysed the relationship between FMS competency and potential benefits. The search identified 21 articles examining the relationship between FMS competency and eight potential benefits (i.e. global self-concept, perceived physical competence, cardio-respiratory fitness [CRF], muscular fitness, weight status, flexibility, physical activity and reduced sedentary behaviour). We found strong evidence for a positive association between FMS competency and physical activity in children and adolescents. There was also a positive relationship between FMS competency and CRF and an inverse association between FMS competency and weight status. Due to an inadequate number of studies, the relationship between FMS competency and the remaining benefits was classified as uncertain. More longitudinal and intervention research examining the relationship between FMS competency and potential psychological, physiological and behavioural outcomes in children and adolescents is recommended.

Fundamental movement skills (FMS) are considered to be the building blocks that lead to specialized movement sequences required for adequate participation in many organized and non-organized physical activities for children, adolescents and adults.^[1,2] Commonly developed in childhood and subsequently refined into context- and sport-specific skills,^[2-4] they include locomotor (e.g. running and hopping), manipulative or object control (e.g. catching and throwing) and stability (e.g. balancing and twisting) skills.^[1] The mastery of FMS has been purported as contributing to children's physical, cognitive and social development^[5] and is thought to provide the foundation for an active lifestyle.^[1,3] Recently, FMS competency has been proposed to interact with perceptions of motor competence and health-related fitness to predict physical activity and subsequent obesity from childhood to adulthood.^[3]

While children may naturally develop a rudimentary form of fundamental movement pattern, a mature form of FMS proficiency is more likely to be achieved with appropriate practice, encouragement, feedback and instruction.^[1,2] Children who do not receive adequate motor skill instructions and practice may demonstrate developmental delays in their gross motor ability.^[6] As such,

early childhood physical activity guidelines, such as the National Association for Sport and Physical Education's (NASPE) *Active Start*, indicate that the development of movement skills should be a key component of early childhood education programmes.^[7] Likewise, FMS competency is identified in National Standards as a primary goal of quality elementary school physical education in the US^[8] and represents an indicator of achievement for elementary school children in England's national physical education curriculum.^[9] Despite this focus, the prevalence of FMS mastery among children in some countries appears inadequately low.^[10,11] For example, in a recent US study of 9- to 12-year-old children, only half of the students assessed demonstrated proficiency in basketball throwing and dribbling motor tasks.^[11] Similarly, an Australian study^[12] involving students from years 4, 6, 8 and 10 (aged 9–15 years) found that the prevalence of mastery only exceeded 40% for one skill in one group (i.e. overarm throw, year 10 boys).

The rationale for promoting the development of FMS in childhood relies on the existence of evidence on the current or future benefits associated with the acquisition of FMS proficiency. Despite support for FMS promotion among motor

behaviourists^[3] and physical educators,^[13] the potential benefits of FMS competency have not yet been methodically evaluated. The purpose of this review is to systematically examine the potential psychological, physiological and behavioural public health benefits associated with FMS competency in children and adolescents.

1. Methods

1.1 Identification of Studies

The Quality of Reporting of Meta-analyses statement (QUOROM)^[14] was consulted and provided the structure for this review. A systematic search of six electronic databases (EMBASE, OVID MEDLINE, PsycINFO, PubMed, Scopus and SportDiscus[®]) was conducted from their year of inception to 22 June 2009. Individualized search strategies for the different databases included combinations of the following keywords: 'child', 'adolescent', 'youth', 'movement skill', 'motor skill', 'actual competence', 'object control', 'locomotor skill' and 'motor proficiency'. Only articles published or accepted for publication in refereed journals were considered for review. Conference proceedings and abstracts were not included. In the first stage of the research, titles and abstracts of identified articles were checked for relevance. In the second stage, full-text articles were retrieved and considered for inclusion. In the final stage, the reference lists of retrieved full-text articles were searched and additional articles known to the authors were assessed for possible inclusion. Eighteen expert informants in the area were also contacted to suggest or provide relevant manuscripts.

1.2 Criteria for Inclusion/Exclusion

Two authors (DRL and DPC) independently assessed the eligibility of the studies for inclusion according to the following criteria: (i) participants were aged 3–18 years (research articles that focused on youth from special populations were not included, e.g. overweight/obese, developmental coordination disorder); (ii) process (i.e. concerned with process or technique also known as qualitative) or product (i.e. concerned with out-

come) assessment of at least two FMS (e.g. run, vertical jump, horizontal jump, hop, dodge, leap, gallop, side gallop, skip, roll, throw, stationary dribble, catch, kick, two-handed strike, static balance); (iii) summary/subtest measure of FMS competency (e.g. locomotor or object control summary score) was used in analyses; (iv) quantitative assessment of potential health benefit of FMS competency (i.e. psychological, physiological or behavioural); (v) quantitative analysis of the relationship between FMS and potential benefits in any of the above domains; (vi) cross-sectional, longitudinal or experimental/quasi-experimental study design; and (vii) published in English. As this review focused on the potential benefits of FMS, which are gross motor skills,^[1] studies that used measurement batteries that included fine motor skills were excluded to preserve internal validity.

1.3 Criteria for Assessment of Study Quality

Two authors (DRL and PJM) independently assessed the quality of the studies that met the inclusion criteria. The criteria for assessing the quality of the studies were adapted from the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement^[15] and the Consolidated Standards of Reporting Trials (CONSORT) statement.^[16] A formal quality score for each study was completed on a 6-point scale by assigning a value of 0 (absent or inadequately described) or 1 (explicitly described and present) to each of the following questions listed: (i) Did the study describe the participant eligibility criteria? (ii) Were the participants randomly selected (or for experimental studies, was the process of randomization clearly described and adequately carried out)? (iii) Did the study report the sources and details of FMS assessment and did the instruments have acceptable reliability for the specific age group? (iv) Did the study report the sources and details of assessment of potential benefits and did all of the methods have acceptable reliability? (v) Did the study report a power calculation and was the study adequately powered to detect hypothesized relationships? (vi) Did the study report the numbers of individuals who completed each of the different measures

and did participants complete at least 80% of FMS and benefit measures? Studies that scored 0–2 were regarded as low quality studies, studies that scored 3–4 were classified as medium quality and those that scored 5–6 were classified as high quality.

1.4 Categorization of Variables and Level of Evidence

The benefits were categorized as follows: psychological (e.g. physical self-perception), physiological (e.g. fitness and healthy weight status) and behavioural (e.g. time spent in physical activity and sedentary behaviours). It should be noted that studies assessing the benefit of fitness in this review will be discussed in terms of whether they used product- or process-oriented motor skill assessments. This is because product-oriented motor skill assessments can view certain fitness constructs (such as strength and speed) as part of the motor skill assessment, unlike process-orientated assessments that are concerned with the quality or technique of the skill execution.

Results were coded using the methods first described by Sallis et al.^[17] and more recently by Hinkley et al.^[18] and Van der Horst et al.^[19] The relationship between FMS competency and each potential benefit was determined by examining the percentage of studies that reported a statistically significant relationship (i.e. between FMS competency and benefit) and is explained in table I. If only 0–33% of the included studies reported a relationship between FMS competency and the benefit, the result was categorized as no association (0). If 34–59% of the studies reported statistically significant relationships between FMS competency and the benefit, the result was categorized as inconsistent

or uncertain (?). If 60–100% of studies reported a positive relationship between FMS competency and the benefit, the result was coded as a positive association (+). The methods of Sallis et al.^[17] were modified to address the issue of study quality and additional coding was conducted based on studies assessed as high quality. If 60–100% of high quality studies (≥ 4) found a positive relationship between FMS competency and the benefit, the result was coded as having strong evidence for a positive association (++).

2. Results

2.1 Overview of Studies

A total of 1793 potentially relevant articles were identified using database searches (figure 1). Following feedback from international experts and checking the reference lists of included studies, a total of 21 articles satisfied the inclusion criteria and were included in the review (table II). The flow of studies through the review process and the reasons for exclusion are reported in figure 1. Of the included articles, 15 reported on cross-sectional studies, four on longitudinal studies and two on experimental studies. Nine studies were conducted in Australia, eight in the US, and one each in Canada, Scotland, Belgium and Germany. The number of study participants ranged from 29^[23] to 4363.^[40]

2.2 Overview of Study Quality

There was 96% agreement between authors on the study assessment criteria and full consensus was achieved after discussion. Results from the study quality assessment are reported in table III. Seven studies were identified as high

Table I. Rules for classifying the association between potential benefits and fundamental movement skills (FMS) competency

Studies supporting association (%)	Summary code	Explanation of code
0–33	0	No association
34–59	?	Inconsistent or uncertain ^a
60–100	–	Negative association
60–100	+	Positive association
60–100	++	Strong evidence for a positive association ^b

a The relationship between benefit and FMS competency was considered uncertain if <4 studies examined the relationship.

b Strong evidence for a positive association is identified when >60% of high quality studies (≥ 4 studies) reported a positive association.




Fig. 1. Flow of studies through the review process. **FMS** = fundamental movement skills.

quality,^[24,29,30,33,34,36,40] 13 studies were rated as medium quality^[11,20-23,25,26,28,32,35,37,39,41] and one study was classified as low quality.^[31] Most of the studies used valid and reliable measures of FMS assessment and also reported the reliability data from their potential benefits. None of the studies reported power calculations to determine if the studies were adequately powered to detect the hypothesized relationships.

2.3 Psychological Benefits

A summary of the associations between FMS competency and potential benefits is reported in table IV. Three studies examined the relationship

between perceived physical competence and FMS competency.^[21,29,34] Perceived competence was associated with at least one aspect of FMS competency in all three studies. Perceived competence refers to an individual's perception of their actual motor proficiency. In a 6-year longitudinal study, Barnett et al.^[34] found that object control competency in childhood was associated with perceived physical competence in adolescence. Only one study assessed the association between FMS competency and global self-concept.^[20] Martinek and colleagues^[20] examined the impact of a motor skill intervention on FMS and self-concept in a sample of 344 children. Although FMS and self-concept improved over the study period, the

Table II. Summary of included studies

Study	Sample; age; school grade; location	Type of study	Analyses	FMS measure ^a		Benefits assessed	Results
				PRODUCT	PROCESS		
Martinek et al. ^[20]	344 children; 6–10 y; NR; US	Experimental	ANCOVA and bivariate correlation	KTK: one-legged obstacle jumping, jumping from side to side as well as sideway movements		Global self-concept (Self Concept Scale for Children)	FMS and self-concept improved in the intervention group over the study period. However, the relationship between self-concept and FMS was nonsignificant at baseline and post-test
Rudisill et al. ^[21]	218 children; 9–11 y; 3, 4 and 5; US	Cross-sectional	Bivariate correlation	Locomotor (standing long jump, 50-yard dash and shuttle run) and object control (two ball throws short and long distance)		Perceived physical competence (Motor Perceived Competence Scale)	Locomotor and object control proficiency associated with perceived competency
Marshall and Bouffard ^[22]	200 children; NR; 1 and 4; Canada	Experimental	ANOVA and bivariate correlation		Test of gross motor development (run, gallop, hop, leap, horizontal jump, slide, skip, striking a stationary ball, stationary dribble, catch, kick, overhand throw and underhand roll), characterized into locomotor and object control subtests	CRF (multi-stage fitness test)	Object control and locomotor FMS competency associated with CRF
Reeves et al. ^[23]	29 children; 5–6 y; kindergarten; US	Cross-sectional	Bivariate correlation	Bruininks Oseretsky Test of Motor Proficiency: (i) running speed and agility; (ii) balance; and (iii) bilateral coordination subtests		CRF (half-mile walk/run)	CRF (half-mile walk/run) was positively associated with balance and bilateral coordination
Okely et al. ^[24]	2026 adolescents; 13–16 y; 8 and 10; Australia	Cross-sectional	Bivariate correlations and linear regression		FMS: A Manual for Classroom Teachers, (run, vertical jump, catch, overhand throw, kick, strike)	CRF (multi-stage fitness test)	FMS competency associated with CRF controlling for gender and grade at school

Continued next page

Table II. Contd

Study	Sample; age; school grade; location	Type of study	Analyses	FMS measure ^a		Benefits assessed	Results
				PRODUCT	PROCESS		
Okely et al. ^[25]	982 adolescents; 13–16 y; 8 and 10; Australia	Cross-sectional	Linear regression analysis (controlling for gender, grade, SES, geographic location)		FMS: A Manual for Classroom Teachers (run, vertical jump, catch, overhand throw, kick, strike)	PA (APARQ)	FMS associated with time in organized PA but not time in non-organized PA controlling for gender and school grade
McKenzie et al. ^[26]	207 children; 4–6 y; NR; US	Longitudinal	Bivariate correlation and linear regression	Lateral jump, catch, and one foot balance		PA (PAR 7-day questionnaire) and adiposity (skinfolds: triceps and subscapular)	Inverse association between adiposity and FMS in boys but not girls Jumping related to PA at age 12 for girls FMS at ages 4–6 did not predict PA at age 12
Okely et al. ^[27]	4363 children and adolescents; NR; 4, 6, 8 and 10; Australia	Cross-sectional	Logistic regression modelling and multiple linear regression		FMS: A Manual for Classroom Teachers (run, vertical jump, catch, overhand throw, kick, strike)	BMI z-score and waist circumference	FMS (locomotor) inversely associated with BMI z-score in children and adolescents
Graf et al. ^[28]	668 children; 6.7 ± 0.4 y; NR; Germany	Cross-sectional	ANCOVA (adjusted for age and gender) and bivariate correlation	KTK: balancing backwards, one-legged obstacle jumping, jumping from side to side as well as sideways movements		BMI z-score, time spent in organized PA (parent questionnaire) and watching TV (child questionnaire)	Inverse association between BMI and FMS Positive association between FMS and PA Nonsignificant association between FMS and TV watching
Southall et al. ^[29]	142 children; 10.8 y; 5 and 6; Australia	Cross-sectional	ANCOVA		Test of Gross Motor Development 2	BMI z-score, perceived physical competence (SPPC)	Overweight children had lower total FMS and locomotor FMS Overweight children had lower perceived physical competence scores No difference between overweight and normal weight children for object control skills
Fisher et al. ^[30]	394 children; 4.2 ± 0.5 y; NR; Scotland	Cross-sectional	Bivariate correlation		Movement Assessment Battery: 15 skills including jumps, balance, skips, ball exercises and throwing	PA (accelerometer)	FMS associated with total PA and MVPA

Continued next page

Table II. Contd

Study	Sample; age; school grade; location	Type of study	Analyses	FMS measure ^a		Benefits assessed	Results
				PRODUCT	PROCESS		
Hamstra-Wright et al. ^[31]	36 children; 8–9 y; NR; US	Cross-sectional	Linear stepwise multiple regression (controlling for gender and age) and bivariate correlation		Test of Gross Motor Development 2 (run, gallop, hop, leap, horizontal jump, slide, striking a stationary ball, stationary dribble, catch, kick, overhand throw and underhand roll), characterized into locomotor and object control subtests	PA (sport experience questionnaire)	Participation in organized and non-organized PA was associated with locomotor competency
Castelli and Valley ^[32]	230 children; 9.5 ± 1.6 y; NR; US	Cross-sectional	Bivariate correlation		SCPEAP: scoring and protocols including: basketball dribble and pass, paddle bat hit and overhand ball throwing to provide a summative score for FMS competency	PA (parent and child 7-d questionnaire, pedometer), BMI z-score, flexibility, CRF (PACER), muscular endurance (curl-ups and push-ups) and flexibility (sit and reach)	FMS competency associated with CRF, muscular endurance, flexibility and PA No relationship between FMS and BMI z-score
Barnett et al. ^[33]	928 children, 244 adolescents (follow-up); 16.4 y; 10 and 11; Australia	Longitudinal (6-y follow-up)	General linear regression model controlling for gender		Get Skilled, Get Active: object control (kick, catch, overhand throw) locomotor (hop, side gallop, vertical jump)	CRF (multi-stage fitness test)	Childhood object control proficiency associated with CRF in adolescence
Barnett et al. ^[34]	928 children, 250 adolescents (PA model); 227 adolescents	Longitudinal (6-y follow-up)	Bivariate correlation and structural equation modelling to test		Get Skilled, Get Active: object control (kick, catch, overhand throw)	APARQ, CRF (multi-stage fitness test) and perceived physical	Childhood object control was associated with adolescent perceived sports competence Childhood object control

Continued next page

Table II. Contd

Study	Sample; age; school grade; location	Type of study	Analyses	FMS measure ^a		Benefits assessed	Results
				PRODUCT	PROCESS		
	(fitness model); 16.4 y; NR; Australia		for mediators		locomotor (hop, side gallop, vertical jump)	competence (PSPP)	associated with adolescent PA Locomotor competency was associated with perceived competence in girls only Locomotor competency was not associated with PA in either girls or boys Locomotor competency was associated with CRF in girls only
Erwin and Castelli ^[11]	180 children; 10.5 ± 0.8 y; 4 and 5; US	Cross- sectional	Bivariate correlation		SCPEAP scoring and protocols including: basketball dribble and pass, overhand ball throwing and gymnastic movement and balance	PA (ACTIVITYGRAM questionnaire), physical fitness (CRF, strength, endurance, flexibility and BMI z-score)	FMS competency associated with PA and physical fitness
Hume et al. ^[35]	248 children; 9–12 y; NR; Australia	Cross- sectional	Linear regression and bivariate correlation		FMS: A Manual for Classroom Teachers, object control (overhand throw, two handed strike, kick) and locomotor (sprint run, dodge and vertical jump)	PA (accelerometer) and BMI z-scores	MPA, VPA and MVPA associated with FMS proficiency in boys VPA associated with FMS proficiency in girls BMI z-scores not associated with FMS in boys or girls
Williams et al. ^[36]	198 children; 3–4 y; NR; US	Cross- sectional	Bivariate correlation		Children's Activity and Movement in preschool study motor skill protocol: locomotor (run, jump, slide, gallop, leap and hop) and object control (throw, roll, kick, catch, strike and dribble)	BMI z-score and PA (accelerometry)	Object control and locomotor proficiency associated with PA in 4-y-olds, but not 3-y-olds BMI z-score not associated with object control or locomotor proficiency in 3- or 4-y-olds

Continued next page

Table II. Contd

Study	Sample; age; school grade; location	Type of study	Analyses	FMS measure ^a		Benefits assessed	Results
				PRODUCT	PROCESS		
Barnett et al. ^[37]	928 children, 276 adolescents (follow-up); 16.4 y; NR; Australia	Longitudinal (6-y follow-up)	General linear model controlling for grade and gender, general linear model controlling for grade and logistic regression		Get Skilled, Get Active: object control (kick, catch, overhand throw) locomotor (hop, side gallop, vertical jump)	PA (APARQ)	Object control proficiency in childhood associated with time in MVPA and time in organized PA Object control proficiency in childhood was associated with probability of participating in VPA but not associated with probability of participating in organized PA Locomotor proficiency did not predict time in or probability of participating in any form of adolescent PA
D'Hondt et al. ^[38]	117 children; 5–10 y; NR; Belgium	Cross-sectional	ANOVA and bivariate correlation		Movement Assessment Battery for Children: ball skills, static and dynamic balance	BMI z-score and PA (accelerometers)	FMS competency (ball skills and balance) was higher in normal and overweight compared with obese children FMS competency (ball skills and balance) associated with PA
Cliff et al. ^[39]	46 children; 4.3 ± 0.7 y; preschool; Australia	Cross-sectional	Bivariate correlation and linear regression		Test of Gross Motor Development 2 (run, gallop, hop, leap, horizontal jump, slide, striking a stationary ball, stationary dribble, catch, kick, overhand throw and underhand roll), characterized into locomotor and object control subtests	PA and sedentary behaviour (accelerometer)	Object control proficiency was associated with moderate PA in boys Locomotor proficiency was not significantly associated with PA in boys Locomotor proficiency and overall FMS proficiency were negatively associated with PA in girls Object control proficiency was not associated with PA in girls FMS not associated with sedentary behaviour in boys or girls

a PRODUCT or PROCESS measure of FMS competency.

ANCOVA = analysis of covariance; **APARQ** = Adolescent Physical Activity Questionnaire; **BMI** = body mass index; **CRF** = cardio-respiratory fitness; **FMS** = fundamental movement skills; **KTK** = Körper Koordinations Test für Kinder; **MVPA** = moderate to vigorous physical activity; **NR** = not reported; **PA** = physical activity; **PACER** = Progressive Aerobic Cardiovascular Endurance Run; **PAR** = physical activity recall questionnaire; **PROCESS** = process assessment of FMS concerned with technique; **PRODUCT** = product assessment of FMS concerned with outcome; **PSPP** = Physical Self-Perception Profile; **SCPEAP** = South Carolina Physical Education Assessment; **SES** = socio-economic status; **SPPC** = Self-Perception Profile for Children; **TV** = television; **VPA** = vigorous physical activity.

Table III. Fundamental movement skills (FMS) study quality checklist with quality scores assigned

Study	Did the study describe the participant eligibility criteria?	Were the participants randomly selected? ^a	Did the study report the sources and details of FMS assessment and did the instruments have acceptable reliability for the specific age group?	Did the study report the sources and details of assessment of potential benefits and did all of the methods have acceptable reliability for the specific age group?	Did the study report a power calculation and was the study adequately powered to detect hypothesized relationships?	Did the study report the numbers of individuals who completed each of the different measures and did participants complete at least 80% of FMS and benefit measures?	Quality score total/6
Martinek et al. ^[20]	1	0	1	1	0	0	3
Rudisill et al. ^[21]	1	0	0	1	0	1	3
Marshall and Bouffard ^[22]	1	0	1	1	0	1	4
Reeves et al. ^[23]	1	0	1	1	0	1	4
Okely et al. ^[24]	1	1	1	1	0	1	5
Okely et al. ^[25]	1	1	1	0	0	0	3
McKenzie et al. ^[26]	1	0	1	0	0	1	3
Okely et al. ^[27]	1	1	1	1	0	1	5
Graf et al. ^[28]	1	1	1	0	0	0	3
Southall et al. ^[29]	1	1	1	1	0	1	5
Fisher et al. ^[30]	1	1	1	1	0	1	5
Hamstra-Wright et al. ^[31]	0	0	1	0	0	0	1
Castelli et al. ^[32]	1	0	1	1	0	0	3
Barnett et al. ^[33]	1	1	1	1	0	1	5
Barnett et al. ^[34]	1	1	1	1	0	1	5
Erwin and Castellji ^[11]	1	0	1	0	0	1	3

Continued next page

Table III. Contd

Study	Did the study describe the participant eligibility criteria?	Were the participants randomly selected? ^a	Did the study report the sources and details of FMS assessment and did the instruments have acceptable reliability for the specific age group?	Did the study report the sources and details of assessment of potential benefits and did all of the methods have acceptable reliability for the specific age group?	Did the study report a power calculation and was the study adequately powered to detect hypothesized relationships?	Did the study report the numbers of individuals who completed each of the different measures and did participants complete at least 80% of FMS and benefit measures?	Quality score total/6
Hume et al. ^[95]	1	0	1	1	0	1	4
Williams et al. ^[96]	1	1	1	1	0	1	5
Barnett et al. ^[97]	1	1	1	1	0	0	4
D'Hondt et al. ^[98]	1	0	1	1	0	1	4
Cliff et al. ^[99]	1	1	1	1	0	0	4

a For intervention studies the criterion was as follows: were participants randomly allocated and was the process of randomization clearly described and adequately carried out (envelope or algorithm)?

relationship between self-concept and FMS was nonsignificant at baseline and post-test.^[20]

2.4 Physiological Benefits

Weight status was the most commonly assessed physiological benefit of FMS competency and was included in nine studies. Body composition was generally estimated using body mass index (BMI) z-score; however, skinfolds were used in one study.^[26] Six of the nine studies found an inverse association between FMS competency and BMI z-score^[11,26-29,39,41] and three studies found no association between FMS competency and weight status.^[32,35,36]

Four studies examined the relationship between FMS competency and CRF. All four found a positive relationship between skill ability and fitness level.^[22-24,33] Three of these studies used a process-oriented motor skill assessment^[22,24,33] and one used a product assessment.^[23] One study found positive associations between FMS competency, muscular fitness and flexibility.^[32] Another study found a positive relationship between FMS competency and a composite physical fitness score (which included CRF, strength, endurance, flexibility and BMI).^[11]

2.5 Behavioural Benefits

Thirteen studies examined the relationship between FMS competency and participation in physical activity. Eight studies used self-report measures of physical activity, four studies used objective measures of physical activity (i.e. accelerometers) and one study used both self-report and pedometers. FMS competency was found to be associated with at least one component of physical activity (e.g. non-organized activity, organized activity, pedometer step counts) in 11 of the cross-sectional studies^[11,25,28,30-32,34-36,39,41] and one of the longitudinal studies.^[37] Longitudinally, McKenzie et al.^[26] found that FMS competency at ages 4–6 years did not predict physical activity at age 12 years. Both studies that examined the association between sedentary behaviour and FMS competency in children^[28,39] did not find a statistically significant relationship.

Table IV. Summary of studies examining the relationship between potential benefits and fundamental movement skill (FMS) competency in youth

Benefits	Associated with FMS		Not associated with FMS references	Summary coding ^a	
	references	association (-/+) ^b		n/N for benefit (%) ^c	association (-/+) ^b
Psychological benefits					
Global self-concept			20	1/1 (100)	?
Perceived physical competence	21, 29, 34 ^d	+		3/3 (100)	?
Physiological benefits					
Weight status (BMI z-score, BMI, skinfolds)	26-29 ^e , 41	-	32, 35, 36	5/8 (63)	-
CRF	22-24, 33 ^f	+		4/4 (100)	+
Muscular fitness	32	+		1/1 (100)	?
Flexibility	32	+		1/1 (100)	?
Physical fitness ^g	11	+		1/1 (100)	?
Behavioural benefits					
Physical activity	11, 25, 28, 30-32, 35-37, 39 ^e , 41, 34 ^h	+	26 ⁱ	11/13 (80)	++
Sedentary behaviour			28, 39	2/2 (100)	?

a Summary code provides an overall summary of the findings for each benefit.

b Association shows the direction of the individual and summary association. A positive or negative association was noted if at least one component of FMS competency was associated with the hypothesized benefit.

c n = number of studies that report support for relationship, N = number of studies that examined and reported possible associations between FMS competency and potential benefit.

d Childhood FMS competency associated with adolescent perceived competence.

e Positive association for boys and negative association for girls.

f Childhood FMS competency associated with CRF in adolescence.

g Composite physical fitness score including CRF, flexibility, strength, muscular fitness and BMI.

h Childhood FMS competency associated with physical activity in adolescence.

i FMS competency at ages 4-6 y did not predict physical activity at age 12 y.

BMI = body mass index; **CRF** = cardio-respiratory fitness; + indicates positive association; ++ indicates strong evidence for a positive association; - indicates negative association; ? indicates inconsistent or uncertain.

3. Discussion

3.1 Overview of Findings

The aim of this systematic review was to identify the health benefits associated with FMS competency in children and adolescents. We found 21 articles that assessed eight potential benefits (i.e. self-concept, perceived physical competence, CRF, muscular fitness, weight status, flexibility, physical activity and sedentary behaviour). We found strong evidence from cross-sectional studies for a positive association between FMS competency and physical activity in children and adolescents. There was also a positive association between FMS competency and CRF, and an inverse association between

FMS competency and weight status. Due to an inadequate number of studies, the relationship between FMS competency and global self-concept, perceived physical competence, muscular fitness, flexibility and sedentary behaviour were classified as uncertain.

It has been suggested that proficiency in a range of FMS provides the foundation for an active lifestyle.^[1,3] The results from this review confirm the cross-sectional relationship between FMS competency and physical activity in children and adolescents. A number of large-scale cross-sectional studies,^[25,30] some of which used objective measures of physical activity,^[30,36] found positive associations between FMS competency and participation in physical activity. One longitudinal study found an association

between childhood object control skill ability and adolescent physical activity.^[34,37] The other longitudinal study in this review found no association between FMS proficiency and physical activity.^[26] This study examined early childhood (ages 4–6 years), three motor skills (lateral jumping, catching a ball, and balancing on one foot) and early adolescent (12 years) physical activity participation (measured via the Seven-Day Physical Activity Recall questionnaire).^[26] However, the study was limited by the use of a physical activity self-report measure and the assessment of only three FMS. Furthermore, two of these skills included what the authors termed ‘a restricted range of measurement’; 0–2 for balancing and 0–6 for catching.^[26] This notion that a more comprehensive skill battery might be needed to accurately test whether skill is associated with physical activity is substantiated by the positive associations found in this review; all the other studies that found positive associations between motor skill and physical activity assessed more than three motor skills.

The other factor that may have precluded the longitudinal study by McKenzie et al.^[26] finding no association, was that skills were measured before the children had been provided with an opportunity to participate in school physical education (PE) and in out-of-school PE and sport programmes. It has been proposed that the relationship between skill ability and physical activity may strengthen over time.^[42] This theory may also be supported in this review, as the one cross-sectional study in which the relationship between physical activity and motor skill ability was most uncertain (both positive and negative associations) was in preschool children.^[39] However, this study may simply be limited by a small sample size, as the other two studies in this age group found positive associations.^[11,43]

We also found a positive association between FMS competency and CRF, and an inverse association between FMS competency and weight status. It has also been suggested that FMS competency might influence fitness levels, as activities that involve FMS also demand high levels of muscular and cardiorespiratory fitness.^[42] More skillful children may increase their time in

physical activity and persist with activities that require high levels of physical fitness,^[42] providing the opportunity for fitness adaptations through progressive overload. Increased time in higher intensity physical activity will contribute to higher levels of CRF and improvements in body composition.^[44]

3.2 Strengths and Limitations

This is the first systematic review of studies examining the relationship between FMS competency and potential health benefits in children and adolescents. The QUOROM statement was consulted and provided the structure for this review, which included an assessment of study quality using criteria adapted from the CONSORT and STROBE statements. However, there are a number of issues that should be noted. First, we did not include studies that combined gross motor skills and fine motor skills in the same composite score. For example, Wrotniak and colleagues^[45] examined the relationship between motor competency and physical activity using the Bruininks-Oseretsky Test of Motor Proficiency (BOTMP) and found a positive association. While the BOTMP is an established measure of general motor ability, the current review was limited to FMS competency and therefore the inclusion of fine motor skills was beyond the scope of this review. It should also be noted that we excluded studies that did not provide a composite FMS score. A number of studies examined the relationship between individual FMS tests and potential benefits but did not provide a summary score.^[46–48] Finally, due to the relatively small number of studies and the inclusion of longitudinal studies, the results for children and adolescents have been combined. As a result, this review could not assess whether the importance of FMS competency varies between childhood and adolescence,^[42] a hypothesis that requires further investigation.

4. Conclusions

Our review included only two longitudinal and two experimental studies. More longitudinal

studies exploring the relationship between changes in FMS competency and potential benefits over time are needed to investigate the causal nature of such relationships. It has been hypothesized that children with high motor skill proficiency will have higher levels of fitness and perceived sports competence, which in turn predict greater participation in physical activity, and vice versa.^[42] This proposed reciprocal relationship could also be investigated in future studies.

In the current review we did not include intervention studies that did not directly examine the relationship between FMS competency and potential benefits. For example, two previous high quality obesity prevention trials^[49,50] evaluated the impact of treatment on changes in FMS competency and BMI z-score in children, but did not report the relationship between such changes. Future physical activity and obesity prevention studies should conduct mediation analyses to identify if FMS competency mediates the impact of interventions on primary outcomes (e.g. BMI z-score, fitness). Few studies have conducted mediation analyses in physical activity interventions among youth^[51] and the importance of FMS competency to future physical activity and other outcomes will be reinforced through this type of analysis. The one study reviewed that did conduct a mediation analysis,^[34] found that perceived sports competence acted as a mediator between skill ability and physical activity.

Due to the limited number of studies it was not feasible to examine how the association between motor skill ability and potential benefits might differ according to gender. Gender differences in motor proficiency have been found, with males generally more proficient than females in object control skill performance.^[35,43,52,53] In locomotor skill performance, some studies report no gender differences,^[35,53,54] while others report males^[55] or females^[53] as more proficient. The potential impact of these differences is important to investigate.

Our findings suggest that FMS development should be included in school- and community-based interventions. Teaching children to become competent and confident performers of FMS may lead to a greater willingness to participate in physical activities that may also provide opportu-

nities to improve fitness levels and reduce the risk of unhealthy weight gain. It is important that such skills are taught during preschool and elementary school years as children are at an optimal age in terms of motor skill learning^[1] and motor skill proficiency tracks through childhood.^[56] In addition, improving the FMS competency of girls should be a priority as many girls lack basic skill proficiency.^[10,11] Existing school physical education programmes have been criticized for not providing a learning environment to develop FMS,^[57] so training and resources should be prioritized to ensure children receive quality instruction in FMS.

FMS have been hypothesized as important to children and adolescents' physical, social and psychological development,^[1,2] and may be the foundation of an active lifestyle. This review has provided evidence supporting the positive association between FMS competency in children and adolescents and physical activity. Furthermore, the positive association between FMS competency and CRF and the inverse relationship between FMS proficiency and weight status suggest that developing competency in movement skills may have important health implications for young people.

Acknowledgements

No external funding was used for this project. The authors would like to thank Emily Hoffman and Kelly Magrann for their assistance in the retrieval of journal articles. The authors have no conflicts of interest that are directly relevant to the content of this review.

References

1. Gallahue DL, Ozmun JC. Understanding motor development: infants, children, adolescents, adults. 6th ed. Boston (MA): McGraw-Hill, 2006
2. Clark JE, Metcalfe JS. The mountain of motor development. In: Clark JE, Humphrey JH, editors. Motor development: research and reviews. Vol. 2. Reston (VA): National Association of Sport & Physical Education, 2002: 163-90
3. Stodden D, Goodway JD, Langendorfer S, et al. A developmental perspective on the role of motor skill competence in physical activity: an emergent relationship. *Quest* 2008; 60: 290-306
4. Clark JE. From the beginning: a developmental perspective on movement and mobility. *Quest* 2005; 57: 37-45

5. Payne VG, Isaacs LD. Human motor development: a life-span approach. 3rd ed. Mountain View (CA): Mayfield, 1995
6. Goodway JD, Branta CF. Influence of a motor skill intervention on fundamental motor skill development of disadvantaged preschool children. *Res Q Exerc Sport* 2003; 74: 36-46
7. National Association for Sport and Physical Education. Active Start: a statement of physical activity guidelines for children birth to five years. Reston (VA): NASPE Publications, 2009
8. National Association for Sport and Physical Education. Moving into the future: national standards for physical education. Reston (VA): McGraw-Hill, 2004
9. Department for Education and Employment. The National Curriculum for England: physical education. London: Crown/Qualifications and Curriculum Authority, 1999
10. Okely AD, Booth ML. Mastery of fundamental movement skills among children in New South Wales: prevalence and sociodemographic distribution. *J Sci Med Sport* 2004; 7 (3): 358-72
11. Erwin HE, Castelli DM. National physical education standards: a summary of student performance and its correlates. *Res Q Exerc Sport* 2008; 79 (4): 495-505
12. Booth M, Okely AD, McLellan L, et al. Mastery of fundamental motor skills among New South Wales school students: prevalence and sociodemographic distribution. *J Sci Med Sport* 1999; 2 (2): 93-105
13. Pangrazi RP. Dynamic physical education for elementary school children. 14th ed. San Francisco (CA): Pearson Education, 2004
14. Moher D, Cook DJ, Eastwood S, et al. Improving the quality of reports of meta-analyses of randomised controlled trials: the QUOROM statement. *Lancet* 1999; 354 (27): 1896-900
15. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Prev Med* 2007; 370 (9596): 1453-7
16. Moher D, Schulz KF, Altman DG. The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials. *Ann Intern Med* 2001; 134: 657-62
17. Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc* 2000; 32 (5): 963-75
18. Hinkley T, Crawford D, Salmon J, et al. Preschool children and physical activity: a review of correlates. *Am J Prev Med* 2008; 34 (5): 435-41
19. Van der Horst K, Paw MJCA, Twisk JWR, et al. A brief review on correlates of physical activity and sedentariness in youth. *Med Sci Sports Exerc* 2007; 39 (8): 1241-50
20. Martinek T, Cheffers J, Zaichkowsky L. Physical activity, motor development and self-concept: race and age differences. *Percept Mot Skills* 1978; 46: 147-54
21. Rudisill ME, Mahar MT, Meaney KS. The relationship between children's perceived and actual motor competence. *Percept Mot Skills* 1993; 76 (3): 895-906
22. Marshall J, Bouffard M. The effects of quality daily physical education on movement competency in obese versus non-obese children. *Adapt Phys Act Q* 1997; 14: 222-37
23. Reeves L, Broeder CE, Kennedy-Honeycutt L, et al. Relationship of fitness and gross motor skills for five-to-six year-old children. *Percept Mot Skills* 1999; 89: 739-47
24. Okely A, Booth ML, Patterson JW. Relationship of cardio-respiratory endurance to fundamental movement skill proficiency among adolescents. *Pediatr Exerc Sci* 2001; 13 (4): 380-91
25. Okely A, Booth ML, Patterson JW. Relationship of physical activity to fundamental movement skills among adolescents. *Med Sci Sports Exerc* 2001; 33 (11): 1899-904
26. McKenzie T, Sallis J, Broyles S, et al. Childhood movement skills: predictors of physical activity in Anglo American and Mexican American adolescents? *Res Q Exerc Sport* 2002; 73 (3): 238-44
27. Okely AD, Booth ML, Chey T. Relationships between body composition and fundamental movement skills among children and adolescents. *Res Q Exerc Sport* 2004; 75 (3): 238-47
28. Graf C, Koch B, Kretschmann-Kandel E, et al. Correlation between BMI, leisure habits and motor abilities in childhood (CHILT-Project). *Int J Obes* 2004; 28: 22-6
29. Southall J, Okely A, Steele J. Actual and perceived physical competence in overweight and non-overweight children. *Pediatr Exerc Sci* 2004; 16: 15-24
30. Fisher A, Reilly JJ, Kelly LA, et al. Fundamental movement skills and habitual physical activity in young children. *Med Sci Sports Exerc* 2005; 37 (4): 684-8
31. Hamstra-Wright K, Swanik B, Sitler M, et al. Gender comparisons of dynamic restraint and motor skill in children. *Clin J Sports Med* 2006; 16 (1): 56-62
32. Castelli D, Valley J. The relationship of physical fitness and motor competence to physical activity. *J Teach Phys Educ* 2007; 26: 358-74
33. Barnett LM, van Beurden E, Morgan PJ, et al. Does childhood motor skill proficiency predict adolescent fitness? *Med Sci Sports Exerc* 2008; 40 (12): 2137-44
34. Barnett LM, Morgan PJ, van Beurden E, Beard JR. Perceived sports competence mediates the relationship between childhood motor skill proficiency and adolescent physical activity and fitness: a longitudinal assessment. *Int J Behav Nutr Phys Act*. Epub 2008 Aug 8
35. Hume C, Okely A, Bagley S, et al. Does weight status influence associations between children's fundamental movement skills and physical activity? *Res Q Exerc Sport* 2008; 79 (2): 158-65
36. Williams HG, Pfeiffer KA, O'Neill JR, et al. Motor skill performance and physical activity in preschool children. *Obesity* 2008; 16: 1421-6
37. Barnett LM, van Beurden E, Morgan PJ, et al. Childhood motor skill proficiency as a predictor of adolescent physical activity. *J Adolesc Health* 2009; 44 (3): 252-9
38. D'Hondt E, Deforche B, De Bourdeaudhuij I, et al. Relationship between motor skill and body mass index in 5- to 10-year-old children. *Adapt Phys Act Q* 2009; 26 (1): 21-37
39. Cliff DP, Okely AD, Smith LM, et al. Relationships between fundamental movement skills and objectively measured

- physical activity in preschool children. *Pediatr Exerc Sci* 2009; 21: 436-49
40. Okely A, Booth M, Chey T. Relationships between body composition and fundamental movement skills among children and adolescents. *Res Q Exerc Sport* 2004; 75 (3): 238-47
 41. D'Hondt E, Deforche B, De Bourdeaudhuij I, et al. Relationship between motor skill and body mass index in 5 to 10 year old children. *Adapt Phys Act Q* 2009; 26: 21-37
 42. Stodden D, Langendorfer S, Robertson MA. The association between motor skill competence and physical fitness in young adults. *Res Q Exerc Sport* 2009; 80 (2): 223-9
 43. Runion BP, Robertson MA, Langendorfer SJ. Forceful overarm throwing: a comparison of two cohorts measured 20 years apart. *Res Q Exerc Sport* 2003; 74 (3): 324-30
 44. American College of Sports Medicine. Physical fitness in children and youth. *Med Sci Sports Exerc* 1988; 20: 422-3
 45. Wrotniak BH, Epstein LH, Dorn JM, et al. The relationship between motor proficiency and physical activity in children. *Pediatr* 2006; 118 (6): e1758-65
 46. Saakslahti A, Numminen P, Niinikoski H, et al. Is physical activity related to body size, fundamental motor skills and CHD risk factors in early childhood? *Pediatr Exerc Sci* 1999; 11: 327-40
 47. Raudsepp L, Liblik R. Relationship of perceived and actual motor competence in children. *Percept Mot Skills* 2002; 94: 1059-70
 48. Raudsepp L, Pall P. The relationship between fundamental motor skills and outside school physical activity of elementary school children. *Pediatr Exerc Sci* 2006; 18: 426-35
 49. Reilly J, Kelly L, Montgomery C, et al. Physical activity to prevent obesity in young children: cluster randomised controlled trial. *BMJ* 2006; 333: 1041-6
 50. Salmon J, Ball K, Hume C, et al. Outcomes of a group-randomized trial to prevent excess weight gain, reduce screen behaviors and promote physical activity in 10-year-old children: Switch-Play. *Int J Obes* 2008; 32: 601-12
 51. Lubans DR, Foster C, Biddle SJH. A review of mediators of behavior in interventions to promote physical activity among children and adolescents. *Prev Med* 2008; 47: 463-70
 52. Raudsepp L, Paasuke M. Gender differences in fundamental movement patterns, motor performances and strength measurements of prepubertal children. *Pediatr Exerc Sci* 1995; 7: 294-304
 53. van Beurden E, Barnett LM, Zask A, et al. Can we skill and activate children through primary school physical education lessons? 'Move it Groove it': a collaborative health promotion intervention. *Prev Med* 2003; 36 (4): 493-501
 54. Goodway J, Crowe H, Ward P. Effects of motor skill instruction on fundamental motor skill development. *Adapt Phys Act Q* 2003; 20: 298-314
 55. Haubenstricker J, Wisner D, Seefeldt V, et al. Gender differences and mixed-longitudinal norms on selected motor skills for children and youth [abstract]. *J Sport Exerc Psych* 1997; 19: S63
 56. Branta C, Haudenstricker J, Seefeldt V. Age changes in motor skills during childhood and adolescence. *Exerc Sport Sci Rev* 1984; 12: 467-520
 57. Morgan PJ, Hansen V. Classroom teachers' perceptions of the impact of barriers to teaching PE on the quality of PE programs delivered in primary schools. *Res Q Exerc Sport* 2008; 79: 506-16

Correspondence: Dr *David Lubans*, University of Newcastle, School of Education, Callaghan Campus, University Drive, NSW 2308, Australia.

E-mail: David.Lubans@newcastle.edu.au