

Deakin Research Online

Deakin University's institutional research repository

This is the published version (version of record) of:

Fitzsimons, James A. 2011, Southward range expansion of the Mourning Gecko *Lepidodactylus lugubris* on mainland Australia and nearshore islands, *Australian zoologist*, vol. 35, no. 3, pp. 619-621.

Available from Deakin Research Online:

<http://hdl.handle.net/10536/DRO/DU:30037003>

Reproduced with the kind permission of the copyright owner.

Copyright : 2011, Royal Zoological Society of New South Wales

Southward range expansion of the Mourning Gecko *Lepidodactylus lugubris* on mainland Australia and nearshore islands

James A. Fitzsimons^{1,2}

¹ School of Life and Environmental Sciences, Deakin University, 221 Burwood Highway, Burwood VIC 3125, Australia. Email: james.fitzsimons@deakin.edu.au

² The Nature Conservancy, Suite 3-04, 60 Leicester Street, Carlton VIC 3053, Australia

Key words: Range expansion, invasive species, Mourning Gecko, *Lepidodactylus lugubris*, Queensland

Introduction

The Mourning Gecko *Lepidodactylus lugubris* is a species that colonises areas of human habitation and its spread is assisted by humans (Lever 2003). The species continues to colonise many tropical and subtropical parts of the world (e.g. Yamashiro *et al.* 2000; Bauer *et al.* 2007; Kraus 2009).

In Australia, the mainland range for *L. lugubris* is described variously as 'coastal areas in eastern Queensland, from islands of Torres Strait to Townsville area' (Wilson and Swan 2008), 'several coastal localities between Innisfail and Port Douglas and from many islands in the Torres Strait' (Cogger 2000), 'Mossman to Innisfail' (Lever 2003) and 'Torres Strait to Bowen' (Couper *et al.* 2000). The species has more recently colonised Heron Island, near Gladstone (Limpus *et al.* 1999; Messenger 2005) and Darwin in the Northern Territory (Scrymgour 2006; McKay and Horner 2007; Wilson and Swan 2008) (Figure 1). It is thought the populations in the Torres Strait are indigenous while those on mainland Australia are introduced (H. Cogger pers. comm. 1996, in Lever 2003). Bomford (2008) considered the establishment risk rank of this species to be 'serious' in Australia.

Observations and Discussion

A young *L. lugubris* was observed and photographed at the resort at South Molle Island (20°15'46"S, 148°50'14"E), off Airlie Beach at 0750 h on 17 August 2008. It was inactive and resting in the open on paving. No other geckos were observed, nor searched for on South Molle Island at the time. The identity of the species was verified by Patrick Couper, Queensland Museum at Brisbane, Australia.

Searches of the Queensland Parks and Wildlife Service's (QPWS) Wildlife Online database in September 2008 revealed no records for this species on Molle Islands National Park (of which South Molle Island is a part) or adjacent island and coastal range national parks and state forests. The most southerly mainland record in both QPWS and Queensland Museum databases (as at September 2008) were from Bowen (20° 01' S, 148°

14' E), some 68 km to the northwest of South Molle Island (Figure 1). The Bowen population was recorded only recently (collections made in 1998 by Queensland Museum), as were records in two distinct localities around Townsville (in 1998 and 1999), some 240 km to the northwest of South Molle Island.

Interestingly, Limpus *et al.* (1999) noted that despite extensive searches for *L. lugubris* in coastal areas and islands between Mission Beach and Townsville in the mid-late 1970's, no geckos were located. This, and the more northerly and restricted mainland ranges previously published for *L. lugubris* in field guides, monographs and journal articles, suggests a recent progressive southerly range expansion on the mainland and nearshore islands. In 1991, the Queensland Museum indicated only one locality for this species on the Queensland coast (near Port Douglas) besides those in the Torres Strait (Covacevich and Couper 1991).

Limpus *et al.* (1999) considered the colonisation of Heron Island by *L. lugubris* to have been more likely by geckos from nearby Pacific Islands to the east rather than from mainland Australia, as it was some ~850 km south of the nearest known mainland record at the time (i.e. Mission Beach). In the case of South Molle Island, the gecko(s) are more likely to have originated from the coastal mainland to the north, as South Molle is less than 3 km from the coast with regular tourist interchange. Although it was not ascertained that the species was established on South Molle Island, its parthenogenetic nature (e.g. Harfmann Short and Petren 2008) suggests its presence is likely to lead to colonisation. The record on South Molle Island suggests the Mourning Gecko is likely to colonise other Whitsunday Islands and adjoining coastal settlements (e.g. Airlie Beach, Shute Harbour), if it has not already done so. Further surveys and research into the occurrence of this species on mainland Australia and nearshore islands (particularly those with tourist resorts) would assist in documenting the rate and likely extent of colonisation in Australia. This may be important for conservation, as the Mourning Gecko has the potential to compete with native Australian geckos (e.g. Cogger 2006).

Acknowledgements

Patrick Couper identified the species and provided information, Mark Antos, Janelle Thomas and an anonymous referee commented on a draft, while Andrew Amey and Andy Baker provided distribution records of the species from the Queensland Museum

and Queensland Parks and Wildlife Service databases, respectively. The South Molle Island record has been added to WildNet, the wildlife database of the now Queensland Department of Environment and Resource Management.




Figure 1. Locations of Mourning Gecko records from Queensland Museum and Queensland Parks and Wildlife Service databases, for range extremes and locations mentioned in published literature (see text plus Kluge 1963; Cameron and Cogger 1992; Turner and Green 1996; Woinarski *et al.* 1999; Sprackland *et al.* 2004), and the location of the South Molle Island record as described in this paper. The Cocos (Keeling) Islands, where a population also occurs (Cogger *et al.* 1983), is not shown.

References

- Bauer, A.M., Jackman, T.R., Greenbaum, E. and Papenfuss, T.J. 2007. First record of *Lepidodactylus lugubris* in Suriname. *Applied Herpetology* 4: 84-85.
- Bomford, M. 2008. *Risk Assessment Models for Establishment of Exotic Vertebrates in Australia and New Zealand*. Invasive Animals Cooperative Research Centre, Canberra.
- Cameron, E.E. and Cogger, H.G. 1992. *The Herpetofauna of the Weipa Region, Cape York Peninsula*. Technical Reports of the Australian Museum No.7. Australian Museum, Sydney.
- Cogger, H.G. 2000. *Reptiles and Amphibians of Australia*. 6th ed. Reed New Holland, Sydney.
- Cogger, H. 2006. National Recovery Plan for Lister's Gecko *Lepidodactylus listeri* and the Christmas Island Blind Snake *Typhlops exocoeti*. Department of the Environment and Heritage, Canberra.
- Cogger, H., Sadlier, R. and Cameron, E. 1983. *The Terrestrial Reptiles of Australia's Island Territories*. Special Publication 11. Australian National Parks and Wildlife Service, Canberra.
- Couper, P., Covacevich, J., Janetzki, H. and McDonald, K. 2000. Lizards. pp 203-233 in: *Wildlife of Tropical North Queensland*, edited by M. Ryan and B. Burwell. Queensland Museum, Brisbane.
- Covacevich, J.A. and Couper, P.J. 1991. The reptile records. pp 45-140 in: *An Atlas of Queensland's Frogs, Reptiles, Birds and Mammals*, edited by G.J. Ingram and R.J. Raven. Queensland Museum, Brisbane.
- Harfmann Short, K. and Petren, K. 2008. Boldness underlies foraging success of invasive *Lepidodactylus lugubris* geckos in the human landscape. *Animal Behaviour* 76: 429-437.
- Kluge, A.G. 1963. The systematic status of certain Australian and New Guinea gekkonid lizards. *Memoirs of the Queensland Museum* 14: 77-86.
- Kraus, F. 2009. *Alien Reptiles and Amphibians: A Scientific Compendium and Analysis*. Springer, New York.
- Limpus, C.J., Limpus, D.J. and Goldizen, A.R. 1999. Recent colonization of Heron Island, Great Barrier Reef, by the Mourning Gecko, *Lepidodactylus lugubris*. *Memoirs of the Queensland Museum* 43: 777-781.
- Lever, C. 2003. *Naturalized Reptiles and Amphibians of the World*. Oxford University Press, Oxford.
- McKay, J.L. and Horner, P. 2007. First records of the Mourning Gecko *Lepidodactylus lugubris* (Dumeril and Bibron, 1836) from the Northern Territory mainland. *Herpetofauna* 37 (2): 75-80.

- Messenger, K. 2005.** Behaviour of *Lepidodactylus lugubris* on Heron Island, Great Barrier Reef and a record of *Gehyra dubia* from that island. *Herpetofauna* 35 (1): 37-39.
- Scrymgour, M. 2006.** 'Invasive Gecko Species Found in Darwin Suburb'. Northern Territory Government Media Release, 11 December 2006. Available: www.nt.gov.au/nreta/publications/media/pdf/2006/12/scrygecko20061211.pdf
- Sprackland, R.G., Sprackland, T. and Diessner, D. 2004.** Reptiles and mammals of Fitzroy Island, Queensland. *Memoirs of the Queensland Museum* 49: 733-739.
- Turner, G. and Green, D. 1996.** Notes on the mourning gecko *Lepidodactylus lugubris* in the Daintree region. *Herpetofauna* 26 (1): 5-7.
- Wilson, S. and Swan, G. 2008.** *A Complete Guide to Reptiles of Australia*, 2nd Edition. New Holland Publishers, Sydney.
- Woinarski, J.C.Z., Horner, P., Fisher, A., Brennan, K., Linder, D., Gambold, N., Chatto R. and Morris I. 1999.** Distributional patterning of terrestrial herpetofauna on the Wessel and English Company Island groups, northeastern Arnhem Land, Northern Territory, Australia. *Australian Journal of Ecology* 24: 60-79.
- Yamashiro, S., Toda, M. and Ota, H. 2000.** Clonal composition of the parthenogenetic gecko, *Lepidodactylus lugubris*, at the northernmost extremity of its range. *Zoological Science* 17: 1013-1020.