REVIEW



WILEY

When an iconic native animal is overabundant: The koala in southern Australia

Desley A. Whisson 💿 | Kita R. Ashman

Deakin University, Geelong, Australia, School of Life and Environmental Sciences, 221 Burwood Highway, Burwood, Victoria, Australia

Correspondence

Desley A. Whisson, Deakin University, Geelong, Australia. School of Life and Environmental Sciences, 221 Burwood Highway, Burwood, Victoria 3125, Australia. Email: desley.whisson@deakin.edu.au,

Abstract

Management of overabundant native species is challenging at the best of times but is even more so when it involves an iconic species that is threatened in part of its range. Here, we examine koala overabundance, a situation that arises in some locations within the species' southern range. We consider the variable conservation status of the koala throughout its range, discuss where and why overabundance occurs, and its consequences. Programs to manage koala overabundance and impacts are generally ineffective, relying on costly and slow acting fertility control and translocation, and implemented only when a crisis already has occurred. A change in management approach is needed. Decisions should be based on an evidence-based and transparent decisionmaking framework that involves predicting where overabundance is likely, where damage costs will be high, and where proactive management is likely to be effective. With more than 65% of Australia's koala population occurring in the south and declining northern populations, a more considered approach to managing southern koalas for conservation of the species is needed.

KEYWORDS

culling, fertility control, overabundance. Phascolarctos cinereus, southern koalas, translocation

1 INTRODUCTION

World-wide, many native wildlife species have become locally abundant or have increased in geographic range communities (O'Connell-Rodwell, Rodwell, Rice, & Hart, (Garrott, White, & Vanderbilt White, 1993). In some situations, this has had negative consequences including expanded its range by almost 70%, resulting in considerimpacts on human health or livelihood, damage to ecosystems through unsustainable browsing or grazing on (Takatsuki, 2009). In Australia, populations of the once preferred resources, introduction or spread of infectious endangered saltwater crocodileCrocodylus porosus) have diseases and parasites to new areas, and a reduction in increased, resulting in an increase in human-crocodile species diversity or the relative abundance of sympatric conflicts (Brien et al., 2017; Fukuda, Manolis, & Appel, species (Garrott et al., 1993; Noss, 1990; Soulé, 1990)2014). An increase in abundance or distribution of a There are numerous examples of such conflicts. In east native species may result from one or more factors

Africa, populations of elephants Loxodonta africana) in some reserves have increased and now are damaging their environment and coming into conflict with local 2000). In Japan, the Sika deer Cervus nippon) has able browsing damage to crops and native vegetation

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in angiorne, provided the original work is properly cited.

© 2020 The Authors. Conservation Science and Practice published by Wiley Periodicals, Inc. on behalf of Society for Conservation Biology

including conservation actions that remove threats (e.g., hunting or predation), human-induced changes in ecosystems and landscapes, and introduction of the spe-koalas. Although this initially halted declines and allowed cies to areas outside of its native range (e.g., Gauthier, populations to recover, threats associated with habitat loss Giroux, Reed, Béchet, & Bélanger, 2005; Lunney et al., and fragmentation, disease, wildfire, changing climates, 2007; Waller & Alverson, 1997).

conflicts, wildlife managers have the difficult task of develoften depend on the iconic nature of the species. For A similar decline in 'southern koalas' has not been docuexample, culling alligators Alligator mississippiensis) in Florida was considered appropriate, whereas proposals to estimated koala numbers at 79,000 (range 33,0063,000) cull African elephants and white-tailed deer were strongly opposed (Hart, Jewell, & Holt, 1981). Similar dilemmas face management of the koalaPhascolarctos cinereus) in Australia. Although populations are declining in the northern part of its range (New South Wales, the Australian Capital Territory and Queensland), in southern states (Victoria and South Australia), koalas can be locally overabundant and have significant impacts on native ecosystems (McAlpine et al., 2015).

1.1 The koala: Northern versus southern populations

The koala is one of Australia's most iconic endemic species, ranking in the top three wildlife species that international tourists wish to see, and generating billions of dollars in tourism revenue (Hundloe & Hamilton, 1997). Its natural range extends from north-eastern Queensland to south-east South Australia (Natural Resources Management Ministerial Council 2009). The koala is a specialist folivore that subsists on Eucalyptus foliage. Its distribution is determined by the presence o*Eucalyptus* species that are high in available nitrogen and moisture, and low in plant secondary metabolites (Cork & Sanson, 1990; Moore & Foley, 2005). Drought and heat waves also sive management (Menkhorst, 2008). In Victoria, koalas are limiting factors (Adams-Hosking, Grantham, Rhodes, McAlpine, & Moss, 2011; Briscoe, Kearney, Taylor, & Wintle, 2016). Koalas are relatively solitary animals that primarily interact during the spring/summer breeding season (Ellis et al., 2015; Watchorn & Whisson, 2019), and generally occur at densities of less than one koala per hectare (Melzer, Carrick, Menkhorst, Lunney, & St. John, 2000).

Hunting and habitat clearance in the late 1800s decimated koala populations throughout their range. It has been estimated that by 1930, only a few hundred remained in New South Wales, up to ten thousand remained in Queensland, and less than a thousand koalas remained on the Victorian mainland (Lewis, 1954;

Menkhorst, 2008; Phillips, 1990). In the late 1920s, each state introduced legislation to stop hunting and to protect vehicle collisions and dog attacks have persisted (Natural When a native species becomes abundant and causesResource Management Ministerial Council, 2009). As a result, koala populations have declined in the last two oping socially acceptable management strategies (Dubois decades by an estimated 53% and 26% in Queensland and et al., 2017; Garrott et al., 1993). Management decisions New South Wales respectively (Adams-Hosking et al., 2016). mented (Adams-Hosking et al., 2016). In 2012, koala experts in Queensland, 36,000 (range 14,00703,000) in New South Wales and the Australian Capital Territory, 183,000 (range 77,000237,000) in Victoria, and 33,000 (range 19,000-51,000) in South Australia (Adams-Hosking et al., 2016). In 2015, only 'northern koalas' were listed as vulnerable under National legislation (Department of the Environment & Energy, 2015).

> Continued declines in koala populations are predicted due to climate change resulting in a greater incidence of drought and heat waves (Briscoe et al., 2016). Distribution modeling suggests that by 2070, the koala's range will contract toward the coast, and that most regions of high habitat suitability will be in the south (Briscoe et al., 2016). An increase in atmospheric Colevels also will impact habitat quality by reducing the nutritional quality of Eucalyptus foliage (Johnson, De Gabriel, & Moore, 2009). The conservation significance of southern koalas therefore is likely to increase (McAlpine et al., 2015). Although southern koalas are thought to be secure, their true status is unknown due to limited monitoring and field studies (Ashman, Watchorn, & Whisson, 2019; McAlpine et al., 2015). Declines may be occurring in some areas, but issues associated with high-density populations receive the most attention (McAlpine et al., 2015).

> Southern koalas have long been the subject of intenfrom island populations established in the late 19th and early 20th century were used to reestablish populations on the mainland where hunting had brought about their near-extinction, and to establish koalas in areas outside their historic range (e.g., Kangaroo Island, South Australia) (Menkhorst, 2008; Wedrowicz, Wright, Schlagloth, Santamaria, & Cahir, 2017). Between 1923 and 2006, over 12,000 koalas were translocated from French and Phillip Islands to the mainland (Menkhorst, 2008). In South Australia, koalas from Kangaroo Island were translocated to other parts of the state including the Eyre Peninsula, Riverlands, southeast SA, and the Adelaide Hills (Robertson, 1978). Although this resulted in the recovery of southern koala populations and

expansion of their range, most populations now have low genetic diversity, resulting from the island source populations being founded by just a few individuals (Cristescu et al., 2009; Houlden et al., 1999; Houlden, England, Taylor, Greville, & Sherwin, 1996).

The iconic nature of the koala combined with its variits range, creates challenges for policy makers and wildlife managers (Natural Resources Management Ministerial Council 2009; Shumway, Lunney, Seabrook, & McAlpine, 2015). On one hand, the koala is portrayed as a species that will be extinct by 2050 (Australian Koala Foundation, 2020); yet this is countered by reports of koala "overabundancë from the southern states, with calls to cull, sterilize, or translocate koalas to reduce high-density populations (Natural Resources Committee, South Australia, 2019; Possingham et al., 1996; South ferred habitats may always have occurred. However, only Australian Government, 2019). In some cases, highdensity populations have caused complete and widespread canopy defoliation resulting in mass starvation of koalas (Menkhorst, 2008; Whisson, Dixon, Taylor, & Melzer, 2016). Although koalas in southern states are not listed as "vulnerable" under National legislation, nor listed in any special conservation category under state legislation, their management is considered differently to other common native wildlife. Whereas permits are issued to cull 'common' native fauna (e.g., kangaroos) that are causing damage to natural environments or property, the same is not done for koalas. South Australian and Victorian state governments abide by national policy to reject culling of "overabundant" koalas (Australian and New Zealand Environment and Conservation Council, 1998; DELWP, 2004; South Australian Government, 2016). Management of overabundant koala populations therefore relies on translocation and fertility control measures that are expensive and may have poor welfare outcomes for koalas (Duka & Masters, 2005; Whisson et al., 2016; Whisson, Holland, & Carlyon, 2012). Furthermore, implementing such measures over large spatial scales where there may be thousands of The reason why overabundance occurs only in the southkoalas is logistically difficult and economically not feasible. As a result, these measures only may be effective in a to a complexity of climate and site-specific factors, or posfew situations (e.g., islands) where management areas are sibly the low genetic diversity of most southern koala small and there is limited potential for immigration (Massei & Cowan, 2014).

WHERE DOES KOALA 2 "OVERABUNDANCE " OCCUR?

We use the term" overabundant here to describe a koala population that occurs at a density that results in defoliation of preferred food trees. If unmanaged, this can lead

to widespread tree death, nutritional stress in the koala population, and even death of koalas due to starvation (e.g., Martin, 1985a, 1985b, 1985c; Whisson et al., 2016; Figure 1). Overabundance can occur where there are only two koalas per hectare (Menkhorst, 2008), but higher densities are typical. For example, at Cape Otway, able conservation status and management needs acrossWhisson et al. (2016) recorded 23 koalas per hectare immediately prior to mass starvation of koalas and a population crash. Koala overabundance has been documented in the scientific literature for 16 locations in Victoria and South Australia (Figure 2, Table 1). There only has been one report of overabundance outside of the koala's southern range, and that was reported for a forest remnant in northern New South Wales (Frith, 1979).

> Martin (1985b) suggested that fluctuations in koala populations (and thus occasional 'overabundance') in pretwo cases of overabundance have been documented for natural (i.e., nonreintroduced) populations (Walkerville and Wilson's Promontory). Kershaw (1934) reported that koalas were "numerous" in several areas of Wilson's Promontory c1905 and in"one or two other localities favored by koalas despite "periodical raids of skinhunters". In fact, translocation of koalas and culls were implemented on Wilson's Promontory c1910 to reduce populations and over browsing of trees (Kershaw, 1934). Most cases of overabundance have occurred in locations where koala populations have been reestablished via translocations (Table 1). For some of these, overabundance has been resolved either through management (e.g., Quail Island, Snake Island, Sandy Point) or as a result of koala overbrowsing itself causing both a crash in koala numbers and a reduction in habitat quality for koalas (e.g., Kennett River/Grey River; Menkhorst, 2008). However, at many sites, koala overabundance is a continuing concern.

3 CAUSES OF OVERABUNDANCE

ern part of the koala's range is not clear and may be due populations. Fecundity rates for many southern koala populations (32 to 81% McLean, 2003) are within the range for northern koalas (e.g., 33 to 85% recorded by Thompson, 2006). At some sites where overabundance occurs, fecundity can be low due to disease associated with Chlamydia pecorum(e.g., 38.8% fecundity for the population in Budj Bim National Park [McLean, 2003]). Overabundance also can occur where koalas face considerable threats. For example, the koala population in the Adelaide Hills and Mt Lofty Ranges, a highly urbanized



FIGURE 1 Koala overabundance at Cape Otway in November 2013. (a) Koalas converged on trees that had not been completely defoliated, and (b) individuals became nutritionally stressed and starved to death




FIGURE 2 Locations of koala overabundance in Victoria and South Australia. The light grey shading shows the distribution of koalas in Australia (Woinarski & Burbidge, 2016)

region, was recently estimated at around 150,000 individ- 2019). In addition to high mortality due to dog attacks uals with densities of up to 13 koalas per hectare and vehicular collisions, this population is impacted by a recorded in some areas (Sequeira, Roetman, Daniels, high incidence of calcium oxalate nephrosis, a condition Baker, & Bradshaw, 2014; South Australian Government, that results in kidney failure (Speight et al., 2013).

Year(s) and Eucalyptus Forest cover numberof koalas Dates of Location species at site within10km introduced "outbreaks" References Mitchell, Bilney, & Martin, Raymond Island E.viminalis pryoriana N/A 1953:42 koalas 1985 Present 1988 (765 ha) Snake Island E.viminalis pryoriana N/A 1945:133 koalas 1990-2012 McLean, 2003 (3,452 ha) 1955, 1963, 1977: Unknown 75% Barrett, 1937 Wilsons promontory E. viminalis N/A - natural 1905-1910 Kershaw, 1915, 1934 Walkerville E. ovata 13% N/A - natural c1980 Martin, 1981, 1985a, 1985b, 1985c Phillip Island 1870-1923: E. viminalis N/A 1941-1978 Lewis, 1954 (10,000 ha) Unknown Gliddon, 1958 Menkhorst, 2008 Wedrowicz et al., 2017 1923 Present French Island E. viminalis N/A c1898: 3 koalas Lewis, 1934 (17,392 ha) Lewis, 1954 McNally, 1960 Quail Island E. viminalis N/A 1930-1933: 1944-1945 Braithwaite, Lumsden, & (530 ha) 165 koalas Dixon, 1980 E. viminalis 17% 1972:20 koalas Sandy point 1985 Martin, 1989 (Mornington Backhouse & Crouch, 1991 peninsula) Kennett river & grey E. viminalis 90% 1958, 1977 and c2000 Menkhorst, 2008 1982:294 koalas river E. cypellocarpa Cape Otway E. viminalis 71% 1981:75 koalas 2013Present Whisson et al., 2016 Menkhorst, Ramsey, O'Brien, Hynes, & Whisson, 2019 Framlingham forest E. viminalis 5% 1970:37 koalas 1987999 Wallis, 2013 E. ovata & Hopkins river 2016-2018 E. camaldulensis 19926003 Menkhorst, 2008 Tower Hill game E. viminalis 1% 1979; 17 koalas reserve Middleton, Walters, Menkhorst, & Wright, 2003 1973:30 koalas 1999-Present Budj Bim NP, stones E. viminalis 27% McLean, 2003 reserve 1982:46 koalas **DELWP**, 2017 & Kutonitj indigenous protected area Far SW Victoria E. viminalis Variable See Budj Bim NP c2013 - present Ashman, Rendall, E. ovata Symonds, & Whisson, 2020 c1965Present Philpott, 1965 Kangaroo Island E. viminalis cygnetensisFlinders chase: 94%1923:18 koalas (440,500 ha) E. ovata Cygnet R: 31% Eberhard, 1972 Robertson, 1978 Masters, Duka, Berris, & Moss, 2004 Adelaide hills & Mt E. viminalis 61% c2016-Present SA Government, 2016 19591969: lofty ranges Unknown

TABLE 1 Sites where koala overabundance and over browsing have been documented (see Figure 2)

primary cause of overabundance (Australian Koala Foundation, 2000) and this may be true on offshore islands or densities than typically occur in other parts of the koala's in habitat 'islands' surrounded by cleared landscapes on range. the mainland (Menkhorst, 2008). However, island populations also occur in the north and have not become overabundant (e.g., McGregor, Kerr, & Krockenberger, 2013). Furthermore, overabundance can occur at sites OVERABUNDANCE surrounded by other suitable koala habitat (e.g., Cape Otway, Kennett River, far south-west Victoria, and Flinders Chase on Kangaroo Island, Table 1). In far southwest Victoria, koala overabundance in remnant woodlands and in roadside vegetation has resulted from the establishment of commercial blue gumE. globulus) plantations (Ashman et al., 2020). Blue gum is highly preferred by koalas and population densities of up to 4.3 koalas per hectare have been documented in plantations (Ashman et al., 2020). When plantations are harvested, issues in native vegetation.

is the presence of E. viminalis (manna gum) and/or E. ovata (swamp qum) (Menkhorst, 2008). These species that bird species diversity and richness declined with contain high levels of available nitrogen and leaf moisture, and low levels of anti-herbivory compounds, thus making them ideal koala food trees (Moore & Foley, 2005). In woodlands comprising a high proportion of these species, koalas have small home ranges (less than two hectares; Mitchell, 1990; Whisson et al., 2016). Furthermore, koalas in these habitats exhibit high site fidelity and appear reluctant to move to alternate habitat types even when trees become completely defoliated (Whisson et al., 2016). Recent research into the koala's within the southern states, there is little or no recognition microbiome suggests that this may be due to koalas becoming highly specialized on a single food tree such that they may have limited ability to switch their diet to another food tree species (Blyton et al., 2019; Brice serving the genetically diverse population in South et al., 2019).

Another factor that may contribute to overabundance is low genetic diversity resulting in behavioral changes in koalas. Except for one population in South Gippsland, southern koalas have extremely low genetic diversity resulting from the near-extinction bottleneck in the 1920s (Cristescu et al., 2009; Houlden et al., 1996). Although a reduction in genetic diversity typically leads to reduced reproductive success in a species (Crnokrak & Roff, 1999; the koala's iconic nature and the limited and costly man-Seymour et al., 2001), this has not been the case for agement tools available. Culling is not endorsed by Com-Carlyon, 2010). However, low genetic diversity also may actions are limited to translocation, fertility control, result in individuals being less aggressive. This has been exclusion methods, and habitat management (Natural observed in house mice (Barnard & Fitzsimons, 1989; Resource Management Ministerial Council, 2009). At a (Tsutsui, Suarez, Holway, & Case, 2000) but remains groups of trees with tree guards or a koala proof fence

Lack of dispersal opportunity has been touted as the untested with koalas. If true for koalas, individuals may better tolerate being near conspecifics, permitting higher

CONSEQUENCES OF 4

Overabundance has negative consequences both for koalas and the ecosystem. Defoliation of trees may result in nutritional stress of koalas and in severe cases, mass starvation (Whisson et al., 2016). Impacts to ecosystems range from the defoliation or death of some preferred trees through to a drastic change in the ecosystem resulting from widespread tree death (Menkhorst, 2008). Such impacts have been greatest in Damp-sands herbrich woodland", a vegetation association that already is koalas must disperse, thus exacerbating overabundance listed as vulnerable (under state legislation) across most of Victoria (Menkhorst, 2008). The flow-on effects of tree One factor in common across sites of overabundance defoliation are poorly understood but likely to be highly negative. Whisson, Orlowski, and Weston (2018) found increasing canopy defoliation. Defoliation also may lead to changes in subcanopy vegetation due to increased light infiltration and changes in microhabitats (Valladares, Laanisto, Niinemets, & Zavala, 2016).

> Koala overabundance also impacts on policy decisions, and allocation of resources for conservation. The relative abundance of southern koalas long hindered recognition of northern koalas as vulnerable under Commonwealth legislation (Shumway et al., 2015). Even of pervasive threats and declining populations in some areas; and limited resources are allocated to the monitoring or management of declining populations or to con-Gippsland (Wedrowicz et al., 2017; Wedrowicz, Mosse, Wright, & Hogan, 2018).

5 | MANAGEMENT OF **OVERABUNDANT KOALAS**

Managing koala overabundance is challenging because of southern koalas (McLean & Handasyde, 2006; Whisson & monwealth or State governments and management Eklund, 1996), salmon (Tiira et al., 2003) and ants local scale, koalas may be excluded from single or small

(for example fence design, see Queensland Department thousands of koalas (Duka & Masters, 2005: Natural ment to reduce the potential for an area to support highdensity koala populations also may be effective. Au et al. first decade of their implementation (Molsher, 2017; (2019) showed that koala abundance at a site is related to Todd et al., 2008), establishment of commercial blue gum the nutritional quality (available nitrogen in canopy foliage) of forests. Consequently, revegetating over- now placing the long-term effectiveness of those probrowsed sites with less palatable tree species may result grams in doubt. On Kangaroo Island, commercial blue in sustainable populations.

Translocation and fertility control can be implemented Australian Government, 2016). Of these, translocation can immediately reduce koala densities at a site; however, the viability of translocation relies on the availability of suitable sites to which koalas can be translocated. In Victoria, sites must containEucalyptusspecies that are preferred by koalas, occur at 700 m elevation, be at least 1,000 ha and at least 1 km from a major road or railway (Menkhorst et al., 2019). Furthermore, for any species, translocation is the koala population (Ashman et al., 2020; Ashman & inherently risky, carrying the risks of spreading disease, causing overabundance in the recipient site, and high animal welfare costs (see Massei, Quy, Gurney, & Cowan, likely. 2010 for review). For koalas, variable survival has been recorded, with success depending on the health of individuals, their treatment prior to translocation (e.g., sterilization method), and the habitat and weather conditions at the release site (see Menkhorst, 2017 for review). In contrast to translocation, fertility control is slow acting, taking between 5 and 10 years for a positive habitat response November 2013 after widespread tree defoliation already (Todd, Forsyth, & Choquenot, 2008). Although surgical sterilization of males and females was used in early programs (e.g., Kangaroo Island; Duka & Masters, 2005), management strategy to prevent a repeat of the situation treatment of female koalas with levonorgestrel implants is now the standard approach (Hynes, Handasyde, Shaw, & world-wide negative media attention (DELWP, 2019a). Renfree, 2010; Menkhorst et al., 2019; Middleton et al., At the same time, the Victorian Government initiated 2003; Todd et al., 2008).

being resolved using these methods. Exceptions are Quail (Ramsey, Tolsma, & Brown, 2017). However, the effec-Island where in 1945, the entire population of 1,308 koalas was translocated to the mainland; Sandy Point ing at sites where overabundance is likely and the ability (1,522 koalas translocated), and Snake Island where a to respond quickly when thresholds for management are combination of translocation (2,607 koalas as of 2006) "triggered". Unfortunately, neither of these conditions and fertility control reduced the population to a sustainable level (Menkhorst, 2008). In most cases, ongoing undertaken in response to a crisis (i.e., starving koalas management is required due to the large geographic scale and widespread tree defoliation and death) in October and size of the koala population, and logistical constraints that limit the proportion of the total population that can be sterilized; and because of immigration of koalas into treated areas (Todd et al., 2008). Long-term mately 120 km north west of Framlingham (DELWP, (>10 years) management programs include those on Raymond Island, Budj Bim National Park, and Kangaroo Island. The Budj Bim NP and Kangaroo Island programs have involved the sterilization and translocation of

of Transport and Main Roads, 2019). Habitat manage- Resources Kangaroo Island, 2019; Todd et al., 2008). Although both of these programs were successful in the plantations in the landscape surrounding treated areas is gum plantations were first established in the 1990s. In 2015, they covered 13,198 ha and were estimated to supby government departments (DELWP, 2004; South port 23,360 ± 3,330 koalas (Molsher, 2017). An increase in the koala population in native vegetation from 14,270 ±759 in 2010 to 25,146 ± 2,646 in 2015 was mostly attributed to immigration of koalas from blue gum plantations (Molsher, 2017). Similarly, Budj Bim NP in far south west Victoria lies within a region where over 80,000 ha of blue gum plantations have been established and have facilitated a considerable wide-scale increase in Watchorn, 2019). Consequently, constant immigration of Budj Bim NP by unsterilized koalas from plantations is

Reactive management in response to public outcry is typical for most cases of koala overabundance and government management actions tend to focus on addressing koala welfare. For example, at Cape Otway, a "welfare intervention" involving the euthanasia of approximately 700 starving koalas was implemented in had occurred and large numbers of koalas already had starved to death (Whisson et al., 2016). A longer-term was not implemented until March 2015, coinciding with much-needed research into habitat triggers for manage-There are relatively few cases of koala overabundance ment" to predict when problems were likely to occur tiveness of this approach relies both on annual monitorwere met for Framlingham where management was 2018. In a one-week program, 194 koalas were captured, with 168 of those translocated (females sterilized with levonorgestrel implants) to two state forests approxi-2019b).

> There are no easy solutions for managing koala overabundance. Many advocate culling as an answer but considering the koala's iconic status and the high level of

public outrage expressed when culling has been suggested island) including 95% of national park land and a large (Bagust, 2010), it seems unlikely that this method would ever receive government endorsement. If management koalas (Department for Environment and Water, 2020). continues to rely on translocation and fertility control, decisions on where and when to apply these tools should approximately 6.369 ha and over 50% of koala habitat. be based on an evidence-based and transparent decisionmaking framework (Dicks, Walsh, & Sutherland, 2014; Dubois et al., 2017). For koalas, this means predicting where and when overabundance will occur, its costs (animal welfare, ecosystem, human dimensions), and evaluating both the cost and potential efficacy of management actions for each location. With this approach, decisions to not implement fertility control and translocation may be warranted in many situations. In such cases, the welfare of koalas and condition of habitat should be monitored, and actions taken to alleviate koala suffering (e.g., euthanasia of koalas) and to restore damaged habitats.

THE FUTURE OF SOUTHERN 6 **KOALAS**

With approximately 65% of Australia's total koala population in southern Australia, and population declines occurring throughout the koala's northern range, maintaining healthy populations of southern koalas is becoming increasingly important for conservation of the species. Although southern koalas currently are considered secure, there is limited understanding of their distribution and abundance (McAlpine et al., 2015). Like northern koalas, they especially are threatened by continued habitat loss, fragmentation and degradation; an increasing frequency of droughts and wildfire as a result of warmer and drier climates; and declining nutritional value of *Eucalyptus* foliage with increasing atmospheric CO₂ (McAlpine et al., 2015).

Habitat loss and fragmentation limit both the distribution and long-term viability of koala populations in a landscape (Januchowski et al., 2008; McAlpine et al., 2006). Victoria and South Australia have suffered high rates of deforestation such that forests now cover approximately 34% and 9% of land in each state respectively (Bradshaw, 2012). Remaining forests tend to be highly fragmented, disturbed or ecologically compromised (Bradshaw, 2012; Gill & Williams, 1996; Norton, 1996). In highly fragmented landscapes where koalas may spend more time on the ground moving between habitat patches, the risk of predation and vehicular collisions are (McAlpine et al., 2006). Wildfires have the potential to decimate entire populations of koalas, and impact on habitat quality. On Kangaroo Island, a fire that started in December 2019 burnt 210,000 ha (almost 50% of the engagement and collaboration with stakeholders, and

proportion of koala habitat supporting thousands of At the same time in Victoria, a fire in Budj Bim NP burnt

In addition to these threats, most populations of southern koalas are inherently at risk of collapse due to their low genetic diversity (Wedrowicz et al., 2018). Such was the case with the Tasmanian devil Sarcophilus harrisii) that also had a history of genetic bottlenecks resulting in low genetic diversity (Siddle et al., 2007). Although abundant prior to 1996, Tasmanian devil populations now have decreased by between 50% and 90% due to the emergence of the highly contagious, devil facial tumor disease (Siddle et al., 2007). Only one southern koala population (in South Gippsland, Victoria) has a relatively high level of genetic diversity which could provide it with a greater chance of survival compared to reintroduced populations (Houlden et al., 1999: Wedrowicz et al., 2018). However, despite the conservation significance of this population, there is limited understanding of the population's extent (abundance and distribution) or threats to its persistence (Wedrowicz et al., 2018).

A comprehensive management strategy for southern koalas is needed. Such a strategy must not just focus on issues of overabundance but aim to address the diversity of issues that impact on koala conservation, and to improve our understanding of the ecolary of the species in its southern range. Compared to northern states, there has been relatively little research on southern koalas, and a stronger research focus on tools for managing overabundance (Ashman et al., 2019). An improved understanding of koala distribution, abundance and population fluctuations, current and future threats, population health and genetic diversity, and gene flow between populations is needed to inform effective conservation actions. Furthermore, management should be considered t the landscape-scale and include both the rehabilitation of degraded habitats, and reforestation to increase haits area and connectivity. Given the propensity for southern koalas to become overabundant in some habitat types, habitat management actions will need to be carefully planned to create habitats that can sustain koala populatons, but that do not result in overabundance. Reforestization is not only important for koala conservation but will improve biodiversity and help mitigate the effects of climate change (Paul et al., 2016).

For a management strategy to be effective in deliverheightened, and energy budgets may be compromised ing on-ground conservation outcomes for koalas, it also must include a process for implementation, and funding to support actions (McAlpine et al., 2015). Implementation of a strategy is often challenging, requiring extensive

www.environment.sa.gov.au/news-hub/news/articles/2020/01/ bushfire-and-recovery.

Department of Environment, Land, Water and Planning (DELWP). (2004). Victoria's koala management strategy. https://www. wildlife.vic.gov.au/__data/assets/pdf_file/0030/27993/Koala.pdf.

Department of Environment, Land, Water and Planning (DELWP). 2017. Kurtonitj Koala management program. Retrieved from https://www.wildlife.v ic.gov.au/our-wildlife/koalas/kurtonitjkoala-management-program-november-2017.

Department of Environment, Land, Water and Planning (DELWP). (2019a). Koalas at Cape Otway: Timeline of events. Retrieved Houlden, B. A., England, P. R., Taylor, A. C., Greville, W. D., & https://www.wildlife.vic.gov.au/ data/assets/pdf file/ from 0020/73631/Koalas-at-Cape-Otway-Fact-Sheet 3 timeline-2017-Autumn-POST-DELIVERY.pdf.

Department of Environment, Land, Water and Planning (DELWP). (2019b). Koalas. Retrieved from https://www.wildlife.vic.gov. au/our-wildlife/koalas.

Department of the Environment and Energy. (2015). Species Profile and Threats Database Phascolarctos cinereus (combined populations of Qld, NSW and the ACT). Retrieved from http:// www.environment.gov.au/cgi-bin/sprat/public/publicspecies. pl?taxon_id=85104.

Dicks, L. V., Walsh, J. C., & Sutherland, W. J. (2014). Organising evidence for environmental management decisions: A '4S' hierarchy. Trends in Ecology and Evolution, 29, 607-613.

Dubois, S., Fenwick, N., Ryan, E. A., Baker, L., Baker, S. E., Beausoleil, N. J.,.. Fraser, D. (2017). International consensus principles for ethical wildlife control. Conservation Biology, 31, 753-760.

Duka, T., & Masters, P. (2005). Confronting a tough issue: Fertility control and translocation for over-abundant koalas on Kangaroo Island, South Australia. Ecological Management & Restoration. 6. 172-181.

Eberhard IH. 1972. Ecology of the koalaPhascolarctos cinereus Goldfuss on Flinders chase, Kangaroo Island. Ph.D. Thesis, University of Adelaide. Retrieved from http://hdl.handle.net/ 2440/19715.

Eklund, A. (1996). The effects of inbreeding on aggression in wild male house mice Mus domesticus). Behaviour, 133, 883-901.

Ellis, W., Fitzgibbon, S., Pve, G., Whipple, B., Barth, B., Johnston, S.,... Bercovitch, F. (2015). The role of bioacoustic signals in koala sexual selection: Insights from seasonal patterns of associations revealed with GPSProximity unit BLoS One, 10, e0130657.

Frith, H. J. (1979). Wildlife conservation (p. 416). Sydney, Australia: Angus and Robertson.

Fukuda, Y., Manolis, C., & Appel, K. (2014). Featured article: Management of human-crocodile conflict in the Northern Territory, Australia: Review of crocodile attacks and removal of problem crocodiles. Journal of Wildlife Management, 78, 1239-1249.

Garrott, R. A., White, P. J., & Vanderbilt White, C. A. (1993). Overabundance: An issue for conservation biologist@nservation Biology, 7, 946-949.

Gauthier, G., Giroux, J. F., Reed, A., Béchet, A., & Bélanger, L. (2005). Interactions between land use, habitat use, and population increase in greater snow geese: What are the consequences for natural wetlands? Global Change Biology, 11, 856-868.

Gill, A. M., & Williams, J. E. (1996). Fire regimes and biodiversity: The effects of fragmentation of southeastern Australian eucalypt forests by urbanisation, agriculture and pine plantations. Forest Ecology Management, 85, 261-278.

Gliddon, J. W. (1958). Phillip Island in picture and story. Phillip Island: The Committee of Trust"Warley", Cowes Bush Nursing Hospital.

Hart, D., Jewell, A., & Holt, S. (1981)Problems in management of locally abundant wild mammals (pp. 1-5). Cambridge, MA: Academic Press.

Houlden, B. A., Costello, B. H., Sharkey, D., Fowler, E., Melzer, A., Ellis, W., Carrick, F. (1999)

Sherwin, W. B. (1996). Low genetic variability of the koala Phascolarctos cinereus in South-Eastern Australia following a severe population bottleneckMolecular Ecology, 5, 269-281.

Hundloe, T., & Hamilton, C. (1997). Koalas and tourism: An economic evaluation. Discussion paper 13. The Australia Institute, Canberra. Retrieved from https://www.tai.org.au/sites/default/ files/DP13 8.pdf.

Hynes, E. F., Handasyde, K. A., Shaw, G., & Renfree, M. B. (2010). Levonorgestrel, not etonogestrel, provides contraception in free-ranging koalas.Reproduction, Fertility and Development, 22, 913-919.

Januchowski, S. R., McAlpine, C. A., Callaghan, J. G., Griffin, C. B., Bowen, M., Mitchell, D., & Lunney, D. (2008). Identifying multiscale habitat factors influencing koala Rhascolarctos cinereus) occurrence and management in Ballarat, Victoria, Australia. Ecological Management and Restoration, 9, 134-142.

Johnson, C. N., De Gabriel, J. L, & Moore, B. D. (2009). Koalas and climate change: hungry for CQ cuts. Retrieved from http:// cmsdata.iucn.org/downloads/fact_sheet_red_list_koala.pdf.

Kershaw, J. A. (1915). Excursion to National Park, Wilsons promontory. Victorian Naturalist, 31, 143-152.

Kershaw, J. A. (1934). The koala on Wilson's promontor Victorian Naturalist, 51, 76-77.

Lewis, F. (1934). The koala in Victoria. Victorian Naturalist, 51, 73-76.

Lewis, F. (1954). The rehabilitation of the koala in VictoriaVictorian Naturalist, 70, 197-200.

Lunney, D., Baker, J., Matthews, A., Waples, K., Dickman, C., & Cogger, H. (2007). Overabundant native vertebrates in New South Wales: Characterising populations, gauging perceptions and developing an ethical management framework. In P. A. Hutchings, D. Lunney, S. Burgin, & P. Eby (Eds.) Pest or Guest (pp. 158-173). Mosman, New South Wales: The Roval Zoological Society of New South Wales.

Martin, R. (1989). Draft management plan for the conservation of the koala (Phascolarctos cinereus) in Victoria. Technical Report Series No. 99. National Parks and Wildlife Division, Department of Conservation, Forests and Lands, Victoria, Melbourne. 64.

Martin, R. W. (1981). Age-specific fertility in three populations of the koala, Phascolarctos cinereus Goldfuss, in Victoria. Wildlife Research, 8, 275-283.

Martin, R. W. (1985a). Overbrowsing and decline of a population of the koala, Phascolarctos cinereus, in Victoria. I. Food preference and food tree defoliation. Wildlife Research, 12, 355-365.

Martin, R. W. (1985b). Overbrowsing and decline of a population of the koala, Phascolarctos cinereus, in Victoria. II. Population condition. Wildlife Research, 12, 367-375.

11 of 12

- Martin, R. W. (1985c). Overbrowsing, and decline of a population of the koala, Phascolarctos cinereus, in Victoria III. Population dynamics. Wildlife Research, 12, 377-385.
- Massei, G., & Cowan, D. P. (2014). Fertility control to mitigate human-wildlife conflicts: A review. Wildlife Research, 41, 1-21.
- Massei, G., Quy, R. J., Gurney, J., & Cowan, D. P. (2010). Can trans- Mitchell, P. J., Bilney, R., & Martin, R. W. (1988). Population struclocations be used to mitigate humanwildlife conflicts? Wildlife Research, 37, 428-439.
- Masters, P., Duka, T., Berris, S., & Moss, G. (2004). Koalas on Kan- Molsher, R. (2017). Kangaroo Island koala population survey 2015. garoo Island: From introduction to pest status in less than a century. Wildlife Research, 31, 267-272.
- McAlpine, C., Lunney, D., Melzer, A., Menkhorst, P., Phillips, S., Phalen, D.,... Close, R. (2015). Conserving koalas: A review of the contrasting regional trends, outlooks and policy challenges. Biological Conservation, 192, 226-236.
- McAlpine, C., Rhodes, J. R., Callaghan, J. G., Bowen, M. E., Natural Resource Management Ministerial Council 2009. National Lunney, D., Mitchell, D. L., ... Possingham, H. P. (2006). The importance of forest area and configuration relative to local habitat factors for conserving forest mammals: A case study of koalas in Queensland, Australia Biological Conservation, 132, 153-165.
- McGregor, D. C., Kerr, S. E., & Krockenberger, A. K. (2013). The distribution and abundance of an Island population of koalas (Phascolarctos cinereus) in the far north of their geographic range.PLoS One, 8, e59713.
- McLean, N. (2003). Ecology and management of overabundant koala (Phascolarctos cinereus) populations. PhD Thesis, The University of Melbourne. Retrieved from: https://minervaaccess.unimelb.edu.au/handle/11343/39249.
- McLean, N., & Handasyde, K. A. (2006). Sexual maturity, factors sons in populations of overabundant Victorian koalas (Phascolarctos cinereus). Australian Journal of Zoology, 54, 385-392.
- McNally J. (1960). Koala management in Victoria. Fisheries and Wildlife Department. Fauna Contribution 8.
- Melzer, A., Carrick, F., Menkhorst, P., Lunney, D., & St. John, B. (2000). Overview, critical assessment, and conservation implications of koala distribution and abundanceConservation Biologv. 14. 619-628.
- Menkhorst, P. (2008). Hunted, marooned, re-introduced, contracepted: A history of koala management in Victoria. Australian Zoologist, 34, 73-92.
- Menkhorst, P. (2017). Literature review: Responses of koalas to translocation. Arthur Rylah Institute for Environmental Research Technical Report Series Number 279. Department of Phillips, W. (1990). Koalas: The little Australian we'd all hate to lose. Environment, Land, Water and Planning, Heidelberg, Victoria. Retrieved from: https://www.ari.vic.gov.au/__data/assets/pdf_ file/0023/64391/ARI-Technical-Report-279-Literature-reviewresponses-of-koalas-to-translocation.pdf.
- Menkhorst, P. W., Ramsey, D. S. L., O'Brien, T., Hynes, E., & Whisson, D. (2019). Survival and movements of koalas translocated from an over-abundant population.Wildlife Research, 46, 557-565.
- Middleton, D. R., Walters, B., Menkhorst, P., & Wright, P. (2003). Fertility control in the koala, Phascolarctos cinereus: The impact of slow-release implants containing levonorgestrel or oestradiol on the production of pouch youngWildlife Research, 30, 207-212.

- Mitchell, P. J. (1990). The home ranges and social activity of koalas-A quantitative analysis. In A. K. Lee. K. A. Handasyde, & G. D. Sanson (Eds. Biology of the koala (pp. 171-187). New South Wales, Australia: Surrey Beatty, Chipping Bay.
- ture and reproductive status of koalas on Raymond Island, Victoria. Wildlife Research, 15, 511-514.
- Natural Resources Kangaroo Island, SA. Retrieved from https:// www.naturalresources.sa.gov.au/files/sharedassets/kangaroo_ island/plants and animals/native animals/koala/kangarooisland-koala-populaton-survey-2015.pdf.
- Moore, B. D., & Foley, W. J. (2005). Tree use by koalas in a chemically complex landscapeNature, 435, 488-490.
- Koala conservation and management strategy 2002014. Retrieved from http://www.environment.gov.au/biodiversity/ threatened/publications/national-koala-conservation-mgtstrategy-2009-2014.
- Natural Resources Committee, South Australia. (2019). Inquiry into management of overabundant and pest species. Third report of the 54th Parliament of the Natural Resources Committee. Retrieved from https://www.environment.sa.gov.au/files/ sharedassets/public/plants_and_animals/abundant_species/ overabundant-pest-species-report.pdf.
- Natural Resources Kangaroo Island. (2019). Koala management. Retrieved from: https://www.naturalresources.sa.gov. au/kangarooisland/plants-and-aimals/native-animals/koalamanagement.
- affecting the breeding season and breeding in consecutive sea- Norton, T. W. (1996). Conserving biological diversity in Australia's temperate eucalypt forests Forest Ecology Management, 85, 21-33.
 - Noss, R. F. (1990). Can we maintain biological and ecological integrity? Conservation Biology, 4, 241-243.
 - O'Connell-Rodwell, C. E., Rodwell, T., Rice, M., & Hart, L. A. (2000). Living with the modern conservation paradigm: Can agricultural communities co-exist with elephants? A five-year case study in East Caprivi, NamibiaBiological Conservation, 93. 381-391.
 - Paul, K. I., Cunningham, S. C., England, J. R., Roxburgh, S. H., Preece, N. D., Lewis, T.,.. Polglase, P. J. (2016). Managing reforestation to sequester carbon, increase biodiversity potential and minimize loss of agricultural land. Land Use Policy, 51, 135-149.
 - Canberra, Australia: Australian National Parks and Wildlife Service. AGPS Press.
 - Philpott, C. M. (1965). The ecology of the koala Phascolarctos cinereus (Goldfuss) on flinders chase, Kangaroo Island. Honours thesis. South Australia: University of Adelaide.
 - Possingham, H., Barton, M., Boxall, M., Dunstan, J., Gibbs, J., Grieg, J.,... St John, B. (1996)Koala management task force final report. Adelaide, SA: University of Adelaide.
 - Queensland Department of Main Roads. 2019. Koala proof fence and gate. Retrieved from https://www.tmr.qld.gov.au/-/media/ busind/techstdpubs/Specifications-and-drawings/Standard-Drawings-Roads/Roadworks-Drainage-Culverts-and-Geotechnical/ SD1603.pdf.

12 of 12 WILEY Conservation Science and Practice

- Ramsey, D. S. L, Tolsma, A. D. & Brown, G. W. (2017). Towards a habitat condition assessment method for guiding the management of overabundant Koala populations. Arthur Rylah Institute for Environmental Research Technical Report Series No. 272. Department of Environment, Land, Water and Planning, Heidelberg, Victoria. Retrieved from https://www.ari.vic. gov.au/__data/assets/pdf_file/0012/44112/ARI-Technical-Report-272-Habitat-assessment-to-guide-management-of-overabundantkoalas.pdf.
- Robertson, A. C. (1978). The koala in South Australia. In T. J. Bergin (Ed.), *The koala: Proceedings of the Taronga sympo*sium on koala biology, management and medicine, Sydney 11th and 12th march, 1976 (pp. 132-143). Sydney: Zoological Parks Board of N.S.W..
- Sequeira, A. M. M., Roetman, P. E. J., Daniels, C. B., Baker, A. K., & Bradshaw, C. J. A. (2014). Distribution models for koalas in South Australia using citizen science-collected data. *Ecology and Evolution*, *4*, 2103-2114.
- Seymour, A. M., Montgomery, M. E., Costello, B. H., Ihle, S., Johnsson, G., St. John, B., Houlden, B. A. (2001). High effective inbreeding coefficients correlate with morphological abnormalities in populations of south Australian koalas *Rhascolarctos cinereus*). *Animal Conservation*, *4*, 211-219.
- Shumway, N., Lunney, D., Seabrook, L., & McAlpine, C. (2015). Saving our national icon: An ecological analysis of the 2011 Australian senate inquiry into status of the koala*Environmental Science and Policy*, *54*, 297–303.
- Siddle, H. V., Kreiss, A., Eldridge, M. D. B., Noonan, E., Clarke, C. J., Pyecroft, S., Belov, K. (2007). Transmission of a fatal clonal tumor by biting occurs due to depleted MHC diversity in a threatened carnivorous marsupial *Proceedings of the National Academy of Sciences of the United States of America*, 104, 16,221-16,226.
- Soulé, M. E. (1990). The onslaught of alien species, and other chal- Whisson, D. A., Dixon, V., Taylor, M. L., & Melzer, A. (2016). Faillenges in the coming decade *Conservation Biology*, *4*, 233-240. ure to respond to food resource decline has catastrophic cons-
- South Australian Government. (2016). The south Australian koala conservation and management strategy. Department of Environment, Water and Natural Resources. Retrieved from https://www.environment.sa.gov.au/files/sharedassets/public/plants_and_animals/koala-conservation-and-management-strategy-gen.pdf.
- South Australian Government. (2019). Environment SA News Koalas and hills habitat to benefit from fertility control. Retrieved from https://www.environment.sa.gov.au/news-hub/ news/articles/2019/05/koala-contraception.
- Speight, K. N., Boardman, W., Breed, W. G., Taggart, D. A., Woolford, L., & Haynes, J. I. (2013). Pathological features of oxalate nephrosis in a population of koalas *Phascolarctos cinereus*) in South Australia. *Veterinary Pathology*, *50*, 299-307.
- Takatsuki, S. (2009). Effects of sika deer on vegetation in Japan: A review. *Biological Conservation*, *142*, 1922-1929.
- Thompson, J. (2006). The comparative ecology and population dynamics of koalas in the koala coast region of south-East Queensland. PhD Thesis, School of Integrative Biology, University of Queensland. Retrieved from https://espace.library.uq. edu.au/view/UQ:158282.

- Ramsey, D. S. L, Tolsma, A. D. & Brown, G. W. (2017). Towards a Tiira, K., Laurila, A., Peuhkuri, N., Piironen, J., Ranta, E., & Primmer, C. R. (2003). Aggressiveness is associated with genetic diversity in landlocked salmon *§almo salar*). *Molecular Ecology*, *12*, 2399-2407.
 - Todd, C. R., Forsyth, D. M., & Choquenot, D. (2008). Modelling the effects of fertility control on koala-forest dynamics*Journal of Applied Ecology*, 45, 568-578.
 - 272-Habitat-assessment-to-guide-management-of-overabundantkoalas.pdf. ertson, A. C. (1978). The koala in South Australia. In T. J. Bergin (Ed.),*The koala: Proceedings of the Taronga sympo-*Reduced genetic variation and the success of an invasive species. *Proceedings of the National Academy of Sciences of the* United States of America, 97(5), 948-5,953.
 - Valladares, F., Laanisto, L., Niinemets, Ü., & Zavala, M. A. (2016). Shedding light on shade: Ecological perspectives of understorey plant life. *Plant Ecology and Diversity*, 9, 237-251.
 - Waller, D. M., & Alverson, W. S. (1997). The White-tailed deer: A keystone herbivore *Wildlife Society Bulletin*, *25*, 217-226.
 - Wallis, R. L. (2013). Koala Phascolarctos cinereus in Framlingham Forest, south-West Victoria: Introduction, translocation and the effects of a bushfire Victorian Naturalist, 130, 37–40.
 - Watchorn, D. J., & Whisson, D. A. (2019). Quantifying the interactions between koalas in a high-density population during the breeding period.*Australian Mammalogy*, 42, 28–37.
 - Wedrowicz, F., Mosse, J., Wright, W., & Hogan, F. E. (2018). Genetic structure and diversity of the koala population in South Gippsland, Victoria: A remnant population of high conservation significance. *Conservation Genetics*, 19, 713-728.
 - Wedrowicz, F., Wright, W., Schlagloth, R., Santamaria, F., & Cahir, F. (2017). Landscape, koalas and people: A historical account of koala populations and their environment in South Gippsland.*Australian Zoologist*, *38*, 518-536.
 - Whisson, D. A., & Carlyon, K. (2010). Temporal variation in reproductive characteristics of an introduced and abundant Island population of koalas.*Journal of Mammalogy*, *91*, 1160-1167.
 - Whisson, D. A., Dixon, V., Taylor, M. L., & Melzer, A. (2016). Failure to respond to food resource decline has catastrophic consequences for koalas in a high-density population in southern Australia. *PLoS One*, 11, e0144348.
 - Whisson, D. A., Holland, G. J., & Carlyon, K. (2012). Translocation of overabundant species: Implications for translocated individuals. *Journal of Wildlife Management*, 76, 1661-1669.
 - Whisson, D. A., Orlowski, A., & Weston, M. A. (2018). Tree canopy defoliation impacts avifauna. *Forest Ecology and Management*, 428, 81–86.
 - Woinarski, J., & Burbidge, A. A. (2016). Phascolarctos cinere@se IUCN Red List of Threatened Species, 2016: e.T16892A21960344. Retrieved from https://www.iucnredlist.org/species/16892/ 21960344.

How to cite this article: Whisson DA, Ashman KR. When an iconic native animal is overabundant: The koala in southern Australia. *Conservation Science and Practice*. 2020;2:e188. https://doi.org/10.1111/csp2.188