

Associations between sweet taste function, oral complex carbohydrate sensitivity, liking and consumption of *ad libitum* sweet and non-sweet carbohydrate milkshakes among female adults

Julia Y. Q. Low^{1‡}, Kathleen E. Lacy², Robert L. McBride^{1†} and Russell S. J. Keast^{1*}

¹Centre for Advanced Sensory Science, School of Exercise and Nutrition Sciences, Deakin University, Burwood, Victoria 3125, Australia

²Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences, Deakin University, Geelong, Victoria 3220, Australia

(Submitted 21 December 2018 – Final revision received 31 May 2019 – Accepted 3 July 2019)

Abstract

Excess energy intake is recognised as a strong contributing factor to the global rise of being overweight and obese. The aim of this paper was to investigate if oral sensitivity to complex carbohydrate relates to *ad libitum* consumption of complex carbohydrate foods in a sample group of female adults. Participants' (n 51 females): age 23.0 (SD 0.6) years (range 20.0–41.0 years); excluding restrained eaters) sensitivity towards maltodextrin (oral complex carbohydrate) and glucose (sweet taste) was assessed by measuring detection threshold (DT) and suprathreshold intensity perception (ST). A crossover design was used to assess consumption of two different iso-energetic preload milkshakes and *ad libitum* milkshakes – (1) glucose-based milkshake, (2) maltodextrin-based milkshake. *Ad libitum* intake (primary outcome) and eating rate, liking, hunger, fullness and prospective consumption ratings were measured. Participants who were more sensitive towards complex carbohydrate (maltodextrin DT) consumed significantly more maltodextrin-based milkshake in comparison with less-sensitive participants ($P = 0.01$) and this was independent of liking. Participants who had higher liking for glucose-based milkshake consumed significantly more glucose-based milkshake in comparison with participants with lower hedonic ratings ($P = 0.049$). The results provide support regarding the role of the oral system sensitivity (potentially taste) to complex carbohydrate and the prospective to overconsume complex carbohydrate-based milkshake in a single sitting.

Key words: Carbohydrate taste: Liking: Starch taste: Sweet taste: Glucose polymers: Dietary intake

Excess energy intake is recognised as a strong contributing factor to the global rise of being overweight and obese^(1,2). The prevalence of obesity worldwide has been increasing over the past years, necessitating an increased understanding of the drivers of food intake. Foods high in dietary carbohydrates in the form of complex carbohydrates and simple carbohydrates represent a major source of energy in our diet. For example, the estimated Acceptable Macronutrient Distribution Ranges related to reduced risk of chronic disease are 45–65 % of total energy intake from carbohydrate, 20–35 % from fat and 15–25 % from protein⁽³⁾. Foods high in dietary carbohydrate (simple carbohydrate, complex carbohydrate) has been shown to have a weaker effect on satiation in comparison with other food groups such as those high in dietary protein^(4,5) and result in overconsumption within a meal.

Individual differences in their ability to perceive complex carbohydrates and the role of oral complex carbohydrate sensitivity in the overconsumption of energy or specific foods associated with the development of obesity deserve more attention. For example, individuals vary in terms of their satiety responses to dietary fat^(6–8), and one possible explanation may be due to the individual's oral and gastrointestinal sensitivity to fatty acids^(8,9). It has been suggested that abnormalities in any or several taste receptors are known to influence intake of specific food components related to the taste receptor⁽¹⁰⁾. For example, it has been well documented in the literature that individuals' abilities to detect bitter tastants at low concentrations (i.e. *n*-6-propylthiouracil (PROP) and phenylthiocarbamide (PTC)) are determined via genetics⁽¹¹⁾ and influence the palatability and

Abbreviations: DT, detection threshold; gLMS, general Labeled Magnitude Scale; ST, suprathreshold intensity perception.

* **Corresponding author:** Dr R. S. J. Keast, email russell.keast@deakin.edu.au

† Deceased 31 August 2016

‡ Present address: Riddet Institute, Massey University, Palmerston North 4474, New Zealand.

consumption of bitter-tasting vegetables such as kale, broccoli and Brussels sprouts⁽¹⁰⁾. This food choice behaviour has also been reported for orally detected compounds such as fatty acids, whereby a negative relationship between habitual fat intake and oral sensitivity to fatty acids has been found, that is, individuals who were less sensitive to fatty acids were found to consume more fatty foods^(8,12). More recent studies found that oral sensitivity to fatty acids was negatively associated with *ad libitum* intake of high-fat meals (i.e. satiation or intrameal satiety in response to fat⁽⁹⁾) and in subsequent meal intake (i.e. satiety responses to fat⁽⁵⁾). In regards to oral complex carbohydrate sensitivity, a recent cross-sectional study from our laboratory observed a positive association between oral complex carbohydrate sensitivity, intake of complex carbohydrate foods and waist measurements (i.e. being more sensitive to complex carbohydrate was associated with greater energy and starch intakes and a bigger waist measurement)⁽¹³⁾. It is uncertain why we observed an opposite direction in our previous work. However, we speculate that the valence of sensing small amounts of simple carbohydrate (sugar) may promote consumption, and perhaps all carbohydrate sensing may be similarly aligned. Of course, sugars do this via appetitive sweetness, but complex carbohydrates may have an unconscious mode of action on consumption. In this way, sensing all carbohydrates including sugars may promote consumption. However, the relation between habitual diet, body composition and sweet taste sensitivity is complicated because most data showed no relationship between these measures⁽¹⁴⁾, suggesting a need to differentiate between simple and complex carbohydrates. As the previous studies used self-reported dietary measures of habitual/usual intake, it is unclear if the differences in dietary intake were solely due to oral perception as consumption of foods in the real world, being a much less controlled environment than a laboratory, could be influenced by many different factors⁽²⁾. It is therefore important to understand whether oral complex carbohydrate sensitivity influences satiation (i.e. meal size or intrameal satiety) from dietary carbohydrate using an experimental approach in controlled laboratory conditions to look at food or energy intake. If there is an effect of complex carbohydrate on satiation, it is unclear whether individual's liking of complex carbohydrate foods influences this effect as foods with higher palatability could trigger overeating⁽¹⁵⁾.

The aim of this paper was to investigate if oral sensitivity to complex carbohydrate relates to *ad libitum* consumption of complex carbohydrate foods. We assessed this by comparing homogenous milkshakes containing a sweet (glucose) and a non-sweet carbohydrate (maltodextrin). A secondary aim was to investigate if liking of carbohydrate (sweet and non-sweet) foods plays a role in *ad libitum* intake of carbohydrate milkshakes. We hypothesise oral complex carbohydrate sensitivity will be positively associated with *ad libitum* consumption of complex carbohydrate foods. Liking towards carbohydrate foods will be positively associated with *ad libitum* consumption of carbohydrate-based foods. For consistency throughout this paper, the terminology 'oral complex carbohydrate sensitivity' refers to all types of complex carbohydrates and its derivatives, while not diminishing the prospect that oral perception of complex carbohydrate could be due to textural differences⁽¹⁶⁾.




Fig. 1. Number of participants who were recruited, screened and completed both sessions. The dietary restraint score was measured according to factor 1 of the Three-Factor Eating Questionnaire⁽¹⁷⁾. Restrained eaters were defined as participants with a score on factor 1 of >11 on the Three-Factor Eating Questionnaire.

Methods

Study design

Participants consumed two different iso-energetic preload milkshakes followed by *ad libitum* intake of milkshakes – (1) sweet milkshake (glucose) and (2) non-sweet carbohydrate milkshake (maltodextrin) in a randomised crossover design. Maltodextrin was chosen as a complex carbohydrate because it dissolves easily in water, whereas glucose was chosen as a simple carbohydrate/sugar because maltodextrin contains a small amount of glucose. Therefore, by measuring participants' sensitivity towards both glucose and maltodextrin, we were able to observe if differences in the amount of milkshakes consumed were due to sensitivity towards sweet taste (glucose) or oral complex carbohydrate sensitivity (maltodextrin). Participants attended two laboratory sessions, separated by at least 7 d of washout period. The outlines of the two sessions are shown in Fig. 1.

As the sessions were part of a laboratory class, each class (seven participants maximum at a time) was randomly assigned to the sequence of sweet (glucose) and non-sweet (maltodextrin) carbohydrate milkshakes using a web-based program (<http://randomizer.org>). In addition, during the same sessions, detection threshold (DT) and suprathreshold intensity perception (ST) for glucose and maltodextrin, hedonic ratings for glucose and maltodextrin solutions, and hedonic ratings for a range of sweet and complex carbohydrate prototypical foods were also determined. Each session lasted approximately 2 h, and participants were given breaks between tasks lasting 15–30 min. Participants were asked to refrain from eating, drinking (except water) or chewing gum for at least 1 h prior to testing.

Demographic information was also collected, including sex, age, height and weight measurements, during session 1. BMI (kg/m^2) was calculated from the height and weight measurements. Participants also completed two online questionnaires: Likes and Dislikes Questionnaire, and a Three-Factor Eating




Fig. 2. Study outline. The left chart represents the session outline for session 1, middle chart represents the session outline for session 2 and the right chart represents the online questionnaires. Each session lasted about 2 h. As the data collection was part of a laboratory class, participants were given intermittent breaks (teaching) lasting 15–30 min between each task. Participants were also asked to cleanse their palate with deionised water before starting each task during sessions 1 and 2. gLMS, general Labeled Magnitude Scale.

Questionnaire within 1 week of sensory testing. The present study was part of a larger study focusing on the psychophysics of oral complex carbohydrate sensitivity, liking and consumption of complex carbohydrate-based foods⁽¹⁸⁾. Psychophysics tasks (DT, ST), consumption of milkshakes, as well as hedonic ratings for a range of sweet (glucose) and complex carbohydrate (maltodextrin) solutions were conducted in computerised, partitioned sensory booths in the Centre for Advanced Sensory Science using Compusense Cloud Software as part of the Compusense Academic Consortium (Compusense Inc.). Hedonic ratings for a range of sweet and complex carbohydrate-based foods were conducted in individual workbenches at our teaching laboratory. The standardising protocols prior to each testing sessions were similar to the ones outlined in Low *et al.*^(16,19). All solutions and prototypical foods were served at room temperature. Milkshakes were served chilled at approximately 3°C.

Participants

Participants were recruited from a convenience sample of 138 students enrolled in a third-year Sensory Evaluation of Foods unit during March 2016 at Deakin University, Melbourne campus, Australia. A total of 132 participants gave written informed consent to take part in the study (response rate = 96%). The exclusion criteria were included in Fig. 2. A dietary restraint score was measured according to factor 1 of the Three-Factor Eating Questionnaire⁽¹⁷⁾. The mean restraint score was 8.9 (SD 3.7).

Ethics

The present study was approved by the institutional review board regulations of Deakin University (2012_162). The experimental protocol was also registered under the Australian New Zealand Clinical Trials Registry (ACTRN12617000551392; www.anzctr.org.au). The present study also complies with the Declaration of Helsinki for Medical Research involving Human Subjects.

Stimuli and test foods

Glucose was used to investigate sweet taste function (DT and ST for sweet taste; for details of stimuli, see Table 1). Maltodextrin was used to investigate oral complex carbohydrate sensitivity (DT and ST for complex carbohydrate). Detailed in Table 1 are the amount of glucose and total sugars (% w/v) present in each maltodextrin DT concentration.

The nutrient compositions of the sweet (glucose-based) and complex carbohydrate (maltodextrin-based) milkshakes were calculated using the Foodworks8 (Xyris Software) (Table 2). The milkshakes were mixed until no lumps were visible using an immersion (stick) blender for 15 s (per 100 g) at 10 000 rpm (KitchenAid KHB2569 Hand Blender, Whirlpool Corporation). All milkshakes were prepared fresh on the day of testing.

Participant training

At the start of session 1, participants were trained to use the general Labeled Magnitude Scale (gLMS) to rate taste intensity using the standard protocol outlined by Green *et al.*^(21,22), except the top of the scale was described as the strongest imaginable sensation of any kind⁽²³⁾. This method has been described in Low *et al.*^(14,24).

BMI

All participants were asked to remove their shoes and any heavy clothing to ensure accurate measurements. Height and weight were measured right after the scale training during the first session after a 2-h fast (food only). This method has been described in Low *et al.*^(14,16).

Detection threshold determination for sweet taste and oral sensitivity to complex carbohydrates

DT was determined using the procedure outlined in the International Standards Organisation (ISO) Method of



Table 1. Sweetener and complex carbohydrate concentrations used for determination of detection thresholds of healthy female adults (*n* 51)

Stimulus	Concentration (% w/v)								
	1	2	3	4	5	6	7	8	9
Glucose*†	0.05	0.09	0.1	0.2	0.4	0.6	1.1	1.8	2.9
Maltodextrin*†	0.1	0.2	0.3	0.6	1.1	1.9	3.6	6.3	11.2
Amount of glucose in maltodextrin ($\times 10^{-3}$)‡	0.9	1.8	2.7	5.4	9.9	17.1	32.4	56.7	100.8
Amount of total sugars in maltodextrin ($\times 10^{-3}$)‡	1.7	3.4	5.1	10.2	18.7	32.3	61.2	107.1	190.4

* The concentration series for glucose and maltodextrin were prepared with successive 0.25 log dilution steps.

† Reference chemical details: glucose (The Melbourne Food Depot); maltodextrin (Star-Dri 5, Tate & Lyle Ingredients Americas). The ninth concentration was presented only when participants were unable to detect a difference from water solution in the previous eight⁽²⁰⁾.

‡ Calculation of the amount of common and total sugars in maltodextrin concentrations were according to the report of analysis by the Australian Government National Measurement Institute from samples used in the present study, where there were a total of 1.7g/100g (1.7 %, w/w) of free sugars for the maltodextrin (glucose: 0.9 %, w/w; fructose and sucrose: <0.2 %, w/w).

Table 2. Nutrient composition (per 100 g) of sweet (glucose) and non-sweet (maltodextrin) carbohydrate milkshakes containing different amounts of glucose and maltodextrin

	Sweet carbohydrate milkshake*	Non-sweet carbohydrate milkshake*
Energy (kJ)	454.3	440.7
Carbohydrate (g)	13.2	11.8
Sugars (g)	12.8	4.5
Starch (g)	0.4	7.3
Protein (g)	2.8	2.8
Fat (g)	5.3	5.3

* The nutrient composition of the milkshakes (8.8 % (w/w) glucose/maltodextrin (The Melbourne Food Depot; Star-Dri 5, Tate & Lyle Ingredients Americas), 63.7 % (w/w) long-life skimmed milk (99.9 % fat free; Devondale Murray Goulburn), 26.5 % (w/w) light thickened cream (approximately 18 % fat; Bulla) and 1 % (w/w) imitation vanilla essence (Queen Fine Foods)) per 100g was calculated using Foodworks8 (Xyris Software).

Investigating Sensitivity of Taste⁽²⁵⁾. The concentration series for glucose and maltodextrin were prepared with successive 0.25 log dilution steps⁽¹⁴⁾ (Table 1).

The eight samples for each stimulus were served in ascending concentration (15 ml per sample), and each stimulus was presented to participants independently. Participants were unaware of the presentation order. Participants were instructed to taste each sample for 5 s then spit and rate whether there was an absence of taste/oral perception (water-like) or if a taste/oral perception was identified but not recognised⁽²⁵⁾. DT was defined as the concentration at which the participants selected the 'taste/oral perception identified, but unknown taste quality/oral perception'⁽²⁵⁾.

Suprathreshold intensity ratings for glucose and maltodextrin

Three concentrations (weak, medium and strong) and a control (blank) solution were prepared to determine perceived ST for glucose and maltodextrin (Table 3)^(13,24). The concentrations for each stimulus ranged from 'weak' to 'strong' on the gLMS. These samples were presented to participants in a randomised order.

Table 3. Concentrations (weak, medium and strong intensity) of glucose and maltodextrin used for determination of suprathreshold intensity of healthy female adults (*n* 51)

	Concentration (% w/v)		
	Weak	Medium	Strong
Glucose	5.3	10.6	21.2
Maltodextrin	3.6	6.3	11.2
Amount of glucose in maltodextrin ($\times 10^{-3}$)*	32.4	56.7	100.8
Amount of total sugars in maltodextrin ($\times 10^{-3}$)*	61.2	107.1	190.4

* Calculations of the amount of common and total sugars in maltodextrin concentrations were according to the report of analysis by the Australian Government National Measurement Institute from samples used in the present study, where there were a total of 1.7g/100g (1.7 %, w/w) of free sugars for the maltodextrin (glucose: 0.9 %, w/w).

Standardisation of general Labeled Magnitude Scale usage with weight ratings

To standardise gLMS usage within participants, a modified version of the method used by Delwiche *et al.*⁽²⁶⁾ was adapted for the present study (see Low *et al.*^(14,24)). There was a significant correlation between the overall mean sweetness ratings for glucose and overall mean heaviness ratings (r 0.38, P < 0.01) indicating that the gLMS ratings were subject to differences in individual scale-use and thus requires standardisation across participants^(20,26,27). Method to determine standardisation factor for each participants was previously described in Keast & Roper⁽²⁷⁾.

Hedonic ratings for sweet and complex carbohydrate solutions and prototypical foods

To measure liking of glucose and maltodextrin solutions, identical concentrations used to assess ST ratings were prepared and presented to participants in a randomised order. To assess liking of sweet and complex carbohydrate prototypical foods, participants were required to rate liking of sixteen food items (eight sweet taste and eight non-sweet carbohydrate foods). The foods included in testing had approximately equivalent fat per 100 g. Participants were given a variety of different sweet and complex carbohydrate-based foods representing a range of

dietary carbohydrate contents per serve (differences in g of sugar or starch per 100 g), approximately equivalent to the concentrations (% w/v) used to measure ST ratings for glucose and maltodextrin. Eight small samples (5–20 g) per tray were served in a randomised order, and each tray was presented to participants independently. The foods included in testing can be viewed in online Supplementary Table S1. Liking of both solutions and foods was measured using a nine-point hedonic scale. All liking evaluations were conducted without the use of nose clips and following psychophysics tests. All solutions/foods were ingested.

Standardisation of hedonic scale usage with non-food items

To control for idiosyncratic scale usage, standardisation of hedonic scale usage method was previously described⁽¹⁸⁾. There was a significant correlation between the overall mean hedonic ratings for food/beverage items and overall mean hedonic ratings for non-food items ($r = 0.22$, $P < 0.05$). As individual hedonic ratings for food/beverage items and non-food items were assumed unrelated, the significant correlation indicated that the hedonic scale ratings were subject to differences in individual scale-use and required standardisation across participants. Therefore, each individual ratings were standardised with his or her personal standardisation factor to account for hedonic scale-use bias⁽¹⁸⁾.

Satiation measures – preload and *ad libitum* intake of milkshakes, drinking rate, and appetite and hedonic ratings

A modified procedure outlined by Rolls & McDermott⁽²⁸⁾ was used to assess the satiation effect of glucose- and maltodextrin-based milkshakes. Participants were first served a cup containing 200 g of milkshake (glucose: 908.6 kJ, maltodextrin: 881.4 kJ) and were instructed to finish the whole cup of milkshake within a minute (maximum time). At 2 min after consumption of the preload milkshake, participants were presented with another serving of the same milkshake (600 g; glucose: 2725.8 kJ; maltodextrin: 2644.2 kJ). For the 600 g milkshake, participants were told to drink until they are comfortably full (maximum time: 5 min). The serving sizes for preload (200 g) and *ad libitum* (600 g) milkshakes were derived through previously published finding by Rolls & McDermott⁽²⁸⁾ using young adult samples. In that study⁽²⁸⁾, a fixed volume of yogurt (300 g) was given to participants as a preload as it was found to be the average amount of yogurt consumed by participants. However, as the participants in the present study were mainly young female adults, we chose to use 200 g as the serving size for preloads to ensure that all participants were given the opportunity to consume similar amount of milkshakes prior to the *ad libitum* experiment. By standardising the same amount of preload milkshakes, the differences in the amount of milkshakes consumed in the *ad libitum* experiment would be due to the satiating and reward effects of the preload milkshakes. A concentration of 8.8% (per 100 g of maltodextrin milkshake) of maltodextrin was derived based on previous published findings of perceptually distinctive sensation concentration without perceivable viscosity^(29,30).

A concentration of 8.8% (per 100 g of glucose milkshake) of glucose was used for sweet milkshakes. The *ad libitum* milkshake intake was calculated as the difference in the weight of the cup of milkshake before and after consumption. The milkshake intake in g was used to determine the energy intake in kJ. Drinking rate (g/s or kJ/s) was calculated by dividing the *ad libitum* milkshake intake in g or kJ by the total drinking duration (s). During the milkshake experiments, participants were asked to start drinking the milkshake as soon as they were instructed to start. The researcher, using a stopwatch, measured the total duration time (seconds) used to drink the *ad libitum* milkshake.

Prior to consuming the preload and *ad libitum* milkshakes, participants completed several questions relating to appetite and hedonic ratings^(9,31–33). When the milkshakes were served, participants were instructed to drink a sip of their milkshake and to rate their liking of it on a nine-point hedonic scale. Participants were also instructed to rate their feelings of hunger, fullness and prospective consumption prior to consumption of both milkshakes (preload and *ad libitum*) on a 100 mm visual analogue scale anchored at each end with descriptors (e.g. 'not hungry at all' at one end and 'very hungry' at the other).

Statistical analyses

According to previous literature⁽⁹⁾, a difference of 10% in intake (in g) would be detected using forty-nine participants in a paired design with the following assumptions: $\alpha = 0.05$, two-sided, power of 80% and a variation of 25%. Statistical analysis was performed using IBM SPSS statistical software version 23.0 (SPSS). Data are presented as means and standard deviations. Significance was accepted at $P < 0.05$. Descriptive statistics were employed to describe demographic information, thresholds and perceived intensity of sweet taste and oral complex carbohydrate sensitivity, hedonic ratings of sweet taste and complex carbohydrate foods (water-based solutions, prototypical foods and milkshakes), intake of milkshakes (g and kJ), drinking rate, appetite ratings and BMI. Due to low number of male participants, seven males were also eliminated from the data set. Potential confounding variable such as order of presentation (being served a glucose/maltodextrin milkshake first) and BMI on sweet taste function and oral complex carbohydrate sensitivity (DT and ST for glucose and maltodextrin), liking and milkshake intake were checked prior to the analyses using independent *t* tests. The order of presentation and BMI had no effect on liking, milkshake intake, and sweet taste function and oral complex carbohydrate sensitivity (see Results) in this data set.

Participants who are termed more sensitive to the carbohydrate compounds tested have a lower DT and higher intensity ratings than less-sensitive participants (higher DT, lower intensity rating). DT and ST for glucose and maltodextrin were treated as grouping variables (tertiles) with participants categorised as more sensitive/who experienced low intensity (1/3), normal sensitive/moderate intensity (2/3) and less sensitive/high intensity (3/3) to explore differences between continuous (milkshake intake, BMI) variables. DT and ST for glucose and maltodextrin were grouped into tertiles to allow comparison

of most and least sensitive groupings or those groups who experienced low and high intensity (i.e. four sets of tertiles were determined: one for DT for glucose and maltodextrin, and one for ST for glucose and maltodextrin)^(12,14). We used an exploratory approach to allow us to observe a clear indication of any effect the variable may have on other attributes of interest. Similarly, individuals' hedonic ratings for sweet and complex carbohydrate solutions and prototypical foods were treated as grouping variables (tertiles) with participants categorised as those who rated low (1/3), moderate (2/3) and high (3/3) on the hedonic scale to explore differences between variables (milkshake intake). Hedonic ratings for sweet and complex carbohydrate solutions and prototypical foods were grouped into tertiles to allow comparison of those groups who rated low and high on the hedonic scale (i.e. four sets of tertiles were determined for hedonic ratings: for sweet solutions, sweet prototypical foods, complex carbohydrate solutions and complex carbohydrate prototypical foods). An independent *t* test was used to detect differences in milkshake intake between more-sensitive and less-sensitive participants, those who experienced low and high intensity, and those who rated low and high on the hedonic scale groups (low- and high-tertile groups). Pearson's product-moment correlations were conducted to also analyse the relationship between sweet taste function and oral complex carbohydrate sensitivity (DT and ST for glucose and maltodextrin), hedonic ratings for sweet and complex carbohydrate solutions and prototypical foods, milkshakes and BMI. Appetite ratings and hedonic ratings for milkshakes from before compared with after preload within a session were assessed using paired *t* tests. Effects of simple carbohydrate and complex carbohydrate on *ad libitum* milkshake intake, drinking rate and liking of milkshakes were compared using paired *t* tests. The effects of simple carbohydrate and complex carbohydrate on Δ appetite ratings and liking of milkshakes (before *ad libitum* intake – rating before preload intake) were compared using paired *t* tests.

Results

Participants

Of the fifty-one female participants who completed the study (age 23.0 (SD 4.0) years (range 20.0–41.0 years), BMI 22.1 (SD 2.5) kg/m² (range 18.0–29.1 kg/m²), eight were classified as overweight/obese (BMI 26.3 (SD 1.2) kg/m² (range 25.2–29.1 kg/m²)).

Ad libitum intake of glucose and maltodextrin milkshakes

There were no significant differences in *ad libitum* consumption of both glucose and maltodextrin milkshakes (all $P > 0.05$) (Fig. 3(a) and (b)). However, there was a trend towards significance where participants consumed more glucose milkshake in comparison with the maltodextrin milkshake ($P = 0.06$; approximately 23% greater). Similarly, no significant differences between BMI groups (lean and overweight/obese participants) and the order of presentation (presented with glucose milkshakes first *v.* maltodextrin milkshakes) in *ad libitum* consumption of milkshakes were found (all $P > 0.05$).




Fig. 3. *Ad libitum* milkshake intakes by weight (g) (a) and energy (kJ) (b) of healthy female adults (n 51) who consumed sweet (glucose) and non-sweet carbohydrate (maltodextrin) milkshakes in random order. Values are means, with standard deviations represented by vertical bars.

Liking of milkshakes, drinking rate and BMI

Liking ratings of preload and *ad libitum* milkshakes showed a significantly higher liking rating for the glucose milkshake than for the maltodextrin milkshake (all $P < 0.05$) (Fig. 4). Following preload consumption of each milkshake, liking ratings decreased significantly for *ad libitum* milkshake (all $P < 0.05$; Table 4). There were no significant differences between both types of milkshakes on decrease in liking (Δ) ($P = 0.78$). *Ad libitum* drinking rate expressed as g/s and kJ/s did not differ between types of milkshakes (all $P > 0.05$; Table 5).

Ad libitum intakes of both milkshakes were positively correlated with drinking rate (g/s; r 0.75 ($P = 0.001$) and r 0.54 ($P = 0.001$) for the glucose and maltodextrin milkshakes, respectively). No significant correlations were observed between drinking rate (g/s), liking ratings (*ad libitum*) and changes in liking ratings for both types of milkshakes (Δ) (all $P > 0.05$). No significant correlations were observed between BMI and *ad libitum* intake of both milkshakes (all $P > 0.05$). Similarly, BMI was not significantly correlated with intake differences (Δ) of both types of milkshakes (all $P > 0.05$). Drinking rates (g/s), liking ratings (preload and *ad libitum*) and changes in liking ratings for both types of milkshakes (Δ) were not correlated with BMI (all $P > 0.05$).




Fig. 4. Hedonic ratings for preload (■) and *ad libitum* (▒) sweet (glucose) and non-sweet carbohydrate (maltodextrin) milkshakes of healthy female adults (n 51). The y-axis is the adjusted hedonic ratings from a nine-point hedonic scale. The x-axis represents the preload and *ad libitum* milkshakes measured. Values are means, with standard deviations represented by vertical bars.

Appetite ratings

No significant differences were observed between ratings of fullness, hunger and prospective consumption before consumption of preload milkshakes (all $P > 0.05$) signifying that participants were in a similar state of satiety before preload intake. Fullness ratings increased, hunger decreased and ratings of prospective consumption decreased significantly following preload intake of both milkshakes (all $P < 0.001$; Table 4). There were no significant differences in terms of Δ fullness, hunger and ratings of prospective consumption of glucose milkshake in comparison with maltodextrin milkshake (i.e. differences in fullness, hunger and ratings of prospective consumption before and after preload consumption between both milkshakes) (all $P > 0.05$; Table 4).

Sweet taste function, oral complex carbohydrate sensitivity and *ad libitum* intake of milkshakes

The DT and mean intensity ratings, standard deviations and ranges for both glucose and maltodextrin are presented in Table 6; for more details see online Supplementary Fig. S1. Significant negative correlations were identified between maltodextrin DT and *ad libitum* consumption of maltodextrin-based milkshakes ($r = -0.36$, $P = 0.01$). However, no significant correlations were identified between any measures of sweet taste function and *ad libitum* consumption of glucose milkshakes (all $P > 0.05$; Table 7).

When stratified into tertile groups (according to the complex carbohydrate and sweetener tested and all taste measures), we observed significant differences in terms of *ad libitum* consumption of maltodextrin-based milkshakes between the participant groups who were more sensitive and less sensitive towards maltodextrin (DT only) (Fig. 5(a)). Participants who were more sensitive towards maltodextrin (DT) consumed significantly more maltodextrin milkshake (mean intake (g) = 150.1 g; mean intake (kJ) = 695.3 kJ) in comparison with less-sensitive participants

(mean intake (g) = 100.1 g; mean intake (kJ) = 463.1 kJ) ($P = 0.01$). Despite differences in maltodextrin milkshake intake (approximately 50% greater energy intake consumed), no significant changes in appetite ratings (i.e. increase in fullness ratings, decrease in hunger and prospective consumption) were observed between the more-sensitive and less-sensitive participants towards maltodextrin DT (all $P > 0.05$). There were no significant differences in terms of *ad libitum* consumption of glucose-based milkshakes between the participant groups who were more sensitive and less sensitive towards glucose (DT) (Fig. 5(a)). Similarly, no significant differences in terms of *ad libitum* consumption of both glucose and maltodextrin-based milkshakes between the participant groups who experienced low intensity or high intensity to glucose (ST) and maltodextrin (ST) (all $P > 0.05$; Fig. 5(b)).

Hedonic ratings for glucose and maltodextrin solutions, prototypical foods, milkshakes and *ad libitum* intake of milkshakes

No significant differences in hedonic ratings for glucose and maltodextrin solutions and prototypical foods were identified between BMI and order of session groups (all $P > 0.05$). The mean hedonic ratings, standard deviations and ranges for both sweet and complex carbohydrate solutions (glucose, maltodextrin) and prototypical foods are presented in Table 8. Liking of glucose milkshake was significantly correlated with glucose milkshake intake ($r = 0.30$; $P = 0.03$). No significant associations were observed between other sweet and complex carbohydrate hedonic measures (solutions, prototypical foods) and *ad libitum* intake of milkshakes (all $P > 0.05$).

When stratified into tertile groups (according to the complex carbohydrate and sweetener tested and all taste measures), we observed significant differences in terms of hedonic ratings for both preload and *ad libitum* glucose-based milkshakes between the participant groups who were more sensitive and less sensitive towards glucose (DT, ST). Participants who were more sensitive towards glucose solutions (DT, ST) rated higher on the hedonic scale for both preload and *ad libitum* glucose milkshakes (DT/ST: mean preload hedonic ratings = 6.6/6.5; mean *ad libitum* hedonic ratings = 6.0/6.0) in comparison with participants with lower hedonic ratings (mean preload hedonic ratings = 5.1/5.5; mean *ad libitum* hedonic ratings = 4.2/4.5) (all $P < 0.05$). However, there were no significant differences in hedonic ratings for glucose milkshakes between oral sensitivity groups towards maltodextrin (both DT, ST) (all $P > 0.05$). Similarly, there were no significant differences in hedonic ratings for maltodextrin milkshakes between more-sensitive and less-sensitive participants or those who experienced high intensity or low intensity to both glucose and maltodextrin (DT, ST) (all $P > 0.05$).

When stratified into tertile groups (according to liking ratings towards solutions, prototypical foods and milkshakes), we observed a trend ($P = 0.09$) towards significant differences in terms of *ad libitum* consumption of glucose milkshakes between participants with high hedonic ratings and low hedonic ratings for glucose solutions (Fig. 5(c)). Significant differences were observed between those with high hedonic ratings and low

Table 4. Hedonic ratings and appetite ratings of healthy female adults (*n* 51) who consumed two types of milkshakes containing different amounts of glucose (sweet carbohydrate milkshake) and maltodextrin (non-sweet carbohydrate milkshake) on two separate days* (Mean values and standard deviations)

	Sweet carbohydrate milkshake		<i>P</i> †	Non-sweet carbohydrate milkshake		<i>P</i> †	<i>P</i> ‡	<i>P</i> §
	Mean	SD		Mean	SD			
Hedonic								
Before preload intake	5.9	1.9		5.0	2.0		<0.01	
Before <i>ad libitum</i> intake	5.1	2.0	<0.001	4.2	1.8	<0.01		
Δ	−0.8	1.8		−0.9	1.4			0.79
Hunger (mm)								
Before preload intake	59.7	24.5		54.4	22.9		0.08	
Before <i>ad libitum</i> intake	40.0	21.8	<0.001	34.1	23.9	<0.001		
Δ	−23.8	19.5		−20.3	19.3			0.26
Fullness (mm)								
Before preload intake	24.4	20.5		25.6	19.8		0.74	
Before <i>ad libitum</i> intake	55.8	22.0	<0.001	56.4	23.0	<0.001		
Δ	31.4	22.0		30.8	21.8			0.89
Prospective consumption (mm)								
Before preload intake	61.0	23.6		57.9	19.6		0.34	
Before <i>ad libitum</i> intake	39.1	23.4	<0.001	37.2	23.1	<0.001		
Δ	−22.0	18.7		−20.8	17.8			0.69

* Δ: rating before *ad libitum* intake – rating before preload intake. Hedonic values are adjusted hedonic ratings from a nine-point hedonic scale.

† *P* values representing differences between before preload intake and before *ad libitum* intake in hedonic, hunger, fullness and prospective consumption ratings (paired *t* tests).

‡ *P* values representing differences between sweet and non-sweet carbohydrate milkshake sessions before preload intake in hedonic, hunger, fullness and prospective consumption ratings (paired *t* tests).

§ *P* values representing differences between sweet and non-sweet carbohydrate milkshakes in terms of changes before preload intake and before *ad libitum* intake (Δ) in hedonic, hunger, fullness and prospective consumption ratings (paired *t* tests).

Table 5. Drinking rates and meal durations of healthy female adults (*n* 51) for *ad libitum* consumption of two types of milkshakes containing different amounts of glucose (sweet milkshake) and maltodextrin (non-sweet carbohydrate milkshake) (Mean values and standard deviations)

	Sweet milkshake		Non-sweet carbohydrate milkshake		<i>P</i> *
	Mean	SD	Mean	SD	
Drinking rate (g/s)	3.8	1.9	3.2	1.6	0.08
Energy intake rate (kJ/s)	17.0	8.9	14.7	7.6	0.07
Meal duration (s)	51.6	47.3	51.0	41.8	0.93

* *P* values representing differences between sweet milkshake and non-sweet carbohydrate milkshake (paired *t* tests).

hedonic ratings for glucose milkshakes and *ad libitum* consumption of glucose milkshakes (*P* = 0.049) (Fig. 6). Participants who had high hedonic ratings for the glucose-based milkshake consumed significantly more for the same milkshake (mean intake (g) = 212.9 g; mean intake (kJ) = 946.3 kJ) in comparison with participants with low hedonic ratings (mean intake (g) = 123.9 g; mean intake (kJ) = 550.8 kJ). There were no significant differences between participants who rated low and high on the hedonic scale according to their liking towards sweet (prototypical foods) and complex carbohydrate hedonic ratings (solutions, prototypical foods, milkshake) for *ad libitum* consumption of milkshakes (all *P* > 0.05; Figs. 5(d) and 6).

Discussion

To our understanding, the present study is the first to investigate if individuals' ability to detect and perceive complex carbohydrates at a range of concentrations is associated with *ad libitum*

intake of energy/foods in the form of liquid. The major finding was that those who were able to detect complex carbohydrate in water at a lower concentration (DT, more-sensitive group) consumed 50 % more non-sweet carbohydrate (maltodextrin-based) milkshake than of those who were less sensitive to complex carbohydrate, and this was independent of liking. Despite differences in intake of non-sweet carbohydrate (maltodextrin-based) milkshake, there were no significant changes in appetite ratings (i.e. decrease in hunger and prospective consumption, increase in fullness) between those who were more sensitive and less sensitive to complex carbohydrate (DT). Presumably liking was driving consumption of the glucose milkshake; however, liking of maltodextrin was not driving consumption of the maltodextrin-based milkshake. In fact, we observed that the maltodextrin-based milkshakes were not liked by participants at all (i.e. average mean hedonic ratings went from neutral in the preload to dislike in the *ad libitum* milkshake). It appears that maltodextrin sensitivity (DT) is associated with increasing consumption although the mechanism remains unknown. We speculate that sensing small amounts of complex carbohydrates (maltodextrin) may promote unconscious consumption due to the activation of specific brain regions involved with taste and reward. For example, previous neuroimaging sequence studies using fMRI to assess cortical responses to a maltodextrin mouth rinse revealed activation within the primary taste cortex and the neural networks (reward) associated with sensory perception^(34,35). All in, these data suggest a novel role of the oral perceptual system to complex carbohydrates in regards to the overconsumption of energy within a meal.

In the present study, participants who were more sensitive to complex carbohydrate (DT) consumed more of the non-sweet carbohydrate milkshake, thus energy intake, than less-sensitive participants did. These findings provide strong evidence to

Table 6. Detection threshold (DT) (% w/v) and mean intensity ratings (general Labeled Magnitude Scale) for glucose and maltodextrin of healthy female adults (*n* 51)* (Mean values, standard deviations and ranges)

	DT				Mean intensity rating			
	<i>n</i>	Mean	SD	Range	<i>n</i>	Mean	SD	Range
Glucose	51	1.0	0.7	0.05–1.8	51	21.3	5.8	10.8–42.7
Lower tertile groups (1/3)	17	0.2	0.1	0.05–0.2	17	15.6	2.0	10.8–18.9
Upper tertile groups (3/3)	17	1.8	0.00	1.8–1.8	17	27.6	4.9	22.7–42.7
Maltodextrin	51	3.2	2.6	0.1–6.3	51	18.1	6.7	8.2–38.5
Lower tertile groups (1/3)	17	0.3	0.2	0.1–0.6	17	11.3	1.7	8.2–14.3
Upper tertile groups (3/3)	18	6.3	0.00	6.3–6.3	17	20.3	5.3	20.3–38.5

* Mean intensity ratings calculated based on the geometric mean score of the three solution ratings (weak, moderate and strong). Participants who are termed more sensitive to the carbohydrate compounds tested have a lower DT and higher intensity ratings than less-sensitive participants (higher DT, lower intensity rating). DT and suprathreshold intensity perception for glucose and maltodextrin were treated as grouping variables (tertiles) with participants categorised as more sensitive/who experienced low intensity (1/3), normal sensitive/moderate intensity (2/3) and less sensitive/high intensity (3/3) to explore differences between continuous (milkshake intake, BMI) variables. DT and suprathreshold intensity perception for glucose and maltodextrin were grouped into tertiles to allow comparison of most and least sensitive groupings or those groups who experienced low and high intensity.

Table 7. Pearson product-moment correlations between detection thresholds, mean intensity ratings and *ad libitum* milkshakes for glucose and maltodextrin of healthy female adults (*n* 51)† (Pearson's *r* correlation coefficient values)

	Sweet (glucose) taste function		Complex carbohydrate (maltodextrin) oral sensitivity	
	Glucose detection thresholds	Glucose intensity ratings	Maltodextrin detection thresholds	Maltodextrin intensity ratings
Sweet (glucose) milkshake intake	–0.20	0.16	–	–
Non-sweet (maltodextrin) milkshake intake	–	–	–0.36*	0.05

* *P* = 0.01.

† For sweet (glucose) detection thresholds and mean intensity ratings, statistical relationships were only calculated for sweet (glucose) milkshakes, and *vice versa* for complex carbohydrate (maltodextrin).

Table 8. Hedonic ratings for sweet and complex carbohydrate solutions and prototypical foods of healthy female adults (*n* 51) (Mean values, standard deviations and ranges)

	Solutions (<i>n</i> 51)*			Prototypical foods (<i>n</i> 51)†		
	Mean	SD	Range	Mean	SD	Range
Sweet	4.8	1.8	2.0–9.6	6.0	1.4	3.0–11.3
Complex carbohydrate	3.2	1.5	0.9–7.8	5.8	1.7	3.2–11.6

* Hedonic rating for solutions calculated based on the geometric mean score of the three solution ratings (weak, medium and strong).

† For hedonic ratings of a range of sweet and complex carbohydrate foods, a geometric mean score of the eight food items was used.

support our first hypothesis where oral complex carbohydrate sensitivity will be positively associated with *ad libitum* consumption of complex carbohydrate foods. Furthermore, they were in line with our previous studies showing that oral sensitivity to complex carbohydrate is positively associated with habitual energy intake and intake of dietary starch⁽¹⁶⁾. Interestingly, these were only observed between DT measures (not intensity perception) and *ad libitum* consumption of milkshakes. As there are multiple perceptual phases of taste with no single measure being able to represent taste function globally⁽²⁰⁾, this emphasises the need to include more than one measure of the oral function to measure the complex relationship between perception and dietary intake.

In the present work, we found that those who rated higher on the hedonic scale for sweet milkshakes had greater consumption of the sweet milkshakes. In contrast, despite an increase in consumption between participants who were more sensitive to complex carbohydrate, no significant associations were found between complex carbohydrate liking, appetite ratings and *ad libitum* intake of the non-sweet carbohydrate milkshakes. This provides a partial support for our second hypothesis, whereby only liking towards sweetness was associated

with *ad libitum* consumption of sweet-carbohydrate-based foods. More importantly, our previous work also showed positive association between oral sensitivity to complex carbohydrate and waist circumference measurements⁽¹⁶⁾. This suggests a possibility of some sub-conscious mechanism relating to oral sensitivity to complex carbohydrates (longer-term outcome of sensitivity), but not conscious liking that encourages consumption.

Regarding sweet taste function, there were no significant differences between participants who were more sensitive and less sensitive or those who experienced low and high intensity to sweet taste (both DT, ST) and *ad libitum* intake of sweet milkshakes. Although some studies found significant associations between sweet taste function, BMI and dietary intake^(36–38), this is in line with a larger body of evidence indicating no significant associations^(14,39–47). In addition, we found that the participants who had higher liking ratings for sweet milkshakes consumed more of the *ad libitum* sweet milkshake than the participants who had lower liking. It is likely that no significant associations were found between sweetness liking, BMI and intake of sweet foods in the previous studies^(41,43,45,46,48–51), as most of these studies looked at self-reported habitual or usual intake rather than satiation/acute intake in a controlled laboratory environment. By looking at satiation/acute intake in a controlled environment, we were able to observe significant differences between hedonic ratings and *ad libitum* consumption of sweet milkshakes, in comparison with other measures such as solution and prototypical foods liking ratings suggesting that satiation measures are




Fig. 5. (a,b) *Ad libitum* milkshake intakes of more-sensitive (■) and less-sensitive (■) participants or those who experienced high (■) and low (■) intensity ratings. (c,d) *Ad libitum* milkshake intakes of participants with high hedonic ratings (■) and low hedonic ratings (■) for both sweet and complex carbohydrate solutions and prototypical foods. For sweet taste function and sweet hedonic ratings, comparisons were only made for sweet (glucose) milkshakes, and *vice versa* for complex carbohydrate (maltodextrin). Values are means, with standard deviations represented by vertical bars. * $P=0.01$. DT, detection threshold.




Fig. 6. *Ad libitum* milkshake intakes for participants with high hedonic ratings (■) and low hedonic ratings (■) for both sweet (glucose) and non-sweet (maltodextrin) carbohydrate milkshakes. For sweet hedonic ratings, comparisons were only made for sweet milkshakes, and *vice versa* for complex carbohydrate. Values are means, with standard deviations represented by vertical bars. * $P=0.049$.

more appropriate/sensitive. Furthermore, it is also possible that no associations were observed between liking and consumption of sweet foods in the previously mentioned studies, as foods high in dietary sugar are most likely accompanied by other taste qualities such as salty, sour, bitter and fatty tastes. Thus, by matching both sweet and non-sweet carbohydrate milkshakes in energy, serving size, protein, fat, as well as salt and fibre levels, we were able to observe the direct influence of liking of sweetness on intake of a sweet milkshake within a meal. Therefore,

foods high in dietary sugar may be one of the many risk factors for overconsumption of energy for individuals with high liking for sweetness due to the sweet taste or flavours present.

The present study needs to be considered alongside limitations, which may have confounded the results. First, the present study measured the intake of only a single food (milkshake). Although laboratory setting research using single foods is the most sensitive approach and provides clear results when quantifying the role of sensory properties on food intake, in reality, however, we normally consume multiple foods in a much less controlled environment as well as foods that consist of a more complex flavour⁽²⁾. Therefore, it is difficult to extrapolate these findings to everyday life. Second, we did not measure appetite ratings and liking ratings after consumption of the *ad libitum* milkshakes, or appetite ratings and intake of other foods in subsequent hours following the milkshake experiment as this was beyond the scope of the present study. Last, but not least, the participants were mainly young female adults within the normal BMI range, thus the present findings may be difficult to generalise to the broader population.

Conclusions

Participants who were orally more sensitive to complex carbohydrate (DT) consumed 50 % more of the non-sweet carbohydrate milkshake than those who were less sensitive to complex carbohydrate and this was independent of liking. However, no relationships were observed between sweet taste

function and *ad libitum* intake of sweet milkshakes. For sweet taste, the present study showed that those who had higher liking ratings for sweet milkshake consumed significantly more sweet milkshakes in comparison with those who had lower liking ratings and this was independent of taste sensitivity to glucose. All in, these results support an association between oral (may be taste system) sensitivity to complex carbohydrate and the potential to overconsume complex carbohydrate foods. The present findings also provide insights into the relationship between liking of sweet taste and the potential to overconsume sweet foods within a meal.

Acknowledgements

Our friend and colleague Dr R. L. McBride passed away during preparation of the paper after a short illness.

The present study was supported by the Centre for Advanced Sensory Science, School of Exercise and Nutrition Sciences of Deakin University, Victoria, Australia.

J. Y. Q. L., K. E. L., R. L. M. and R. S. J. K. designed the research; J. Y. Q. L. conducted the research, analysed the data and wrote the main part of the manuscript; K. E. L., R. L. M. and R. S. J. K. provided expert input and were involved with the drafting of the manuscript; and R. S. J. K. had primary responsibility for final content. J. Y. Q. L., K. E. L. and R. S. J. K. read and approved the final manuscript.

All authors disclose no conflicts of interest.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S0007114519001703>

References

1. Swinburn BA, Sacks G, Hall KD, *et al.* (2011) The global obesity pandemic: shaped by global drivers and local environments. *Lancet* **378**, 804–814.
2. Drewnowski A & Bellisle F (2007) Liquid calories, sugar, and body weight. *Am J Clin Nutr* **85**, 651–661.
3. National Health and Medical Research Council, Australian Government Department of Health and Ageing, New Zealand Ministry of Health (2006) *Nutrient Reference Values for Australian and New Zealand Including Recommended Dietary Intakes*. Canberra: National Health and Medical Research Council. <https://www.nhmrc.gov.au/sites/default/files/images/nutrient-reference-dietary-intakes.pdf> (accessed September 2019).
4. Bertenshaw EJ, Luch A & Yeomans MR (2008) Satiating effects of protein but not carbohydrate consumed in a between-meal beverage context. *Physiol Behav* **93**, 427–436.
5. Keast RS, Azzopardi KM, Newman LP, *et al.* (2014) Impaired oral fatty acid chemoreception is associated with acute excess energy consumption. *Appetite* **80**, 1–6.
6. Blundell JE, Burley V, Cotton J, *et al.* (1993) Dietary fat and the control of energy intake: evaluating the effects of fat on meal size and postmeal satiety. *Am J Clin Nutr* **57**, 772S–777S.
7. Blundell JE, Stubbs R, Golding C, *et al.* (2005) Resistance and susceptibility to weight gain: individual variability in response to a high-fat diet. *Physiol Behav* **86**, 614–622.
8. Stewart JE, Seimon RV, Otto B, *et al.* (2011) Marked differences in gustatory and gastrointestinal sensitivity to oleic acid between lean and obese men. *Am J Clin Nutr* **93**, 703–711.
9. Bolhuis DP, Costanzo A, Newman LP, *et al.* (2016) Salt promotes passive overconsumption of dietary fat in humans. *J Nutr* **146**, 838–845.
10. Dinehart M, Hayes J, Bartoshuk L, *et al.* (2006) Bitter taste markers explain variability in vegetable sweetness, bitterness, and intake. *Physiol Behav* **87**, 304–313.
11. Guo S-W & Reed DR (2001) The genetics of phenylthiocarbamide perception. *Ann Human Biol* **28**, 111–142.
12. Stewart JE, Feinle-Bisset C, Golding M, *et al.* (2010) Oral sensitivity to fatty acids, food consumption and BMI in human subjects. *Br J Nutr* **104**, 145–152.
13. Low JY, Lacy KE, McBride RL, *et al.* (2017) Carbohydrate taste sensitivity is associated with starch intake and waist circumference in adults. *J Nutr* **147**, 2235–2242.
14. Low JY, Lacy KE, McBride R, *et al.* (2016) The association between sweet taste function, anthropometry, and dietary intake in adults. *Nutrients* **8**, 241.
15. Drewnowski A, Mennella JA, Johnson SL, *et al.* (2012) Sweetness and food preference. *J Nutr* **142**, 1142S–1148S.
16. Low JY, Lacy KE, McBride RL, *et al.* (2017) Evidence supporting oral sensitivity to complex carbohydrates independent of sweet taste sensitivity in humans. *PLOS ONE* **12**, e0188784.
17. Stunkard AJ & Messick S (1985) The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger. *J Psychosomat Res* **29**, 71–83.
18. Low JY, Lacy KE, McBride RL, *et al.* (2018) The associations between oral complex carbohydrate sensitivity, BMI, liking, and consumption of complex carbohydrate based foods. *J Food Sci* **83**, 2227–2236.
19. Low JY, McBride RL, Lacy KE, *et al.* (2017). Psychophysical evaluation of sweetness functions across multiple sweeteners. *Chem Senses* **42**, 111–120.
20. Webb J, Bolhuis DP, Cicerale S, *et al.* (2015) The relationships between common measurements of taste function. *Chemosens Percept* **8**, 11–18.
21. Green BG, Shaffer GS & Gilmore MM (1993) Derivation and evaluation of a semantic scale of oral sensation magnitude with apparent ratio properties. *Chem Senses* **18**, 683–702.
22. Green BG, Dalton P, Cowart B, *et al.* (1996) Evaluating the 'Labeled Magnitude Scale' for measuring sensations of taste and smell. *Chem Senses* **21**, 323–334.
23. Bartoshuk LM (2000) Comparing sensory experiences across individuals: recent psychophysical advances illuminate genetic variation in taste perception. *Chem Senses* **25**, 447–460.
24. Low JY, McBride RL, Lacy KE, *et al.* (2016) Psychophysical evaluation of sweetness functions across multiple sweeteners. *Chem Senses* **42**, 111–120.
25. ISO3972 (1991) *Sensory Analysis-Methodology-Method of Investigating Sensitivity of Taste*. Geneva: International Organisation for Standardization.
26. Delwiche JF, Buletic Z & Breslin PA (2001) Relationship of papillae number to bitter intensity of quinine and PROP within and between individuals. *Physiol Behav* **74**, 329–337.
27. Keast RS & Roper J (2007) A complex relationship among chemical concentration, detection threshold, and suprathreshold intensity of bitter compounds. *Chem Senses* **32**, 245–253.
28. Rolls BJ & McDermott TM (1991) Effects of age on sensory-specific satiety. *Am J Clin Nutr* **54**, 988–996.
29. de Ataíde e Silva T, Di Cavalcanti Alves de Souza ME, de Amorim JF, *et al.* (2013) Can carbohydrate mouth rinse improve performance during exercise? A systematic review. *Nutrients* **6**, 1–10.

30. Lapis TJ, Penner MH & Lim J (2014) Evidence that humans can taste glucose polymers. *Chem Senses* **39**, 737–747.
31. Rolls BJ, Rolls ET, Rowe EA, *et al.* (1981) Sensory specific satiety in man. *Physiol Behav* **27**, 137–142.
32. Rolls BJ, Hetherington M & Burley VJ (1988) The specificity of satiety: the influence of foods of different macronutrient content on the development of satiety. *Physiol Behav* **43**, 145–153.
33. Havermans RC, Janssen T, Giesen JC, *et al.* (2009) Food liking, food wanting, and sensory-specific satiety. *Appetite* **52**, 222–225.
34. Turner CE, Byblow WD, Stinear CM, *et al.* (2014) Carbohydrate in the mouth enhances activation of brain circuitry involved in motor performance and sensory perception. *Appetite* **80**, 212–219.
35. Chambers E, Bridge M & Jones D (2009) Carbohydrate sensing in the human mouth: effects on exercise performance and brain activity. *J Physiol* **587**, 1779–1794.
36. Jayasinghe S, Kruger R, Walsh D, *et al.* (2017) Is sweet taste perception associated with sweet food liking and intake? *Nutrients* **9**, 750.
37. Proserpio C, Laureati M, Bertoli S, *et al.* (2015) Determinants of obesity in Italian adults: the role of taste sensitivity, food liking, and food neophobia. *Chem Senses* **41**, 169–176.
38. Bartoshuk LM, Duffy VB, Hayes JE, *et al.* (2006) Psychophysics of sweet and fat perception in obesity: problems, solutions and new perspectives. *Philos Trans R Soc Lond B Biol Sci* **361**, 1137–1148.
39. Ciceralo S, Riddell LJ & Keast RS (2012) The association between perceived sweetness intensity and dietary intake in young adults. *J Food Sci* **77**, H31–H35.
40. Grinker J, Hirsch J & Smith DV (1972) Taste sensitivity and susceptibility to external influence in obese and normal weight subjects. *J Pers Soc Psychol* **22**, 320.
41. Malcolm R, O'Neil P, Hirsch A, *et al.* (1979) Taste hedonics and thresholds in obesity. *Int J Obes* **4**, 203–212.
42. Frijters JE & Rasmussen-Conrad EL (1982) Sensory discrimination, intensity perception, and affective judgment of sucrose-sweetness in the overweight. *J Gen Psychol* **107**, 233–247.
43. Wooley OW, Wooley SC & Dunham RB (1972) Calories and sweet taste: effects on sucrose preference in the obese and non-obese. *Physiol Behav* **9**, 765–768.
44. Rodin J, Moskowitz HR & Bray GA (1976) Relationship between obesity, weight loss, and taste responsiveness. *Physiol Behav* **17**, 591–597.
45. Rodin J (1975) Effects of obesity and set point on taste responsiveness and ingestion in humans. *J Comp Physiol Psychol* **89**, 1003.
46. Thompson DA, Moskowitz HR & Campbell RG (1977) Taste and olfaction in human obesity. *Physiol Behav* **19**, 335–337.
47. Bertoli S, Laureati M, Battezzati A, *et al.* (2014) Taste sensitivity, nutritional status and metabolic syndrome: implication in weight loss dietary interventions. *World J Diabetes* **5**, 717.
48. Thompson DA, Moskowitz HR & Campbell RG (1976) Effects of body weight and food intake on pleasantness ratings for a sweet stimulus. *J Appl Physiol* **41**, 77–83.
49. Simone M & Pangborn R (1957) Consumer acceptance methodology-one vs 2 samples. *Food Technol* **11**, A25–A29.
50. Witherly S, Pangborn RM & Stern JS (1980) Gustatory responses and eating duration of obese and lean adults. *Appetite* **1**, 53–63.
51. Drewnowski A, Kurth CL & Rahaim JE (1991) Taste preferences in human obesity: environmental and familial factors. *Am J Clin Nutr* **54**, 635–641.