

The Official Journal of the American College of Sports Medicine

. . . Published ahead of Print

Extended Sleep Maintains Endurance Performance better than Normal or Restricted Sleep

Spencer S H Roberts¹, Wei-Peng Teo^{1,2}, Brad Aisbett¹, Stuart A Warmington¹

¹Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences. Deakin University, Burwood, Victoria, Australia; ²Physical Education and Sports Science Academic Group, National Institute of Education, Nanyang University, Singapore

Accepted for Publication: 9 June 2019

Medicine & Science in Sports & Exercise Published ahead of Print contains articles in unedited manuscript form that have been peer reviewed and accepted for publication. This manuscript will undergo copyediting, page composition, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered that could affect the content.

Extended Sleep Maintains Endurance Performance better than

Normal or Restricted Sleep

Spencer S H Roberts¹, Wei-Peng Teo^{1,2}, Brad Aisbett¹, Stuart A Warmington¹

¹Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences. Deakin University, Burwood, Victoria, Australia; ²Physical Education and Sports Science Academic Group, National Institute of Education, Nanyang University, Singapore

Corresponding author:

Spencer S. H. Roberts, PhD Candidate

School of Exercise and Nutrition Sciences, Deakin University.221 Burwood Hwy, Burwood, Victoria, Australia, 3125

Email: rspen@deakin.edu.au

The results of this study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. Results do not constitute endorsement by the American College of Sports Medicine. Spencer Roberts, Wei-Peng Teo, Brad Aisbett, and Stuart Warmington declare they have no conflict of interest. The authors acknowledge the School of Exercise and Nutrition Sciences at Deakin University for funding the study.

ABSTRACT

Purpose: The cumulative influence of sleep time on endurance performance remains unclear. This study examined effects of three consecutive nights of both sleep extension and restriction on endurance cycling performance. Methods: Endurance cyclists/triathletes (n=9) completed a counterbalanced crossover experiment with three conditions; sleep restriction (SR), normal sleep (NS), and sleep extension (SE). Each condition comprised seven days/nights of data collection (-2, -1, D1, D2, D3, D4, +1). Sleep was monitored using actigraphy throughout. Participants completed testing sessions on days D1-D4 that included an endurance time-trial (TT), mood, and psychomotor vigilance assessment. Perceived exertion (RPE) was monitored throughout each TT. Participants slept habitually prior to D1, however, time in bed was reduced by 30% (SR), remained normal (NS), or extended by 30% (SE) on nights D1, D2, and D3. Data were analysed using Generalised Estimating Equations. Results: On nights D1, D2, and D3, total sleep time was longer (P<0.001) in the SE condition (8.6±1.0; 8.3±0.6; 8.2±0.6h, respectively), and shorter (P<0.001) in the SR condition (4.7±0.8; 4.8±0.8; 4.9±0.4h) compared with NS (7.1±0.8; 6.5±1.0; 6.9±0.7h). Compared with NS, TT performance was slower (P<0.02) on D3 of SR (58.8±2.5 vs 60.4 ± 3.7 min) and faster (P<0.02) on D4 of SE (58.7 ±3.4 vs 56.8 ±3.1 min). RPE was not different between or within conditions. Compared with NS, mood disturbance was higher-, and psychomotor vigilance impaired, following SR. Compared with NS, psychomotor vigilance improved following SE. Conclusion: Sleep extension for three nights led to better maintenance of endurance performance compared with normal and restricted sleep. Sleep restriction impaired performance. Cumulative sleep time affects performance by altering the perceived exertion of a given exercise intensity. Endurance athletes should sleep >8 hours per night to optimise performance. **Keywords:** Recovery, fatigue, athlete. extra sleep, sports

INTRODUCTION

Endurance athletes experience high levels of physical and psychological stress during training and competition (1). For example, elite road cyclists pedal more than 30,000 km per year, and during stage races, will compete for 4-6 hours per day on consecutive days (1). Sleep is considered an important recovery behaviour that may help athletes tolerate such demands (2), however, the influence of sleep on endurance performance remains unclear.

No study, to our knowledge, has investigated the effects of sleep extension (i.e., increased habitual total sleep time) on endurance performance. In non-endurance athletes, sleep extension has been reported to improve the serving accuracy of tennis players (3), and the shooting accuracy and sprint times of basketballers (4). However, in the latter study, the absence of a control arm suggests improvements may have been attributable to training adaptations rather than sleep extension (4). Studies investigating effects of sleep *restriction* (i.e., decreased habitual total sleep time) on endurance performance have reported equivocal findings (5-10). Moreover, these studies have often recruited untrained participants (5, 6, 9), assessed performance using relatively brief (<30 minutes) intermittent (5, 6) or graded exercise (9) tests, or examined effects of a single night of sleep restriction (5, 6, 8, 10). Given many endurance athletes (e.g., road cyclists) train or compete for prolonged periods (≥ 60 minutes), and on consecutive days, and in light of evidence that athletes' sleep is often disturbed during training and competition (11), further investigation of the cumulative effects of sleep time on endurance performance is required. The present study examined the effects of both sleep extension and restriction across three consecutive nights endurance cycling performance. on

METHODS

Participants

Nine males (mean \pm SD, Age: 30 \pm 6 years, $\dot{V}O_{2 max}$: 63 \pm 6 mL·kg⁻¹·min⁻¹) were recruited from cycling (n=7) and triathlon (n=2) clubs. Athletes were considered 'trained' according to adapted criteria for classifying cyclists (\geq 1 year competitive racing, \geq 3 training sessions per week, $\dot{V}O_{2 max} \geq$ 55 mL·kg⁻¹·min⁻¹) (12). To screen for sleep problems and high anxiety, inclusion criteria required a score \leq 5 in the Pittsburgh Sleep Quality Index (PSQI) (13), and \leq 40 in the State-Trait Anxiety Inventory (STAI-Y) (14). Participants did not habitually consume high levels of caffeine (mean \pm SD, caffeine products per day: 2 \pm 1). The Morningness-Eveningness Questionnaire determined participants were mostly 'moderate morning' types (n=5), with the remainder being 'definite morning' (n=2) or 'intermediate' (n=2) types (15). The study was approved the Deakin University Human Research Ethics Committee and informed consent was obtained prior to participation.

Overview

Participants completed a counterbalanced crossover experiment with three conditions; sleep restriction (SR), normal sleep (NS), and sleep extension (SE). Beforehand, participants had their habitual sleep monitored for a minimum four nights, and undertook two familiarisation sessions comprising an incremental exercise test and a practice time-trial (TT), respectively. Each condition comprised seven consecutive days/nights (-2, -1, D1, D2, D3, D4, +1) of data collection (Figure 1). Participants undertook four testing sessions (D1-D4) at the Deakin University Human Research and Performance Laboratory. During these sessions, participants completed an endurance TT, subjective mood evaluation, and a psychomotor vigilance task. For all conditions, participants slept habitually prior to D1. However, for the three subsequent

'intervention' nights (D1, D2, D3), habitual 'time in bed' was either reduced by 30% (SR), extended by 30% (SE), or remained normal (NS). Required time in bed for the intervention nights was calculated according to participants' habitual sleep recorded prior to the experiment. Participants were prescribed bed-, and get-up times on nights D1, D2, and D3 to ensure the required time in bed was achieved. Bed- and get-up times were tailored to individual chorotype in order to maximise the likelihood of modifying 'total sleep time' rather than simply 'time in bed'. For example, sleep extension for a 'morning type' was prescribed by predominantly advancing bedtime rather than delaying get-up time. To minimise the effect of circadian variations on performance, all testing commenced between 6:00-9:00 am. Testing start times were consistent for each participant on D1 of each condition, and on D2, D3, and D4 of the NS condition (mean \pm SD, start-time, 7:08 am \pm 31 min). Testing start times on D2, D3, and D4 of the SE condition were slightly later to allow for prescribed time in bed increases (mean \pm SD, start time, 7:48 am \pm 37 min). Testing start times on D2, D3, and D4 of the SR condition were slightly earlier to reduce idle time after waking and thus minimise the risk falling back asleep (start-time, 6:32 am \pm 30 min). No circadian variation in prolonged (e.g., 60-minute) endurance performance has been established for time of day differences such as those that occurred in the present study (e.g., 6:30am vs 7:50am) (16). All participants had either morning or intermediate chronotypes, and all routinely trained in the morning. Thus, all testing was undertaken at a time when participants would normally be awake (15), and when they would often be training. Consumption of caffeine and alcohol was prohibited on days -1 to D4. Athletes were experienced racers, so dietary requirements were self-determined. However, to prevent discrepancies in energy availability, athletes recorded (e.g., 7 am; 1 cup oats with milk) and replicated their dietary intake for each condition. Exercise was prohibited on days -1 to D4 (other

than that required for the experiment). However, to accommodate preferred preparation strategies between conditions, participants were permitted to exercise lightly on days -2, +1, and +2, and were required to replicate this exercise between conditions. Participants recorded all exercise so load could be quantified (17). No differences between conditions were noted on days prior to, or post laboratory testing [see Table, Supplemental Digital Content 1, Self-reported training load (arbitrary units) calculated as the product of exercise time (min) and session perceived exertion (0-10 scale), http://links.lww.com/MSS/B660]. A minimum seven-day washout period was required between D4 of a condition and D1 of the next condition.

Familiarisation sessions

Incremental test

On a cycle ergometer (Excalibur Sport, Lode, Groningen, Netherlands) controlled using compatible software (Lode Ergometry Manager 9, Lode, Groningen, Netherlands) participants cycled for three minutes at 75, 125, and 175 W respectively, before workload increased by 25 W every minute until volitional exhaustion. An Innocor metabolic system (DK-5260, Innovision, Odense, Denmark) determined maximal oxygen uptake ($\dot{V}O_{2max}$), and anaerobic threshold (AT).

Time-trial protocol

Target work for the TT was the estimated work expended when cycling at AT for one hour:

work
$$(kJ) = \frac{(W_{AT} \times 3600)}{1000}$$

Power at AT (W_{AT}) was determined from a regression of the relationship between oxygen uptake ($\dot{V}O_2$) and power (W) for the first three workloads of the incremental test. Pedalling resistance was calculated according to the formula:

$$W_{AT}$$
 = linear factor × preferred pedal rate²

where the linear factor ensured W_{AT} ocurred at the participant's preferred pedal rate per minute (rev·min⁻¹). A strong correlation has been demonstrated between W_{AT} and one-hour TT performance (r = 0.8, P<0.05) (18). Participants completed one practice TT to refine their pacing strategy. During the TT, work completed (kJ) was displayed on a computer screen. No other feedback/encouragement was provided.

Experimental conditions

Sleep

Participants wore activity monitors (Actical MiniMitter / Philips Respironics, Bend, OR) on their non-dominant wrist from day -2 to +2 to monitor sleep (19, 20). Activity counts were recorded in one-minute epochs and downloaded using a device specific interface unit (ActiReader, Philips Respironics, Bend, OR). Raw data was processed with a validated manufacturer proprietary v3.10) algorithm (Actical set to а medium sleep-wake threshold (<40 counts.min⁻¹ scored sleep) (19, 20). This threshold has shown 87% agreement with polysomnography when identifying sleep and wake states in elite cyclists (20). In order to verify, or identify misclassified sleep / wake states, participants completed a sleep diary that required them to record the time of day (i.e., to the nearest minute) they 'began attempting to sleep', and the time of day they 'woke up for the last time' for all sleep episodes (21). No daytime naps were permitted from day -1 until completion of testing on D4. For all sleep episodes, the total amount of sleep obtained (i.e., total sleep time – TST), and the percentage of time in bed spent asleep (i.e., sleep efficiency) were determined. For analysis, TST was aggregated from the end of one night's main sleep to the end of the next night's main sleep. Mean sleep efficiency was calculated for all sleep episodes during the same period. Subjective sleep quality (SQ) was recorded in the sleep diary upon waking each morning on a 5-point Likert scale (i.e., 1=Very Good, 2=Good, 3=Average, 4=Poor, 5=Very Poor).

Time-trial

Overall finishing time (minutes) was recorded. Target work was divided into four equal splits and perceived exertion (6-20 scale) recorded during the final minute of splits 1-3, and upon completion of split four (22).

Pre time-trial testing

Prior to the TT, upon arriving at the laboratory, participants completed psychometric testing. The Profile of Mood States (POMS) assessed the feelings of participants "right now" across 65 mood descriptors, providing scores for total mood disturbance, tension, depression, anger, vigour, fatigue, and confusion (23). Participants completed a touchscreen version of the psychomotor vigilance task (PVT) on a tablet device using the application sleep-2-Peak (v2.2.1, Proactive Life LLC, New York, NY). This version of the PVT has been validated against traditional PVT methods (24). The PVT measured reaction times to visual stimuli occurring at varying intervals over 10 minutes. Mean response time and the number of lapses >500 milliseconds were recorded.

Statistical analysis

Mean and SD were calculated for all variables. Generalised Estimating Equations with exchangeable correlation structures and robust standard errors analysed mean changes in outcome variables. Initial models tested for period and carryover effects, however no such effects were found (P>0.05). Models analysed two-, or three-way interactions for the factors 'condition', 'day', and 'split' (RPE only). Where interactions were significant (P<0.05), pairwise models were run for each 'day'. A p-value <0.025 was used to account for multiple comparisons. Additional models analysed main effects of 'day' for each condition. A p-value <0.05 was used. For sleep variables, nights -2 and -1 served as baseline values in separate models. For all other variables, D1 served as a baseline value. Analyses were performed in IBM SPSS statistics for Windows (v24.0,Armonk,NY).

RESULTS

Sleep

Total sleep time (Figure 2a) on nights D1, D2, and D3 was longer (P<0.001) in the SE condition (8.6 \pm 1.0; 8.3 \pm 0.6; 8.2 \pm 0.6 h, respectively), and shorter (P<0.001) in the SR condition (4.7 \pm 0.8; 4.8 \pm 0.8; 4.9 \pm 0.4 h), compared with NS (7.1 \pm 0.8; 6.5 \pm 1.0; 6.9 \pm 0.7 h). On night -2 (i.e., two nights prior to commencement of laboratory testing) TST was longer (P<0.01) in the SR condition (7.4 \pm 1.0 h) compared with SE (6.9 \pm 1.0 h). On night D4 (i.e., following the final laboratory testing session) TST was longer in the SR condition (7.5 \pm 0.8 h) compared with SE (6.6 \pm 0.9 h, P<0.001) and NS (7.1 \pm 0.7 h, P<0.02), while TST was also longer (P<0.02) in the NS condition compared with SE (6.6 \pm 1.3 h).

Within the SR condition, TST was shorter (P<0.01) on nights D1, D2, and D3 compared with nights -2 and -1, longer (P<0.02) on night +1 compared with night -1, and shorter (P<0.01) on

night -1 compared with night -2. Within the NS condition, TST was shorter (P<0.05) on nights -1 and D2 compared with night -2, and longer (P<0.05) on nights D1 and D4 compared with night -1. Within the SE condition, TST was longer (P<0.01) on nights D1, D2, and D3 compared with nights -2 and -1.

On night D2, sleep efficiency (Figure 2b) was lower (P<0.01) in the SE condition ($88\pm5\%$) compared with SR ($91\pm3\%$) and NS ($91\pm4\%$). On night D3, sleep efficiency was lower (P<0.025) in the SE condition ($86\pm5\%$) compared with SR ($90\pm4\%$) and NS ($90\pm5\%$). Within the SR condition, sleep efficiency was lower (P<0.01) on night D2 compared with baseline night -2.

On night D3, SQ (Figure 2c) was better (P<0.01) in the NS condition (2.7 ± 1.0) compared with SE (3.3 ± 0.7) . On night D4, SQ tended to be better (P=0.039) in the SR condition (2.8 ± 1.3) compared with SE (3.6 ± 0.9) . Within the SR condition, SQ was better (P<0.05) on night D3 compared with baseline night -2. Within the NS condition, SQ was worse (P<0.05) on night D4 compared with baseline night -1 Within the SE condition, SQ was worse (P<0.05) on night D4 compared with baseline night -1 Within the SE condition, SQ was worse (P<0.05) on night D4 compared with baseline night -2 and -1. [See Table, Supplemental Digital Content 2, Bedtime, get-up time, time in bed (TIB), total sleep time (TST), sleep efficiency (SE), subjective sleep quality (SQ), time-trial (TT) finishing time, and TT mean power output for each experimental condition, http://links.lww.com/MSS/B661.]

Time-Trial Performance

Shown in Figure 3, time was slower (P<0.02) on D3 of SR (60.4 ± 3.7 min) compared with NS (58.8 ± 2.5 min). Time was slower (P<0.02) on D4 of SR (62.0 ± 5.2 min) and NS (58.7 ± 3.4 min) compared with SE (56.8 ± 3.1 min). Within the SR condition, time was slower (P<0.05) on D2 and D4 compared with D1, and tended to be slower (P=0.053) on D3 compared with D1. [See Table,

Supplemental Digital Content 2, Bedtime, get-up time, time in bed (TIB), total sleep time (TST), sleep efficiency (SE), subjective sleep quality (SQ), time-trial (TT) finishing time, and TT mean power output for each experimental condition, http://links.lww.com/MSS/B661.]

Time-Trial Perceived Exertion

There was no difference in perceived exertion for any split between conditions, or any split between days within conditions (Table 1)

Psychomotor Vigilance Task

Mean response time (Table 2) was faster (P<0.025) on D3 and D4 of SE compared with SR and NS, and faster (P<0.025) on D4 of NS compared with SR. Within the SR condition, mean response time was slower (P<0.05) on D2, D3, and D4 compared with D1. Within the NS condition, mean response time was slower (P<0.05) on D2 and D4 compared with D1. Within the SE condition, mean response time was faster (P<0.05) on D4 compared with D1. Lapses were fewer (P<0.025) on D3 and D4 of SE compared with SR and NS. Lapses were fewer on D4 of NS compared with SR. Within the SR condition, lapses were greater (P<0.05) on D3 and D4 compared with D1.

Profile of Mood States

Total mood disturbance (Table 2) was higher (P<0.025) on D3 and D4 of SR compared with NS and SE. Within the SR condition, total mood disturbance was higher (P<0.05) on D2, D3, and D4 compared with D1. Confusion was higher (P<0.025) on D3 and D4 of SR compared with NS and SE. Within the SR condition, confusion was higher (P<0.05) on D3 and D4 compared with D1. Fatigue was higher (P<0.025) on D2, D3, and D4 of SR compared with SE, and higher (P<0.025) on D3 and D4 of SR compared with NS. Within the SR condition, fatigue was higher (P<0.05) on D2, D3, and D4 compared with D1. Within the NS condition, fatigue was higher (P<0.05) on D4 compared with D1. Within the SE condition, fatigue was higher (P<0.05) on D3 and D4 compared with D1. Vigour was lower (P<0.025) on D2, D3, and D4 of SR compared with SE, and lower on D3 of SR compared with NS. Vigour was higher (P<0.025) on D4 of SE compared with NS. Within the SR condition, vigour was lower (P<0.05) on D2, D3, and D4 of SR compared with D4 of SR compared with NS. Within the SR condition, vigour was lower (P<0.05) on D2, D3, and D4 of SE compared with D1. Within the NS condition, vigour was lower (P<0.05) on D3 and D4 compared with D1. Within the NS condition, vigour was lower (P<0.05) on D3 and D4 compared with D1.

DISCUSSION

Three nights of sleep extension better maintained endurance performance compared with both normal and restricted sleep. Compared with normal sleep, an extra ~90 minutes of sleep per night, for three consecutive nights, improved performance by 3%, or ~2 minutes across a ~60-minute TT. In contrast, reducing sleep for two consecutive nights by 144 and 102 minutes respectively, slowed TT performance by 3%, or ~1.5 minutes. Within the sleep restriction condition, performance was slower on day two and four compared with day one. However, performance was consistent over time in the normal and extended sleep conditions.

Sleep extension and endurance performance

Few studies have examined the effects of sleep extension on athletic performance. While extending sleep has been reported to improve sport-specific skill execution and sprint times (3, 4), this is the first study to examine the performance of endurance athletes. Moreover, previous studies examining sleep extension in athletes have used self-reported sleep times (3), or have not included a control arm (4). In contrast, the present study objectively monitored sleep and adopted a three-armed crossover design. In the present study, athletes habitually slept ~6.5-7.0 hours per night, similar to sleep durations reported in elite athletes (11). While a minimum seven hours of

sleep per night is recommended for good health (25), our findings suggest this may not be sufficient to optimise endurance performance. In fact, on sleep extension nights, athletes slept, on average, 8.4 hours per night (Figure 2a), similar to previous studies reporting improved athletic performance when sleep time was extended to 8.4 (4), and 8.9 (3) hours per night. Therefore, we recommend athletes sleep >8 hours per night to optimise performance. Sleep efficiency was consistently above 85% (Figure 2b), the minimum efficiency recommended for good health (26). However, sleep extension led to lower sleep efficiency compared with normal and restricted sleep, and poorer subjective sleep quality over time, perhaps indicative of reduced homeostatic sleep pressure (i.e., sleep 'need') (27). Therefore, sleep extension led to better maintenance of performance despite reductions in sleep efficiency. While future research should examine the precise impact of sleep quality on endurance performance, we recommend practitioners, with the help of valid sleep monitoring/assessment tools (20, 28), work with athletes to optimise both sleep quantity *and* quality.

Sleep restriction and endurance performance

The extent of accumulated sleep pressure may moderate the effect of sleep restriction on endurance performance. Compared with normal sleep, we found performance was unaffected by one night-, but impaired following two nights, of sleep restriction (i.e., ~5 hours TST per night). Previously, a severe sleep restriction protocol whereby cyclists slept 2.4 hours for one night, led to slower 3 km TT performance compared with 7.1 hours of sleep (10). In endurance athletes, the maximal workload achieved during a graded exercise test was unaffected when the prior night's sleep opportunity was reduced by three hours (8), but was lower when sleep opportunity was reduced by four hours (7). In taekwondo athletes, reducing sleep by 3-4 hours for one night did not affect distance covered during an intermittent test in the morning (5), but reduced distance

covered in the evening (6). Collectively, these findings suggest performance is likely impaired as sleep pressure/debt accumulates. Apparently contrary to this hypothesis, one study found time to exhaustion during a graded exercise test was unaffected following three consecutive nights of 2.5 hours sleep (9). Moreover, in the present study, we found performance was not statistically slower (P=0.09) on day four of sleep restriction, compared with normal sleep. This may reflect, on the part of at least some of the athletes tested, a subconscious increase in motivation for the final TT of the sequence as the fear of premature fatigue diminishes, akin to the 'end-spurt' effect demonstrated within endurance tasks (29). Nonetheless, within the sleep restriction condition, performance *was* slower on day two and four compared with day one. Therefore, collectively, the present findings suggest athletes should avoid short or restricted sleep, particularly on consecutive nights, for optimal endurance performance.

Effects of cumulative sleep time on perceived exertion

Cumulative sleep time did not affect perceived exertion despite differences in TT finishing times between conditions (Table 1). According to the linear nature of the TT protocol, finishing times correspond to mean power output (see Table, Supplemental Digital Content 2, which shows time and power output for each TT, http://links.lww.com/MSS/B661), thus compared with normal sleep, athletes' perceived exertion for a given power output was higher following sleep restriction (e.g., D3), and lower following sleep extension (e.g., D4). Perceived exertion reflects the effort required to overcome fatigue, and according to the psychobiological model of exercise tolerance, athletes disengage from an endurance task when perceived effort is greater than the maximum effort they are willing to exert, or believe they are capable of exerting (30). Our findings suggest total sleep obtained over 2-3 nights appears to alter the intensity (i.e., power output) at which these 'effort thresholds' occur. Increased perceived exertion during exercise has

been associated with mental fatigue (31). While we did not measure mental fatigue *per se*, we speculate that prior cumulative sleep time affects the level of mental fatigue experienced, or tolerated, during an endurance task. In fact, sleep extension has been shown to increase pain tolerance (i.e., ability to withstand pain) in healthy adults (32), which may explain higher power outputs for a given RPE after three nights of sleep extension. Evidence that sleep restriction *impaired* mood and psychomotor vigilance, while sleep extension *improved* vigour and psychomotor vigilance (Table 2), further supports speculation that mental/psychological determinants of endurance performance (e.g., attentional focus on pacing, response inhibition etc.) were likely affected by sleep extension and restriction (33).

Limitations

Participants were well-trained male endurance athletes, therefore, inferences for elite and/or female athletes may require caution. Caffeine withdrawal symptoms peak 20-51 hours post-abstinence (34), therefore, symptoms may have impaired performances on D1. However, given the crossover nature of the experiment this is unlikely to affect findings. Participants slept ~30 minutes more on night -2 of SR compared with SE, potentially confounding results. However, total sleep time for the 48 hours prior to D1 was no different (~ 14 hours, see Table, Supplementary Digital Content 2) between conditions, therefore, differences on night -2 are unlikely to affect findings. On D2, D3, and D4, mean start times of testing sessions differed slightly between conditions (see methods 'overview' section), potentially confounding results due to circadian variation in endurance capacity. However, performance differences between conditions did not manifest until after consecutive days of either sleep restriction (e.g., D3) or extension (e.g., D4). Thus, circadian effects cannon explain findings as any effects on performance should have occurred as soon as start times differed (e.g., D2). In addition, findings

from studies examining time of day effects on *prolonged endurance performances* (e.g., 60 minutes) have been equivocal (16), and any effects of small time of day changes, such as those occurring in the current study (e.g.,~ 40 minute difference between start times of the NS condition and the SR / SE conditions), have not been established.

Conclusions

Sleep extension for three consecutive nights better maintained prolonged self-paced endurance performance compared with both normal and restricted sleep. Sleep restriction impaired endurance performance. Sleep time accumulated over 2-3 nights appears to influence performance by altering perceived exertion during exercise. Athletes should aim to sleep >8 hours per night to optimise endurance performance.

DECLARATION

The results of this study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. Results do not constitute endorsement by the American College of Sports Medicine.

CONFLICT OF INTEREST

Spencer Roberts, Wei-Peng Teo, Brad Aisbett, and Stuart Warmington declare they have no conflict of interest.

ACKNOWLEDGEMENTS

The authors acknowledge the School of Exercise and Nutrition Sciences at Deakin University for funding the study. In addition, the authors thank the Deakin University Biostatistics Unit, particularly Dr Mohammadreza Mohebbi and Dr Steve Bowe, for their assistance with statistical analysis.

REFERENCES

 Lucía A, Hoyos J, Chicharro JL. Physiology of professional road cycling. Sports Med. 2001;31(5):325-37.

2. Fullagar HHK, Skorski S, Duffield R, Hammes D, Coutts AJ, Meyer T. Sleep and athletic performance: the effects of sleep loss on exercise performance, and physiological and cognitive responses to exercise. Sports Med. 2015;45(2):161-86.

3. Schwartz J, Simon RD. Sleep extension improves serving accuracy: A study with college varsity tennis players. Physiol Behav. 2015;151:541-4.

4. Mah CD, Mah KE, Kezirian EJ, Dement WC. The effects of sleep extension on the athletic performance of collegiate basketball players. Sleep. 2011;34(7):943.

5. Mejri MA, Hammouda O, Zouaoui K, Chaouachi A, Chamari K, Rayana MCB, et al. Effect of two types of partial sleep deprivation on Taekwondo players' performance during intermittent exercise. Biol Rhythm Res. 2014;45(1):17-26.

6. Mejri MA, Yousfi N, Mhenni T, Tayech A, Hammouda O, Driss T, et al. Does one night of partial sleep deprivation affect the evening performance during intermittent exercise in Taekwondo players? J Exerc Rehabil. 2016;12(1):47-53.

7. Mougin F, Bourdin H, Simon-Rigaud M, Nhu UN, Kantelip J, Davenne D. Hormonal responses to exercise after partial sleep deprivation and after a hypnotic drug-induced sleep. J Sports Sci. 2001;19(2):89-97.

 Mougin F, Simon-Rigaud M, Davenne D, Renaud A, Garnier A, Kantelip J, et al. Effects of sleep disturbances on subsequent physical performance. Eur J Appl Physiol. 1991;63(2):77-82.

9. Reilly T, Deykin T. Effects of partial sleep loss on subjective states, psychomotor and physical performance tests. J Hum Mov Studies. 1983;9:157-70.

10. Chase JD, Roberson PA, Saunders MJ, Hargens TA, Womack CJ, Luden ND. One Night of Sleep Restriction Following Heavy Exercise Impairs 3-km Cycling Time Trial Performance in the Morning. Appl Physiol Nutr Metab. 2017; 42(9):909-915.

11. Roberts SSH, Teo W-P, Warmington SA. Effects of training and competition on the sleep of elite athletes: a systematic review and meta-analysis. Br J Sports Med. 2019;53(8):513-522.

 Jeukendrup AE, Craig NP, Hawley JA. The bioenergetics of world class cycling. J Sci Med Sport. 2000;3(4):414-33.

 Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. Psychiatry Res. 1989;28(2):193-213.

14. Spielberger CD, Gorsuch R. State-Trait Anxiety Inventory for Adults: Manual and Sample: Manual, Instrument and Scoring Guide: Consulting Psychologists Press; 1983.

15. Horne JA, Östberg O. A self-assessment questionnaire to determine morningnesseveningness in human circadian rhythms. Int J Chronobiol. 1976;4:97-110.

16. Thun E, Bjorvatn B, Flo E, Harris A, Pallesen S. Sleep, circadian rhythms, and athletic performance. Sleep Med Rev. 2015;23:1-9.

17. Foster C. Monitoring training in athletes with reference to overtraining syndrome. Med Sci Sports Exerc. 1998;30(7):1164-8.

18. Lorenzo S, Minson CT, Babb TG, Halliwill JR. Lactate threshold predicting time-trial performance: impact of heat and acclimation. J Appl Physiol. 2011;111(1):221-7.

19. Kosmadopoulos A, Sargent C, Zhou X, Darwent D, Roach GD. Two for one? Validating an energy expenditure monitor against polysomnography for sleep/wake measurement. In: Zhou X, Sargent C (Eds). Sleep of Different Populations. 2012:16-20.

20. Sargent C, Lastella M, Halson SL, Roach GD. The validity of activity monitors for measuring sleep in elite athletes. J Sci Med Sport. 2016;19(10):848-53.

 Ancoli-Israel S, Martin JL, Blackwell T, Buenaver L, Liu L, Meltzer LJ, et al. The SBSM Guide to Actigraphy Monitoring: Clinical and Research Applications. Behav Sleep Med. 2015;13(sup1):S4-S38.

22. Borg GA. Psychophysical bases of perceived exertion. Med Sci Sports Exerc.1982;14(5):377-81.

23. McNair DM, Lorr M, Droppleman LF. Profile of Mood States Manual. Educational and Industrial Testing Service, San Diego. 1971.

24. Brunet J-F, Dagenais D, Therrien M, Gartenberg D, Forest G. Validation of sleep-2-Peak: A smartphone application that can detect fatigue-related changes in reaction times during sleep deprivation. Behav Res Meth. 2017;49(4):1460-9.

25. Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. Sleep Health. 2015;1(1):40-3.

26. Ohayon M, Wickwire EM, Hirshkowitz M, Albert SM, Avidan A, Daly FJ, et al. National Sleep Foundation's sleep quality recommendations: first report. Sleep Health. 2017;3(1):6-19.

27. Levine B, Lumley M, Roehrs T, Zorick F, Roth T. The effects of acute sleep restriction and extension on sleep efficiency. Int J Neurosci. 1988;43(3-4):139-43.

28. Samuels C, James L, Lawson D, Meeuwisse W. The Athlete Sleep Screening
Questionnaire: a new tool for assessing and managing sleep in elite athletes. Br J Sports Med.
2016;50(7):418-22.

29. Tucker R, Noakes TD. The anticipatory regulation of performance: the physiological basis for pacing strategies and the development of a perception-based model for exercise performance. Br J Sports Med. 2009;43(6):392-400.

30. Marcora SM. Do we really need a central governor to explain brain regulation of exercise performance? Eur J Appl Physiol. 2008;104(5):929.

31. Van Cutsem J, Marcora S, De Pauw K, Bailey S, Meeusen R, Roelands B. The effects of mental fatigue on physical performance: a systematic review. Sports Med. 2017;47(8):1569-88.

32. Simonelli G, Mantua J, Gad M, St Pierre M, Moore L, Yarnell AM, et al. Sleep extension reduces pain sensitivity. Sleep Med. 2019;54:172-6.

33. Brick NE, MacIntyre TE, Campbell MJ. Thinking and action: a cognitive perspective on self-regulation during endurance performance. Front Physiol. 2016;7:159.

Juliano LM, Griffiths RR. A critical review of caffeine withdrawal: empirical validation of symptoms and signs, incidence, severity, and associated features. Psychopharmacol. 2004;176(1):1-29.

CAPTIONS

Figure 1 Overview of data collection across the eight days/seven nights of each condition. Training (Tr.) load and diet were self-reported every day. Total sleep time, sleep efficiency, and subjective sleep quality were monitored throughout (-2 to +1). Bedtimes were prescribed on nights D1 to D3 according to the condition being undertaken. Laboratory testing was undertaken on days D1 to D4

Figure 2 Total sleep time (A), sleep efficiency (B) and subjective Sleep Quality (C) for sleep restriction (red line), normal sleep (black line) and sleep extension (green line) conditions. Shaded area represents the nights where bedtime interventions were implemented. [#] Different (P<0.025) to both normal sleep and sleep restriction. * Different (P<0.025) to both normal sleep and sleep restriction. * Different (P<0.025) to both normal sleep and sleep restriction and sleep extension only. ^ Difference (P<0.025) between normal sleep and sleep extension only. ^{a,b} Differences (P<0.05) within sleep restriction condition compared with -1 (a) and -2 (b). ^{c,d} Differences (P<0.05) within normal sleep extension condition compared with -1 (c) and -2 (d). ^{e,f} Differences (P<0.05) within normal sleep condition compared with -1 (e) and -2 (f)

Figure 3 Finishing time (mean \pm SD) for each time-trial across the four days (D1-D4) of testing. Sleep restriction (red line), normal sleep (black line), and sleep extension (green line). * Different (P<0.025) to sleep restriction. + Different (P<0.025) to sleep extension. ^ Different (P<0.05) to D1 of the same condition

SUPPLEMENTARY DIGITAL CONTENT

Supplementary Digital Content 1.pdf—Self-reported training load (arbitrary units) calculated as the product of exercise time (min) and session perceived exertion (0-10 scale)

Supplementary Digital Content 2.pdf—Bedtime, get-up time, time in bed (TIB), total sleep time (TST), sleep efficiency (SE), subjective sleep quality (SQ), time-trial (TT) finishing time, and TT mean power output for each experimental condition.



24

Figure 2







| Table 1. | Ratings of | perceived | exertion | recorded | for each | split | during | the tim | e-trials. | Data | presented | as mean | $s \pm SD$ |
|----------|------------|-----------|----------|----------|----------|-------|--------|---------|-----------|------|-----------|---------|------------|
|----------|------------|-----------|----------|----------|----------|-------|--------|---------|-----------|------|-----------|---------|------------|

| Day | D1 | | | | D2 | | | | D3 | | | | | |
|--------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| Time- | 1 | 2 | 3 | 4 | 1 | 2 | 3 | 4 | 1 | 2 | 3 | 4 | 1 | 2 |
| Sleep | 14±1 | 16±1 | 17±1 | 19±1 | 14±2 | 16±2 | 17±2 | 19±1 | 14±2 | 16±1 | 17±2 | 19±1 | 14±1 | 16±1 |
| Normal | 14±1 | 16±1 | 17±1 | 19±1 | 15±1 | 16±1 | 17±1 | 19±1 | 15±1 | 16±1 | 17±2 | 19±1 | 15±1 | 16±2 |
| Sleep | 14±1 | 16±1 | 17±1 | 19±1 | 15±1 | 16±1 | 17±1 | 19±1 | 15±2 | 16±1 | 17±2 | 19±1 | 15±2 | 16±2 |

D1-D4, testing days one to four. No significant differences for any split between conditions (P>0.025), or any split between days within conditions (P>0.05).

| | | Sleep Re | striction | | | | | | |
|------------------------------|---------|---------------------|--|--|--------|--------------------|----------------|----------------|---------|
| | D1 | D2 | D3 | D4 | D1 | D2 | D3 | D4 | D1 |
| Profile of Mood States | | | | | | | | | |
| Total mood disturbance | 2±11 | $10 \pm 10^{^{-1}}$ | 25±14 ^{+^} | $28\pm12^{+^{+}}$ | 3±13 | 3±14 | $9\pm14^{*}$ | 13±18* | 2±10 |
| Anger | 2±2 | 2 ± 1 | 3±2 | 2±2 | 3±2 | 2±1 | 2±2 | 2±2 | 3±2 |
| Confusion | 4±2 | 6±3 | $8\pm2^{+^{+}}$ | $9\pm3^{+^{+}}$ | 5±3 | 4±3 | 5±4* | 5±4* | 4±2 |
| Depression | 2±2 | 2±2 | $4\pm\!4$ | 3±3 | 2±2 | 2±3 | 3±2 | 1±1 | 3±2 |
| Fatigue | 4 ± 4 | $7\pm 4^{+^{+}}$ | $11 \pm 4^{+^{+^{+^{+^{+^{+^{+^{+^{+^{+^{+^{+^{+^$ | $14\pm5^{+^{+^{+^{+^{+^{+^{+^{+^{+^{+^{+^{+^{+^$ | 4±3 | 6±4 | $7\pm5^*$ | 9±6*^ | 4±2 |
| Tension | 8±6 | 5±3 | 6±4 | 7 ± 4 | 6±4 | 4±3 | 6±3 | 6±3 | 6±4 |
| Vigour | 17±4 | 13±3 ^{+^} | $7\pm5^{+^{+}}$ | $7\pm3^{+^{+}}$ | 18±5 | 14±6 | $13\pm5^*$ | $10 \pm 8^+$ | 17±6 |
| Psychomotor Vigilance | | | | | | | | | |
| Mean response time (ms) | 347±26 | 365±30 [^] | $374 \pm 31^{+}$ | $392 \pm 40^{+}$ | 348±34 | $363\pm30^{\circ}$ | $360 \pm 28^+$ | $363 \pm 28^+$ | 349±32 |
| Lapses (>500ms) | 2±1 | 3±2 | $4\pm2^{+}$ | $5\pm5^{+}$ | 2±1 | 3±2 | $3 \pm 1^{+}$ | $3\pm 2^{+*}$ | 2 ± 1 |

Table 2. Outcomes of preliminary testing conducted prior to time-trials. Data presented as means \pm SD.

D1-D4, laboratory testing days one to four.* Different (P<0.025) compared with sleep restriction. ⁺ Different (P<0.025) compared with sleep extension. [^]Different (P<0.05) compared with D1 of condition.