

Solving for X: evidence for sex-specific autism biomarkers across multiple transcriptomic studies

Samuel C. Lee^{1,+,*}, Thomas P. Quinn^{1,2,3,+,*}, Jerry Lai⁴, Sek W. Kong⁵, Irva Hertz-Picciotto⁶, Stephen J. Glatt⁷, Tamsyn M. Crowley^{2,3,8}, Svetha Venkatesh¹, and Thin Nguyen¹

¹Centre for Pattern Recognition and Data Analytics (PRaDA), Deakin University, Geelong, 3220, Australia

²Centre for Molecular and Medical Research, Deakin University, Geelong, 3220, Australia

³Bioinformatics Core Research Group, Deakin University, Geelong, 3220, Australia

⁴Deakin eResearch, Deakin University, Geelong, 3220, Australia | Intersect Australia, Sydney, 2000, Australia

⁵Computational Health Informatics Program, Boston Children's Hospital, Boston, MA, USA | Department of Pediatrics, Harvard Medical School, Boston, MA, USA

⁶Department of Public Health Sciences and UC Davis MIND Institute, School of Medicine, Davis, California

⁷Psychiatric Genetic Epidemiology and Neurobiology Laboratory (PsychGENe Lab) | SUNY Upstate Medical University, Syracuse, NY, USA

⁸Poultry Hub Australia, University of New England, Armidale, New South Wales, 2351, Australia

+ contributed equally, * samleenz@me.com; contacttomquinn@gmail.com

Abstract

Autism spectrum disorder (ASD) is a markedly heterogeneous condition with a varied phenotypic presentation. Its high concordance among siblings, as well as its clear association with specific genetic disorders, both point to a strong genetic etiology. However, the molecular basis of ASD is still poorly understood, although recent studies point to the existence of sex-specific ASD pathophysiologies and biomarkers. Despite this, little is known about how exactly sex influences the gene expression signatures of ASD probands. In an effort to identify sex-dependent biomarkers (and characterise their function), we present an analysis of a single paired-end post-mortem brain RNA-Seq data set and a meta-analysis of six blood-based microarray data sets. Here, we identify several genes with sex-dependent dysregulation, and many more with sex-independent dysregulation. Moreover, through pathway analysis, we find that these sex-independent biomarkers have substantially different biological roles than the sex-dependent biomarkers, and that some of these pathways are ubiquitously dysregulated in both post-mortem brain and blood. We conclude by synthesizing the discovered biomarker profiles with the extant literature, by highlighting the advantage of studying sex-specific dysregulation directly, and by making a call for new transcriptomic data that comprise large female cohorts.

1 Introduction

Autism Spectrum Disorder (ASD) is a markedly heterogeneous condition with a varied phenotypic presentation and a spectrum of disability for those affected. As a neurodevelopmental disorder, the ASD syndrome is characterised by social abnormalities, language abnormalities, and stereotyped behavioural patterns [3]. The presence of a genetic link in ASD etiology is well-established [38, 39], first evidenced by ASD concordance among siblings and by a clear association between ASD and specific genetic disorders (e.g., Fragile X mental retardation) [3]. This link has prompted a number of transcriptomic studies (e.g., [20, 17, 19]) to identify gene expression signatures (i.e., as a kind of biomarker) that might help elucidate the etiology of ASD and aid in its diagnosis (an important objective since early diagnosis and therapy is shown to improve outcomes in ASD [13]). However, despite the number of transcriptomic studies performed, the pathophysiology and biomarker profile of ASD are still not known. Rather, these studies have tended to produce inconsistent results, suggesting wide heterogeneity among both the individual patients and the study populations. Indeed, ASD may not have one signature at all, but instead multiple diverging signatures [53].

Transcriptomic studies of ASD probands typically use cells collected from either post-mortem brains or blood in order to estimate the mRNA abundance for thousands of gene transcripts (by way of microarray technology or massively parallel high-throughput sequencing (RNA-Seq)). Since many expressed transcripts are a precursor to structural or functional proteins, these studies can provide an insight into the functional state of a cell, capturing the common pathway for hereditary predisposition and environmental exposure. Although post-mortem brain studies have an advantage in that they look directly at the tissue of interest, blood-based studies can identify clinically useful biomarkers while also serving as a reliable proxy for gene expression in the brain [55] (though a complete understanding of ASD pathophysiology and its biomarker profile will likely require careful consideration of both lines of evidence). To date, more than a dozen studies have measured the transcriptomic profiles of ASD probands (and controls), the results of which have been summarised by two separate meta-analyses [10, 42] and one “mega-analysis” [53].

Sex is often called a risk factor for ASD, and it is stated that the risk for a male to have ASD is four to five times higher than that for females [56, 11] (although the magnitude of this difference may be partly due to diagnostic biases [28]). A similar observation, that the increased male risk is even higher among high-functioning ASD probands [16], likewise suggests that sex-specific mechanisms could influence ASD pathophysiology and its biomarker profile. Further evidence for sex-specific mechanisms is found in recent transcriptomic and functional-imaging studies. For example, Tylee et al., using transformed lymphoblastoid cell lines, found evidence for sex-specific differential regulation of genes and pathways among ASD probands [52]. Similarly, Trabzuni et al. found sex-specific differences in alternative splicing in adult human brains, including for a well-known ASD risk gene NRXN3 [50]. Functional brain connectivity studies using fMRI imaging have also identified sexual heterogeneity among ASD probands, showing dysregulation in sexually dimorphic brain regions across two large studies [15, 27]. Taken together, it seems plausible that sex could interact with other genetic and environmental factors to create sex-specific ASD pathophysiologies and biomarker profiles.

As ASD is more common in males, it suggests that females may have some underlying protection whereby a higher risk load is required for them to become afflicted [44]. One hypothesis posits that ASD itself reflects a shift towards “extreme maleness” such that males are necessarily predisposed [4]. In support of this, females with ASD do harbour more (and larger) copy number variants than males with ASDs [32], and moreover exhibit differential penetrance given the same genetic etiology [36]. Unfortunately, however, the increased prevalence of ASD in males has led to the exclusion of females from many transcriptomic studies (e.g., [21, 45, 1]), making it difficult to understand the male skew in ASD prevalence. Indeed, individual studies are often underpowered to detect subtle sex-specific differences (if they contain female subjects at all). When female subjects are included, sex is typically modelled as a simple covariate rather than an interaction term (i.e. the ASD-sex interaction), meaning that only sex-independent (and not sex-dependent) biomarkers are discovered. When male ASD is contrasted with female ASD, it typically involves loosely comparing simple sex-specific differences (e.g., differential expression present in males but not females, and *vice versa*) in a statistically anticonservative manner. To our knowledge, no study has looked at whether gene expression signatures show a sex-autism interaction across multiple studies and human tissues.

Using a single paired-end post-mortem brain RNA-Seq data set and a meta-analysis of six blood-based microarray data sets, we present a analysis of transcriptomic data that focuses on comparing sex-dependent and sex-independent ASD biomarkers (and the functional profiles thereof) across multiple tissues. By modelling the interaction of sex and ASD directly, we identify biomarkers (as well as functional pathways) that show sex-differences in ASD probands that are different than those in control subjects. Then, for those biomarkers that show no interaction, we pool male and female probands for a secondary sex-independent analysis. Our results suggest that, despite low power, some genes have FDR-adjusted significant sex-dependent interactions, while even more have significant sex-independent main effects. Subsequent pathway analysis further shows that these sex-independent biomarkers have substantially different biological roles than the sex-dependent biomarkers, and that some of these pathways are ubiquitously dysregulated in both post-mortem brain and blood.

2 Methods

2.1 Data acquisition

2.1.1 RNA-Seq data

We searched for relevant publicly available RNA-Seq data using the Gene Expression Omnibus (GEO) [5] with the term ("expression profiling by high throughput sequencing"[DataSet Type] AND ("autism spectrum disorder"[MeSH Terms] OR "autistic disorder"[MeSH Terms])) AND "homo sapiens"[Organism] (query made January 2018). We restricted eligible data sets to those sequenced with paired-end and non-poly-A-selected libraries. After excluding any data sets that used cell lines or did not have female cases, only one experiment, GSE107241 [57], remained. These data comprise a RiboZero Gold paired-end RNA-Seq data set from 52 postmortem dorsolateral prefrontal cortex tissue samples.

Prior to alignment and quantification, raw RNA-Seq reads were trimmed using Trimmomatic (docker image quay.io/biocontainers/trimmomatic:0.36-4) [9] and quality control metrics were recorded (before and after trimming) using FastQC (docker image biocontainers/fastqc:0.11.5) [2]. We aligned trimmed reads and quantified expression using Salmon (docker image combinelab/salmon:0.9.0) [43] as run in pseudo-quantification mode with a k-mer index of length 31. For the reference, we concatenated a human coding reference (i.e., GRCh38.90.cds) with the corresponding non-coding reference (i.e., GRCh38.90.ncrna).

2.1.2 Microarray data

We collected multiple microarray data sets to perform a meta-analysis of sex-autism interactions and main effects of ASD (i.e., sex-independent effects, where males and females are pooled). We referenced two prior meta-analyses [10, 42], and one "mega-analysis" [53], to prepare a list of data sets to study. Of these data sets, we excluded any study that (a) measured transcript expression from brain tissue, (b) had no female cases, (c) used cell lines (i.e., GSE37772 and GSE43076), or (d) treated cells with PPA (i.e., GSE32136). Six data sets remained after exclusion, as described in Table 1.

Data acquired from the Gene Expression Omnibus (GEO) [5] (i.e., GSE6575 [18] and GSE18123 [26]) were acquired already normalised and were not modified further. The other data sets (i.e., the Glatt et al. Wave I and Wave II data [17], the CHARGE study data [20], and the Kong et al. 2013 data [25]) each underwent RMA normalization, quantile normalization, and base-2 logarithm transformation. All subjects with a labelled condition other than typically developed (TD) were assigned to the autism spectrum disorder (ASD) group, except for the two Glatt et al. data sets where "Type-1 errors" were assigned to the TD group. Note that, in crafting this dichotomy, some subjects assigned to the ASD group have delays that fall outside of the "spectrum" *per se*.

2.2 Differential expression analysis of RNA-Seq data

We used DESeq2 (Version 3.6) [37] to test for differential transcript expression within the Salmon-generated counts. We applied a conservative expression filter (i.e., at least 10 estimated counts per-gene in every sample) to the raw count matrix to ensure that the high variability of lowly expressed transcripts did not bias results due to the small group sizes. For each transcript that passed the expression filter, a model was fit using the formula $\sim \text{ASD} * \text{Sex} + \text{Age}$ (where Age is the age of death). Interaction and sex-independent main effects (i.e., of the ASD condition) were then extracted from the model by specifying the relevant contrasts to the DESeq2::results function. We corrected for multiple testing using the Benjamini-Hochberg procedure [7].

2.3 Meta-analysis of microarray data

Before proceeding with the meta-analysis, we established a set of probes (i.e., for each microarray platform) that represent genes also represented by probes in the other platforms. In other words, we established a final probe set based on the intersection of unique gene symbols present in all microarray platforms under study. Note that we resolved one-to-many mapping ambiguities by excluding any probe that mapped to multiple gene symbols.

For each microarray data set, and for each probe (i.e., of those representing genes found in all data sets), we performed differential expression analysis using limma (Version 3.34) [46], applying

the following steps: (1) fit a model with the formula $\sim \text{ASD} * \text{Sex} + \text{Age}$ where ASD and Sex are each two-level factors (except GSE6575, where the Age covariate is unknown), (2) define contrasts for the sex-autism interaction and for the sex-independent main effects (i.e., of the ASD condition), and (3) measure the differential expression for each contrast using the eBayes procedure.

Next, we transformed platform-specific probe p-values to HGNC symbol p-values using AnnotationDbi (available from Bioconductor [22]). We resolved many-to-one mapping ambiguities by FDR-adjusting the minimum p-value of all probes for a given gene symbol (i.e., calculating a within-gene FDR correction). We then used Fisher’s method to perform a meta-analysis of the p-values obtained from the differential expression analysis. For K studies, Fisher’s method scores each gene based on (negative two times) the sum of the logarithm of the p-values:

$$\chi_{2K}^2 = -2 \sum_i^K \log p_i \quad (1)$$

This score follows a χ^2 distribution with $2K$ degrees of freedom [41]. Thus, for each gene, we computed a p-value directly from this score. We corrected for multiple testing using the Benjamini-Hochberg procedure [7].

2.4 Adjustment of latent batch effects

To ensure that latent batch effects did not inflate the discovery of false positives, we performed all analyses above with adjustment for batch effects using sva (Version 3.26) [31, 30], applying the following steps: (1) estimate the number of surrogate variables while specifying the ASD * Sex interaction as the variable of interest and Age as an adjustment variable, (2) use the sva function (or, in the case of Salmon-generated counts, the svaseq function) to estimate the surrogate variables, and (3) include the surrogate variables in the differential expression model(s) described above. Generally speaking, using sva yielded more conservative results than not using sva. All tables and figures show results generated with sva except where otherwise noted.

2.5 Pathway analysis and knowledge integration

We performed pathway analysis using GSEA (Version 3.0) [48] in PreRanked mode with classic enrichment and 1,000 permutations. Enrichment scores were calculated for specific MSigDB (Version 6.1) [47, 34] gene sets, including the curated KEGG (c2.cp.kegg[23]), Gene Ontology Biological Process (c5.bp) [49], Reactome (c2.cp.reactome) [14], and MSigDB Hallmark (h.all) [35] sets.

Based on the nature of the analyses, input rank lists were prepared differently for the RNA-Seq and microarray results. For the RNA-Seq analysis, we ranked transcripts based on the p-value, p , and the magnitude of the fold-change, FC:

$$\text{Rank} = -\log_{10}(p) \times \text{sign}(\log_2(\text{FC})) \quad (2)$$

Then, these transcript-level ranks were converted into gene-level ranks based on the top transcript-level rank. For the microarray meta-analysis, we ranked genes using the χ^2 test statistic (as calculated from Fisher’s method). Note that since this latter metric is agnostic to the direction of expression changes, we focused here on pathways enriched with a positive score (effectively making this pathway enrichment test one-tailed).

3 Results

3.1 Evidence for sex-dependent autism biomarkers

By modelling the sex-autism interaction directly, we can detect gene expression signatures that have differential dysregulation in male ASD probands when compared with female ASD probands. In other words, we can find sexually dimorphic ASD biomarkers (e.g., a gene up-regulated in male ASD but not in female ASD, or *vice versa*). Despite small study sizes (and disproportionately fewer females), we find some evidence for a sex-autism interaction among biomarkers, especially throughout the microarray meta-analysis data.

From the analysis of the RNA-Seq data derived from post-mortem brain tissue, we find no transcripts with significant (FDR-adjusted p-value < 0.05) sex-dependent dysregulation, although one

of these transcripts showed a significant interaction prior to batch correction with *sva*. To illustrate what a sex-autism interaction might look like, Figure 1 shows the per-group expression profiles for the two transcripts with the largest interaction effect (i.e., those with the smallest corrected p-value). Table 2 characterises those transcripts with the most sex-dependent dysregulation.

From the meta-analysis of the blood-based microarray data, we find two genes with significant (FDR-adjusted) sex-dependent dysregulation: *TTF2* and *UTY*. Table 3 characterises those genes with the most sex-dependent dysregulation. Since for a meta-analysis by Fisher's method, a large departure from the null (i.e., a very small p-value) in only one of several studies could cause the meta-analysis to post a significant result (i.e., even after FDR-adjustment) [51], it is useful to inspect visually how each study contributed to the results of the meta-analysis. For this, Figure 2 shows how each study contributed to the meta-analysis findings by plotting the aggregate Fisher score for each gene (of those with large sex-dependent dysregulation) along with the study-wise nominal significance (unadjusted p-value < 0.05). Notably, several of the most significantly dysregulated genes are at least nominally significant in more than one study.

3.2 Evidence for sex-independent autism biomarkers

In situations where a sex-autism interaction is not detectable, we can proceed to measure main condition (i.e., sex-independent) effects by pooling male ASD probands with female ASD probands (and male controls with female controls), without having to model sex as a covariate. Genes with significant sex-independent main effects (i.e., of the ASD condition) have large unidirectional effect sizes in male ASD probands, female ASD probands, or both. Yet, because the interaction is tested first, we can interpret the main condition effects as sex-independent.

From the analysis of the RNA-Seq data derived from post-mortem brain tissue, we find seven transcripts with significant (FDR-adjusted p-value < 0.05) sex-independent differential expression. Of these, only one transcript showed significant up-regulation in ASD. Figure 3 shows the expression profile for the two transcripts with the most significant sex-independent main effects (i.e., of the ASD condition). Table 4 characterises those transcripts with significant sex-independent dysregulation. Interestingly, several of the transcripts called differentially expressed by the analysis are annotated as non-coding RNA species.

From the meta-analysis of blood-based microarray data, we find 21 genes with significant (FDR-adjusted) sex-independent dysregulation. Table 5 characterises those genes with the most sex-independent dysregulation. As in Figure 2, Figure 4 shows how each study contributed to the meta-analysis findings by plotting the aggregate Fisher score for each gene (i.e., of those with large sex-independent dysregulation) along with the study-wise nominal significance (unadjusted p-value < 0.05). Again, most genes selected as statistically significant by the meta-analysis are at least nominally significant in more than one study.

3.3 Pathway enrichment of ASD biomarkers

In an effort to summarise the biological relevance of the biomarker profiles generated above, we used the complete ranked lists of the differentially expressed transcripts (and genes) in four separate gene set enrichment analyses to identify common differentially regulated pathways. Four enrichment profiles were generated using the sex-dependent RNA-Seq (brain) biomarkers, sex-independent RNA-Seq (brain) biomarkers, sex-dependent microarray (blood) biomarkers, and sex-independent microarray (blood) biomarkers.

Figure 5 shows the KEGG pathways enriched by the biomarkers as ranked by the analysis of the RNA-Seq data. For the sex-dependent biomarkers, nine pathways showed significant (FDR-adjusted p-value < 0.15) enrichment. For the sex-independent biomarkers, five pathways showed significant enrichment. Interestingly, all significant enrichment occurred in the same direction.

Figure 6 shows the KEGG pathways enriched by the biomarkers as ranked by the analysis of the microarray data. For the sex-dependent biomarkers, one pathway (i.e., Alanine Aspartate and Glutamate Metabolism) showed significant (FDR-adjusted p-value < 0.30) enrichment. For the sex-independent biomarkers, thirty-six pathways showed significant enrichment. Note that because only positive (i.e., one-tailed) enrichments are considered for these data, an FDR-adjusted p-value < 0.30 is used here (see Methods for more details).

Figure 7 compares the overlap between these significant pathways. For the sex-dependent analyses, no pathways are enriched in both the RNA-Seq and microarray data. However, for the sex-independent analyses, two pathways are enriched in both data. Interestingly, this agreement

exists despite differences in the ranked lists, suggesting that ASD biomarker profiles may show some degree of higher-order conservation at the pathway-level that exists not only across multiple studies, but across multiple tissues (as well as multiple transcript quantification assays). Note that we also tested for enrichment among the Gene Ontology Biological Process, Reactome, and MSigDB Hallmarks gene sets, all of which show more examples of overlap between the separate sex-independent analyses (see the Supplementary Information for more details).

4 Discussion

In this report, we present an analysis of several ASD transcriptomic studies, including an analysis of RNA-Seq data derived from post-mortem brain and a meta-analysis of six blood-based microarray data sets. Specifically, we focus on identifying both sex-dependent and sex-independent biomarker profiles for ASD by modelling the sex-autism interaction directly and secondarily measuring main effects of the ASD condition (i.e., sex-independent effects where males and females are pooled). In addition to identifying transcript (and gene) biomarkers, we use gene set enrichment analysis to summarise the observed dysregulation at the pathway level, contrasting sex-dependent pathway enrichment with sex-independent pathway enrichment. In doing so, we find evidence that ASD biomarker profiles may show some degree of higher-order conservation at the pathway level that exists not only across multiple studies, but across multiple tissues (and across multiple transcript quantification assays).

Despite small sample sizes in all studies, we found evidence for the existence of some sex-dependent biomarkers in human tissue. The meta-analysis identified two genes, *TTF2* and *UTY*, with sexually dimorphic expression in the blood. One of these, *TTF2*, plays an important role in normal thyroid development [12]. Interestingly, a loss of thyroid hormone homeostasis has been linked to ASD [8, 24]. Since it is well-known that thyroid diseases have a sex-specific presentation [6], it seems plausible that thyroid abnormalities could contribute to a sexually dimorphic ASD signature. The other, *UTY*, is a Y-chromosome gene, making any interpretation of its differential dysregulation difficult. Two other genes, *KCNJ8* and *MAP1B*, had FDR-adjusted p-values very close to the pre-defined significance cutoff, warranting follow-up in another study. Although the RNA-Seq analysis did not yield any significant interactions, it is not surprising considering this data set contained only three female ASD probands. Nevertheless, the large (albeit non-significant) effect sizes warrant repeat studies with bigger cohorts and more female ASD probands.

By modelling the sex-autism interaction directly, we are able to follow-up the sex-dependent analysis with a secondary sex-independent analysis for any transcript (or gene) whose expression did not significantly interact with sex. In this scenario, we contrast the pooled male ASD probands and female ASD probands against the pooled male controls and female controls to calculate the main effects (which we can thus interpret as sex-independent biomarkers). Here, over twenty transcripts and genes exceeded the threshold for FDR-adjusted significance. Interestingly, for the RNA-Seq data, several of the significant biomarkers are not protein-coding genes (highlighting the value of using non-poly-A-selected libraries to quantify both coding and non-coding transcripts). For the microarray meta-analysis, several of the sex-independent biomarkers are associated with key neurodevelopmental processes, including some X-chromosome genes. For example, *MAGED2*, differentially expressed in ASD probands, is located on an X-linked intellectual disability hotspot (i.e., Xp11.2) [29, 40] (which, if causally relevant, could contribute to the male risk bias).

For both the RNA-Seq analysis and the microarray meta-analysis, we tested the ranked sex-dependent and sex-independent biomarker profiles separately for pathway-level enrichment. We found some pathway enrichment for the sex-dependent profiles, and even more for the sex-independent profiles. Importantly, very few of the enriched pathways were the same for both the interaction and main effects. This suggests that males and females exhibit unique pathway-level signatures that, if causally relevant, might further suggest the existence of both sex-specific and common ASD pathophysiologies. Although few KEGG pathways are enriched among the sex-dependent results, there are dozens of significantly enriched sex-dependent pathways across other tested gene sets (see Supplementary Information for more details). Among the sex-independent enriched pathways (for the meta-analysis results), there are a number of pathways for known neurodevelopmental and neurodegenerative diseases, including Huntingtons, Parkinsons, Alzheimers, and amyotrophic lateral sclerosis (ALS), suggesting that at least some of these ASD biomarkers may have functions important to general brain health. Considering that both unique and shared signatures (i.e., at the biomarker-level and pathway-level) exist among ASD probands, it seems plausible that molecular

diagnostics could benefit from modelling sex-specific processes directly.

Although we found pathway enrichment to differ considerably between the sex-dependent and sex-independent biomarker profiles, we found that several sex-independent pathways (i.e., based on KEGG and other genes sets) were enriched across both the RNA-Seq and microarray data. Interestingly, this overlap exists despite the fact that analyses were performed on different human tissues (and with different transcript quantification assays). In fact, more than fifty Gene Ontology pathways were enriched among both sets of ranked sex-independent biomarkers (even though no gene products showed significant differential expression in both data). This overlap supports the hypothesis that there may exist common diagnostic (and perhaps etiological) signatures across the widely heterogeneous population of ASD probands. If true, it seems plausible that molecular diagnostics could further benefit from modelling pathway-level dysregulation directly (i.e., in addition to modelling conventional transcriptomic biomarkers).

When we compare our pathway enrichments to the previous ASD “mega-analysis” pathway enrichments [54], we observe several complementary results. First, we found positive enrichment of the MAPK pathway in our sex-dependent RNA-Seq results, agreeing with the male-specific enrichment of Mek targets found in the Tylee et al. study [54]. Second, we found an enrichment of the ribosome-related pathway in both of our sex-independent analyses, agreeing with the ribosome-related pathway enrichment identified by the sex-independent “mega-analysis” [54]. Third, we found an enrichment of the Toll-like receptor (TLR) signalling pathway in our sex-independent meta-analysis results, agreeing with the TLR 3 and 4 signalling pathway enrichment identified by the sex-independent “mega-analysis” [54]. Importantly, these complementary results exist despite considerable differences in statistical methodology and data set inclusion.

Our analysis is not without limitations. First, although we used *sva* to adjust for latent batch effects, it is still possible that any number of remaining factors (or batch effects) could coincide with the diagnostic label (e.g., undocumented co-morbidities or medication use), thereby confounding the discovered biomarker profile. Second, as with any observational study, it is impossible to conclude whether the gene expression signatures (and their biological pathways) are causally related to ASD (or, likewise, the sex-autism interaction). Third, this analysis is likely under-powered to detect both sex-autism interactions and main effects, owing to the small sample sizes and disproportionately smaller female cohorts. Yet, based on the extant literature (which clearly highlights sex as an ASD risk factor) and the results published here, we believe that modelling the sex-autism interaction should become a mainstay of ASD transcriptomic research. Advantageously, as shown here, interaction modelling is compatible with the most commonly used softwares for batch-effect correction [31], RNA-Seq analysis [37], and microarray analysis [46]. Yet, this analytical technique cannot offer any benefit if transcriptomic studies continue to systematically exclude female subjects ([21, 45, 1]). Although there seems to exist a strong skew in the prevalence of male ASD, this very fact underlies the importance of studying female ASD at equal proportions: a complete understanding of the molecular basis of ASD will require the intentional study of both sex-dependent and sex-independent mechanisms, as well as their differences and commonalities.

Acknowledgements

This research was partially supported by the Australian Government through the Australian Research Council’s Linkage Projects funding scheme (project LP140100240). SWK is supported by a grant from the National Institute of Health (R01MH107205).

References

- [1] Mark D. Alter, Rutwik Kharkar, Keri E. Ramsey, David W. Craig, Raun D. Melmed, Theresa A. Grebe, R. Curtis Bay, Sharman Ober-Reynolds, Janet Kirwan, Josh J. Jones, J. Blake Turner, Rene Hen, and Dietrich A. Stephan. Autism and increased paternal age related changes in global levels of gene expression regulation. *PloS One*, 6(2):e16715, February 2011.
- [2] S Andrews. FastQC a Quality Control Tool for High Throughput Sequence Data [Online], 2010.

- 359 [3] A. Bailey, W. Phillips, and M. Rutter. Autism: towards an integration of clinical, genetic,
360 neuropsychological, and neurobiological perspectives. *Journal of Child Psychology and Psy-*
361 *chiatry, and Allied Disciplines*, 37(1):89–126, January 1996.
- 362 [4] Simon Baron-Cohen. The extreme male brain theory of autism. *Trends in cognitive sciences*,
363 6(6):248–254, 6 2002.
- 364 [5] Tanya Barrett and Ron Edgar. Gene Expression Omnibus (GEO): Microarray data storage,
365 submission, retrieval, and analysis. *Methods in enzymology*, 411:352–369, 2006.
- 366 [6] Michael Bauer, Tasha Glenn, Maximilian Pilhatsch, Andrea Pfennig, and Peter C Whybrow.
367 Gender differences in thyroid system function: relevance to bipolar disorder and its treatment.
368 *Bipolar Disorders*, 16(1):58–71, 2 2014.
- 369 [7] Yoav Benjamini and Yosef Hochberg. Controlling the False Discovery Rate: A Practical and
370 Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*
371 *(Methodological)*, 57(1):289–300, 1995.
- 372 [8] Pere Berbel, Daniela Navarro, and Gustavo C Román. An evo-devo approach to thyroid
373 hormones in cerebral and cerebellar cortical development: etiological implications for autism.
374 *Frontiers in endocrinology*, 5:146, 2014.
- 375 [9] Anthony M. Bolger, Marc Lohse, and Bjoern Usadel. Trimmomatic: A flexible trimmer for
376 Illumina sequence data. *Bioinformatics*, 30(15), 2014.
- 377 [10] Carolyn Ch’ng, Willie Kwok, Sanja Rogic, and Paul Pavlidis. Meta-Analysis of Gene Expres-
378 sion in Autism Spectrum Disorder. *Autism Research*, 8(5):593–608, 10 2015.
- 379 [11] Deborah L. Christensen, Jon Baio, Kim Van Naarden Braun, Deborah Bilder, Jane Charles,
380 John N. Constantino, Julie Daniels, Maureen S. Durkin, Robert T. Fitzgerald, Margaret
381 Kurzius-Spencer, Li-Ching Lee, Sydney Pettygrove, Cordelia Robinson, Eldon Schulz, Chris
382 Wells, Martha S. Wingate, Walter Zahorodny, and Marshelyn Yeargin-Allsopp. Prevalence
383 and Characteristics of Autism Spectrum Disorder Among Children Aged 8 Years — Autism
384 and Developmental Disabilities Monitoring Network, 11 Sites, United States, 2012. *MMWR.*
385 *Surveillance Summaries*, 65(3):1–23, 4 2016.
- 386 [12] Mario De Felice and Roberto Di Lauro. Thyroid Development and Its Disorders: Genetics
387 and Molecular Mechanisms. *Endocrine Reviews*, 25(5):722–746, 10 2004.
- 388 [13] Jennifer Harrison Elder, Consuelo Maun Kreider, Susan N Brasher, and Margaret Ansell.
389 Clinical impact of early diagnosis of autism on the prognosis and parent-child relationships.
390 *Psychology research and behavior management*, 10:283–292, 2017.
- 391 [14] Antonio Fabregat, Steven Jupe, Lisa Matthews, Konstantinos Sidiropoulos, Marc Gillespie,
392 Phani Garapati, Robin Haw, Bijay Jassal, Florian Korninger, Bruce May, Marija Milacic, Co-
393 rina Duenas Roca, Karen Rothfels, Cristoffer Sevilla, Veronica Shamovsky, Solomon Shorser,
394 Thawfeek Varusai, Guilherme Viteri, Joel Weiser, Guanming Wu, Lincoln Stein, Henning
395 Hermjakob, and Peter D’Eustachio. The Reactome Pathway Knowledgebase. *Nucleic Acids*
396 *Research*, 46(D1):D649–D655, 1 2018.
- 397 [15] Dorothea L. Floris, Meng-Chuan Lai, Tanmay Nath, Michael P. Milham, and Adriana Di Mar-
398 tino. Network-specific sex differentiation of intrinsic brain function in males with autism.
399 *Molecular Autism*, 9(1):17, 12 2018.
- 400 [16] E Fombonne. The epidemiology of autism: a review. *Psychological medicine*, 29(4):769–86, 7
401 1999.
- 402 [17] Stephen J. Glatt, Ming T. Tsuang, Mary Winn, Sharon D. Chandler, Melanie Collins, Linda
403 Lopez, Melanie Weinfeld, Cindy Carter, Nicholas Schork, Karen Pierce, and Eric Courchesne.
404 Blood-based gene expression signatures of infants and toddlers with autism. *Journal of the*
405 *American Academy of Child and Adolescent Psychiatry*, 51(9):934–944.e2, September 2012.
- 406 [18] Jeffrey P. Gregg, Lisa Lit, Colin A. Baron, Irva Hertz-Picciotto, Wynn Walker, Ryan A.
407 Davis, Lisa A. Croen, Sally Ozonoff, Robin Hansen, Isaac N. Pessah, and Frank R. Sharp.
408 Gene expression changes in children with autism. *Genomics*, 91(1):22–29, January 2008.

- 409 [19] Simone Gupta, Shannon E. Ellis, Foram N. Ashar, Anna Moes, Joel S. Bader, Jianan Zhan,
410 Andrew B. West, and Dan E. Arking. Transcriptome analysis reveals dysregulation of innate
411 immune response genes and neuronal activity-dependent genes in autism. *Nature Communi-*
412 *cations*, 5:5748, December 2014.
- 413 [20] Irva Hertz-Picciotto, Lisa A. Croen, Robin Hansen, Carrie R. Jones, Judy van de Water, and
414 Isaac N. Pessah. The CHARGE study: an epidemiologic investigation of genetic and environ-
415 mental factors contributing to autism. *Environmental Health Perspectives*, 114(7):1119–1125,
416 July 2006.
- 417 [21] Valerie W. Hu, Tewarit Sarachana, Kyung Soon Kim, AnhThu Nguyen, Shreya Kulkarni,
418 Mara E. Steinberg, Truong Luu, Yinglei Lai, and Norman H. Lee. Gene expression profiling
419 differentiates autism case-controls and phenotypic variants of autism spectrum disorders: ev-
420 idence for circadian rhythm dysfunction in severe autism. *Autism Research: Official Journal*
421 *of the International Society for Autism Research*, 2(2):78–97, April 2009.
- 422 [22] Wolfgang Huber, Vincent J Carey, Robert Gentleman, Simon Anders, Marc Carlson, Benil-
423 ton S Carvalho, Hector Corrada Bravo, Sean Davis, Laurent Gatto, Thomas Girke, Raphael
424 Gottardo, Florian Hahne, Kasper D Hansen, Rafael A Irizarry, Michael Lawrence, Michael I
425 Love, James MacDonald, Valerie Obenchain, Andrzej K Oleś, Hervé Pagès, Alejandro Reyes,
426 Paul Shannon, Gordon K Smyth, Dan Tenenbaum, Levi Waldron, and Martin Morgan. Or-
427 chestrating high-throughput genomic analysis with Bioconductor. *Nature Methods*, 12(2):115–
428 121, 2 2015.
- 429 [23] Minoru Kanehisa, Miho Furumichi, Mao Tanabe, Yoko Sato, and Kanae Morishima. KEGG:
430 New perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Research*,
431 45(D1):D353–D361, 2017.
- 432 [24] A Khan, J W Harney, A M Zavacki, and E M Sajdel-Sulkowska. Disrupted brain thyroid
433 hormone homeostasis and altered thyroid hormone-dependent brain gene expression in autism
434 spectrum disorders. *Journal of physiology and pharmacology : an official journal of the Polish*
435 *Physiological Society*, 65(2):257–72, 4 2014.
- 436 [25] S. W. Kong, Y. Shimizu-Motohashi, M. G. Campbell, I. H. Lee, C. D. Collins, S. J. Brewster,
437 I. A. Holm, L. Rappaport, I. S. Kohane, and L. M. Kunkel. Peripheral blood gene expres-
438 sion signature differentiates children with autism from unaffected siblings. *Neurogenetics*,
439 14(2):143–152, May 2013.
- 440 [26] Sek Won Kong, Christin D. Collins, Yuko Shimizu-Motohashi, Ingrid A. Holm, Malcolm G.
441 Campbell, In-Hee Lee, Stephanie J. Brewster, Ellen Hanson, Heather K. Harris, Kathryn R.
442 Lowe, Adrianna Saada, Andrea Mora, Kimberly Madison, Rachel Hundley, Jessica Egan,
443 Jillian McCarthy, Ally Eran, Michal Galdzicki, Leonard Rappaport, Louis M. Kunkel, and
444 Isaac S. Kohane. Characteristics and predictive value of blood transcriptome signature in
445 males with autism spectrum disorders. *PloS One*, 7(12):e49475, 2012.
- 446 [27] M.-C. Lai, M. V. Lombardo, J. Suckling, A. N. V. Ruigrok, B. Chakrabarti, C. Ecker, S. C. L.
447 Deoni, M. C. Craig, D. G. M. Murphy, E. T. Bullmore, and S. Baron-Cohen. Biological sex
448 affects the neurobiology of autism. *Brain*, 136(9):2799–2815, 9 2013.
- 449 [28] Meng-Chuan Lai, Michael V Lombardo, Bonnie Auyeung, Bhismadev Chakrabarti, and Si-
450 mon Baron-Cohen. Sex/gender differences and autism: setting the scene for future research.
451 *Journal of the American Academy of Child and Adolescent Psychiatry*, 54(1):11–24, 1 2015.
- 452 [29] K. Langnaese, D.U. Kloos, M. Wehnert, B. Seidel, and P. Wieacker. Expression pattern
453 and further characterization of human MAGED2 and identification of rodent orthologues.
454 *Cytogenetic and Genome Research*, 94(3-4):233–240, 2001.
- 455 [30] Jeffrey T. Leek. svaseq: removing batch effects and other unwanted noise from sequencing
456 data. *Nucleic Acids Research*, 42(21):e161–e161, 12 2014.
- 457 [31] Jeffrey T. Leek, W. Evan Johnson, Hilary S. Parker, Andrew E. Jaffe, and John D. Storey.
458 The sva package for removing batch effects and other unwanted variation in high-throughput
459 experiments. *Bioinformatics*, 28(6):882–883, March 2012.

- [32] Dan Levy, Michael Ronemus, Boris Yamrom, Yoon-ha Lee, Anthony Leotta, Jude Kendall, Steven Marks, B. Lakshmi, Deepa Pai, Kenny Ye, Andreas Buja, Abba Krieger, Seungtai Yoon, Jennifer Troge, Linda Rodgers, Ivan Iossifov, and Michael Wigler. Rare De Novo and Transmitted Copy-Number Variation in Autistic Spectrum Disorders. *Neuron*, 70(5):886–897, June 2011.
- [33] Alexander Lex, Nils Gehlenborg, Hendrik Strobelt, Romain Vuillemot, and Hanspeter Pfister. UpSet: Visualization of Intersecting Sets. *IEEE Transactions on Visualization and Computer Graphics*, 20(12):1983–1992, 12 2014.
- [34] A. Liberzon, A. Subramanian, R. Pinchback, H. Thorvaldsdottir, P. Tamayo, and J. P. Mesirov. Molecular signatures database (MSigDB) 3.0. *Bioinformatics*, 27(12):1739–1740, 6 2011.
- [35] Arthur Liberzon, Chet Birger, Helga Thorvaldsdóttir, Mahmoud Ghandi, Jill P. Mesirov, and Pablo Tamayo. The Molecular Signatures Database Hallmark Gene Set Collection. *Cell Systems*, 1(6):417–425, 12 2015.
- [36] Anath C. Lionel, Kristiina Tammimies, Andrea K. Vaags, Jill A. Rosenfeld, Joo Wook Ahn, Daniele Merico, Abdul Noor, Cassandra K. Runke, Vamsee K. Pillalamarri, Melissa T. Carter, Matthew J. Gazzellone, Bhooma Thiruvahindrapuram, Christina Fagerberg, Lone W. Laulund, Giovanna Pellicchia, Sylvia Lamoureux, Charu Deshpande, Jill Clayton-Smith, Ann C. White, Susan Leather, John Trounce, H. Melanie Bedford, Eli Hatchwell, Peggy S. Eis, Ryan K. C. Yuen, Susan Walker, Mohammed Uddin, Michael T. Geraghty, Sarah M. Nikkel, Eva M. Tomiak, Bridget A. Fernandez, Noam Soreni, Jennifer Crosbie, Paul D. Arnold, Russell J. Schachar, Wendy Roberts, Andrew D. Paterson, Joyce So, Peter Szatmari, Christina Chrysler, Marc Woodbury-Smith, R. Brian Lowry, Lonnie Zwaigenbaum, Divya Mandyam, John Wei, Jeffrey R. Macdonald, Jennifer L. Howe, Thomas Nalpathamkalam, Zhuozhi Wang, Daniel Tolson, David S. Cobb, Timothy M. Wilks, Mark J. Sorensen, Patricia I. Bader, Yu An, Bai-Lin Wu, Sebastiano Antonino Musumeci, Corrado Romano, Diana Postorivo, Anna M. Nardone, Matteo Della Monica, Gioacchino Scarano, Leonardo Zoccante, Francesca Novara, Orsetta Zuffardi, Roberto Ciccone, Vincenzo Antona, Massimo Carella, Leopoldo Zelante, Pietro Cavalli, Carlo Poggiani, Ugo Cavallari, Bob Argiropoulos, Judy Chernos, Charlotte Brasch-Andersen, Marsha Speevak, Marco Fichera, Caroline Mackie Ogilvie, Yiping Shen, Jennelle C. Hodge, Michael E. Talkowski, Dimitri J. Stavropoulos, Christian R. Marshall, and Stephen W. Scherer. Disruption of the ASTN2/TRIM32 locus at 9q33.1 is a risk factor in males for autism spectrum disorders, ADHD and other neurodevelopmental phenotypes. *Human Molecular Genetics*, 23(10):2752–2768, May 2014.
- [37] M I Love, W Huber, and S Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12):550, 2014.
- [38] Judith H Miles. Autism spectrum disorders—a genetics review. *Genetics in medicine : official journal of the American College of Medical Genetics*, 13(4):278–294, 4 2011.
- [39] Shingo Miyauchi and Irina Voineagu. Autism susceptibility genes and the transcriptional landscape of the human brain. *International review of neurobiology*, 113:303–318, 2013.
- [40] Ching Moey, Susan J Hinze, Louise Brueton, Jenny Morton, Dominic J McMullan, Benjamin Kamien, Christopher P Barnett, Nicola Brunetti-Pierri, Jillian Nicholl, Jozef Gecz, and Cheryl Shoubbridge. Xp11.2 microduplications including IQSEC2, TSPYL2 and KDM5C genes in patients with neurodevelopmental disorders. *European Journal of Human Genetics*, 24(3):373–380, 3 2016.
- [41] Frederick Mosteller and R. A. Fisher. Questions and Answers. *The American Statistician*, 2(5):30–31, 1948.
- [42] L. F. Ning, Y. Q. Yu, E. T. GuoJi, C. G. Kou, Y. H. Wu, J. P. Shi, L. Z. Ai, and Q. Yu. Meta-analysis of differentially expressed genes in autism based on gene expression data. *Genetics and molecular research: GMR*, 14(1):2146–2155, March 2015.
- [43] R Patro, G Duggal, M I Love, R A Irizarry, and C Kingsford. Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods*, 14(4):417–419, 2017.

- 512 [44] Elise B Robinson, Paul Lichtenstein, Henrik Anckarsäter, Francesca Happé, and Angelica
513 Ronald. Examining and interpreting the female protective effect against autistic behavior. *Pro-*
514 *ceedings of the National Academy of Sciences of the United States of America*, 110(13):5258–62,
515 3 2013.
- 516 [45] Tewarit Sarachana, Rulun Zhou, Guang Chen, Hussein K. Manji, and Valerie W. Hu. In-
517 vestigation of post-transcriptional gene regulatory networks associated with autism spectrum
518 disorders by microRNA expression profiling of lymphoblastoid cell lines. *Genome Medicine*,
519 2(4):23, April 2010.
- 520 [46] Gordon K. Smyth. Linear models and empirical bayes methods for assessing differential
521 expression in microarray experiments. *Statistical Applications in Genetics and Molecular*
522 *Biology*, 3:Article3, 2004.
- 523 [47] A. Subramanian, P. Tamayo, V. K. Mootha, S. Mukherjee, B. L. Ebert, M. A. Gillette,
524 A. Paulovich, S. L. Pomeroy, T. R. Golub, E. S. Lander, and J. P. Mesirov. Gene set enrich-
525 ment analysis: A knowledge-based approach for interpreting genome-wide expression profiles.
526 *Proceedings of the National Academy of Sciences*, 102(43):15545–15550, 2005.
- 527 [48] Aravind Subramanian, Pablo Tamayo, Vamsi K. Mootha, Sayan Mukherjee, Benjamin L.
528 Ebert, Michael A. Gillette, Amanda Paulovich, Scott L. Pomeroy, Todd R. Golub, Eric S.
529 Lander, and Jill P. Mesirov. Gene set enrichment analysis: A knowledge-based approach for
530 interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences*,
531 102(43):15545–15550, October 2005.
- 532 [49] The Gene Ontology Consortium. Expansion of the Gene Ontology knowledgebase and re-
533 sources. *Nucleic Acids Research*, 45(D1):D331–D338, 1 2017.
- 534 [50] Daniah Trabzuni, Adaikalavan Ramasamy, Sabaena Imran, Robert Walker, Colin Smith,
535 Michael E. Weale, John Hardy, Mina Ryten, and North American Brain Expression Con-
536 sortium. Widespread sex differences in gene expression and splicing in the adult human brain.
537 *Nature Communications*, 4:2771, 11 2013.
- 538 [51] George C. Tseng, Debashis Ghosh, and Eleanor Feingold. Comprehensive literature review and
539 statistical considerations for microarray meta-analysis. *Nucleic Acids Research*, 40(9):3785–
540 3799, May 2012.
- 541 [52] Daniel S. Tylee, Alfred J. Espinoza, Jonathan L. Hess, Muhammad A. Tahir, Sarah Y. McCoy,
542 Joshua K. Rim, Totadri Dhimal, Ori S. Cohen, and Stephen J. Glatt. RNA sequencing
543 of transformed lymphoblastoid cells from siblings discordant for autism spectrum disorders
544 reveals transcriptomic and functional alterations: Evidence for sex-specific effects. *Autism*
545 *Research*, 10(3):439–455, 3 2017.
- 546 [53] Daniel S. Tylee, Jonathan L. Hess, Thomas P. Quinn, Rahul Barve, Hailiang Huang, Yanli
547 Zhang-James, Jeffrey Chang, Boryana S. Stamova, Frank R. Sharp, Irva Hertz-Picciotto,
548 Stephen V. Faraone, Sek Won Kong, and Stephen J. Glatt. Blood transcriptomic comparison
549 of individuals with and without autism spectrum disorder: A combined-samples mega-analysis.
550 *American Journal of Medical Genetics. Part B, Neuropsychiatric Genetics: The Official Pub-*
551 *lication of the International Society of Psychiatric Genetics*, 174(3):181–201, April 2017.
- 552 [54] Daniel S. Tylee, Jonathan L. Hess, Thomas P. Quinn, Rahul Barve, Hailiang Huang, Yanli
553 Zhang-James, Jeffrey Chang, Boryana S. Stamova, Frank R. Sharp, Irva Hertz-Picciotto,
554 Stephen V. Faraone, Sek Won Kong, and Stephen J. Glatt. Blood transcriptomic comparison
555 of individuals with and without autism spectrum disorder: A combined-samples mega-analysis.
556 *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 174(3):181–201, 4
557 2017.
- 558 [55] Daniel S. Tylee, Daniel M. Kawaguchi, and Stephen J. Glatt. On the outside, looking in:
559 a review and evaluation of the comparability of blood and brain "-omes". *American Jour-*
560 *nal of Medical Genetics. Part B, Neuropsychiatric Genetics: The Official Publication of the*
561 *International Society of Psychiatric Genetics*, 162B(7):595–603, October 2013.

- 562 [56] Donna M. Werling, Neelroop N. Parikshak, and Daniel H. Geschwind. Gene expression in
563 human brain implicates sexually dimorphic pathways in autism spectrum disorders. *Nature*
564 *Communications*, 7:10717, 2 2016.
- 565 [57] C Wright, J H Shin, A Rajpurohit, A Deep-Soboslay, L Collado-Torres, N J Brandon, T M
566 Hyde, J E Kleinman, A E Jaffe, A J Cross, and D R Weinberger. Altered expression of
567 histamine signaling genes in autism spectrum disorder. *Translational psychiatry*, 7(5):e1126,
568 5 2017.

List of Figures

569			
570	1	These violin plots show the base-2 logarithm-transformed expression for the two	
571		transcripts with the largest interaction effect from the RNA-Seq data (i.e., those	
572		with the smallest corrected p-value). The solid lines show sex-specific mean ex-	
573		pression differences. The dashed line shows the sex-independent (i.e., pooled) mean	
574		expression difference.	14
575	2	This figure shows the genes with the most significant sex-dependent dysregulation	
576		(i.e., a sex-autism interaction) according to the meta-analysis of the microarray data.	
577		Above, the bar plot shows the χ^2 score for each gene as calculated using Fisher's	
578		method (where the dark bars indicate that the gene has an FDR-adjusted p-value	
579		< 0.05). Below, the dot plot shows whether a gene showed a nominally significant	
580		sex-dependent dysregulation at an unadjusted p-value < 0.05 for a given study. Note	
581		that most genes selected for by the meta-analysis show at least nominal significance	
582		across multiple studies.	15
583	3	These violin plots show base-2 logarithm-transformed expression for the two most	
584		significant main effects (i.e., of the ASD condition) from the RNA-Seq data. The	
585		solid lines show sex-specific mean expression differences. The dashed line shows the	
586		sex-independent (i.e., pooled) mean expression difference.	16
587	4	This figure shows the genes with the most significant sex-independent main effects	
588		(i.e., of the ASD condition) according to the meta-analysis of the microarray data.	
589		Above, the bar plot shows the χ^2 score for each gene as calculated using Fisher's	
590		method (where the dark bars indicate that the gene has an FDR-adjusted p-value	
591		< 0.05). Below, the dot plot shows whether a gene showed a nominally significant	
592		sex-independent main effect at an unadjusted p-value < 0.05 for a given study. Note	
593		that most genes selected for by the meta-analysis show at least nominal significance	
594		across multiple studies.	17
595	5	This dot plot shows results from a GSEA of the RNA-Seq data against the MSigDB	
596		KEGG pathways. For the two sets of results (i.e., the sex-autism interaction and	
597		the main effect), a KEGG pathway (y-axis) has a circle (or triangle) if it is enriched	
598		(or depleted). The size of the points indicates the absolute normalised enrichment	
599		score. The colour indicates the FDR. Note that only points with an FDR < 0.3 are	
600		plotted (see Methods).	18
601	6	This dot plot shows results from a GSEA of the meta-analysis data against the	
602		MSigDB KEGG pathways. For the two sets of results (i.e., the sex-autism interaction	
603		and the main effect), a KEGG pathway (y-axis) has a circle if it is enriched. The	
604		size of the points indicates the absolute normalised enrichment score. The colour	
605		indicates the FDR. Note that only points with an FDR < 0.3 are plotted (see	
606		Methods).	19
607	7	This UpSet plot [33] shows set intersections (and their sizes) from a GSEA of four	
608		results against the MSigDB KEGG pathways. Set identity is indicated by the joined	
609		lines. Set size is indicated by the top bar chart. The bar chart on the left shows the	
610		total set size for each individual GSEA run. Results are filtered using a liberal FDR	
611		threshold of FDR < 0.15 for the RNA-Seq data and FDR < 0.3 for the meta-analysis	
612		data (see Methods).	20




Figure 1: These violin plots show the base-2 logarithm-transformed expression for the two transcripts with the largest interaction effect from the RNA-Seq data (i.e., those with the smallest corrected p-value). The solid lines show sex-specific mean expression differences. The dashed line shows the sex-independent (i.e., pooled) mean expression difference.




Figure 2: This figure shows the genes with the most significant sex-dependent dysregulation (i.e., a sex-autism interaction) according to the meta-analysis of the microarray data. Above, the bar plot shows the χ^2 score for each gene as calculated using Fisher's method (where the dark bars indicate that the gene has an FDR-adjusted p-value < 0.05). Below, the dot plot shows whether a gene showed a nominally significant sex-dependent dysregulation at an unadjusted p-value < 0.05 for a given study. Note that most genes selected for by the meta-analysis show at least nominal significance across multiple studies.




Figure 3: These violin plots show base-2 logarithm-transformed expression for the two most significant main effects (i.e., of the ASD condition) from the RNA-Seq data. The solid lines show sex-specific mean expression differences. The dashed line shows the sex-independent (i.e., pooled) mean expression difference.




Figure 4: This figure shows the genes with the most significant sex-independent main effects (i.e., of the ASD condition) according to the meta-analysis of the microarray data. Above, the bar plot shows the χ^2 score for each gene as calculated using Fisher's method (where the dark bars indicate that the gene has an FDR-adjusted p-value < 0.05). Below, the dot plot shows whether a gene showed a nominally significant sex-independent main effect at an unadjusted p-value < 0.05 for a given study. Note that most genes selected for by the meta-analysis show at least nominal significance across multiple studies.




Figure 5: This dot plot shows results from a GSEA of the RNA-Seq data against the MSigDB KEGG pathways. For the two sets of results (i.e., the sex-autism interaction and the main effect), a KEGG pathway (y-axis) has a circle (or triangle) if it is enriched (or depleted). The size of the points indicates the absolute normalised enrichment score. The colour indicates the FDR. Note that only points with an FDR < 0.3 are plotted (see Methods).




Figure 6: This dot plot shows results from a GSEA of the meta-analysis data against the MSigDB KEGG pathways. For the two sets of results (i.e., the sex-autism interaction and the main effect), a KEGG pathway (y-axis) has a circle if it is enriched. The size of the points indicates the absolute normalised enrichment score. The colour indicates the FDR. Note that only points with an FDR < 0.3 are plotted (see Methods).




Figure 7: This UpSet plot [33] shows set intersections (and their sizes) from a GSEA of four results against the MSigDB KEGG pathways. Set identity is indicated by the joined lines. Set size is indicated by the top bar chart. The bar chart on the left shows the total set size for each individual GSEA run. Results are filtered using a liberal FDR threshold of $FDR < 0.15$ for the RNA-Seq data and $FDR < 0.3$ for the meta-analysis data (see Methods).

List of Tables

613			
614	1	This table details all studies included in the meta-analysis, and the number of probes	
615		available after establishing a final probe set. All subjects with a labelled condition	
616		other than typically developed (TD) were assigned to the autism spectrum disorder	
617		(ASD) group, except for the two Glatt et al. data sets where “Type-1 errors” were	
618		assigned to the TD group.	22
619	2	This table shows SVA-adjusted results for the sex-autism interaction for the RNA-	
620		Seq data (sorted by FDR-adjusted p-value). Note that FDR-adjusted p-values are	
621		also shown for an analysis performed without the adjustment of latent batch effects.	23
622	3	This table shows genes with the most sex-dependent dysregulation (and their chro-	
623		mosomal position), sorted by Fisher score and adjusted p-value. In addition, this	
624		table shows the Fisher score and adjusted p-value calculated for an analysis repeated	
625		without the adjustment of latent batch effects.	24
626	4	This table shows SVA-adjusted results for the main effects (i.e., of the ASD con-	
627		dition) for the RNA-Seq data (sorted by FDR-adjusted p-value). Note that FDR-	
628		adjusted p-values are also shown for an analysis performed without the adjustment	
629		of latent batch effects.	25
630	5	This table shows genes with the most sex-independent dysregulation (and their	
631		chromosomal position), sorted by Fisher score and adjusted p-value. In addition,	
632		this table shows the Fisher score and adjusted p-value calculated for an analysis	
633		repeated without the adjustment of latent batch effects.	26

Study ID	Probes (Intersect)	Females (TD)	Males (TD)	Females (ASD)	Males (ASD)
GSE6575	39561	3	9	8	36
GSE18123	19532	34	48	24	80
Glatt et al. Wave I	28424	28	40	23	88
Glatt et al. Wave II	28424	35	56	28	85
CHARGE	39561	15	75	15	103
Kong et al. 2013	19532	7	10	7	46

Table 1: This table details all studies included in the meta-analysis, and the number of probes available after establishing a final probe set. All subjects with a labelled condition other than typically developed (TD) were assigned to the autism spectrum disorder (ASD) group, except for the two Glatt et al. data sets where “Type-1 errors” were assigned to the TD group.

Transcript ID	Gene symbol	Transcript biotype	Log 2 FC	P-adj SVA	P-adj (no SVA)
ENST00000354042	SLC13A4	protein_coding	3.27	0.293	0.1136846
ENST00000379802	DSP	protein_coding	3.19	0.293	0.6534814
ENST00000262551	OGN	protein_coding	2.97	0.299	0.8169099
ENST00000371625	PTGDS	protein_coding	1.74	0.299	0.0329544
ENST00000223357	AEBP1	protein_coding	1.85	0.529	0.8713166

Table 2: This table shows SVA-adjusted results for the sex-autism interaction for the RNA-Seq data (sorted by FDR-adjusted p-value). Note that FDR-adjusted p-values are also shown for an analysis performed without the adjustment of latent batch effects.

	Location	Fisher	Fisher p-adj	Fisher (no SVA)	Fisher p-adj (noSVA)
TTF2	1p13.1	52.16404	0.0105053	28.88686	1.0000000
UTY	Yq11.221	50.59543	0.0198876	45.76688	0.1378710
KCNJ8	12p12.1	48.17048	0.0528841	38.16932	1.0000000
MAP1B	5q13.2	48.15632	0.0531822	47.94878	0.0578051
RAP2C	Xq26.2	47.70446	0.0637312	24.82099	1.0000000
PAFAH1B1	17p13.3	45.45517	0.1559409	17.84249	1.0000000
CRHR1	17q21.31	44.98624	0.1876599	43.46097	0.3416423
NCS1	9q34.11	44.79693	0.2021903	30.46521	1.0000000
CHST11	12q23.3	44.75342	0.2056750	21.19593	1.0000000
SH3BGR	21q22.2	44.61154	0.2174809	32.59585	1.0000000
BNC2	9p22.3-p22.2	44.40363	0.2360031	39.81245	1.0000000
RORA	15q22.2	43.52113	0.3335702	34.11125	1.0000000
HECA	6q24.1	43.22311	0.3747481	33.12178	1.0000000
FBRSL1	12q24.33	43.04625	0.4015007	35.53452	1.0000000
PAK3	Xq23	43.03339	0.4034965	43.20181	0.3780235
ZC3H7B	22q13.2	42.95711	0.4156536	35.03776	1.0000000
CAMK1D	10p13	42.80430	0.4411269	24.56439	1.0000000
TMED10	14q24.3	42.55614	0.4858196	17.45529	1.0000000

Table 3: This table shows genes with the most sex-dependent dysregulation (and their chromosomal position), sorted by Fisher score and adjusted p-value. In addition, this table shows the Fisher score and adjusted p-value calculated for an analysis repeated without the adjustment of latent batch effects.

Transcript ID	Gene symbol	Transcript biotype	Log 2 FC	P-adj (SVA)	P-adj (no SVA)
ENST00000390930	SNORD17	snoRNA	-2.98	1.54e-05	0.0000102
ENST00000410396	RNU2-2P	snRNA	-4.76	4.04e-05	0.0000000
ENST00000613119		snRNA	-3.23	9.18e-05	0.0000000
ENST00000258526	PLXNC1	protein_coding	0.48	0.00468	0.4273372
ENST00000393775	IGSF11	protein_coding	-1.18	0.00468	1.0000000
ENST00000459255	SCARNA10	snoRNA	-1.71	0.00468	0.0014803
ENST00000618786	RN7SL1	misc_RNA	-1.35	0.0124	0.0026454

Table 4: This table shows SVA-adjusted results for the main effects (i.e., of the ASD condition) for the RNA-Seq data (sorted by FDR-adjusted p-value). Note that FDR-adjusted p-values are also shown for an analysis performed without the adjustment of latent batch effects.

	Location	Fisher	Fisher p-adj	Fisher (no SVA)	Fisher p-adj (noSVA)
ARHGAP35	19q13.32	69.17663	0.0000083	59.97651	0.0004125
GIMAP8	7q36.1	58.39735	0.0008000	52.71485	0.0083436
UCHL3	13q22.2	55.85012	0.0023073	31.88589	1.0000000
SPART	13q13.3	53.75888	0.0054659	43.79029	0.2920570
HNRNPA3P1	10q11.21	53.16493	0.0069742	54.55326	0.0039291
ZNF721	4p16.3	53.02620	0.0073817	45.32102	0.1608751
MAGED2	Xp11.21	52.57098	0.0088931	31.43801	1.0000000
UBE2A	Xq24	52.15816	0.0105264	24.84369	1.0000000
KIF13B	8p12	51.80723	0.0121459	44.99172	0.1830060
POLR1A	2p11.2	51.21815	0.0154371	35.12970	1.0000000
PATL1	11q12.1	50.53892	0.0203385	37.55012	1.0000000
COX19	7p22.3	50.48910	0.0207524	51.68452	0.0126954
GNG5	1p22.3	50.42442	0.0213024	21.04799	1.0000000
HNRNPF	10q11.21	50.20526	0.0232786	52.19956	0.0102957
MUM1	19p13.3	50.09134	0.0243757	38.59229	1.0000000
MTERF4	2q37.3	49.77445	0.0277066	40.36576	1.0000000
KLF1	19p13.13	49.50019	0.0309497	35.07655	1.0000000
SART3	12q23.3	48.93549	0.0388576	51.89275	0.0116656
EIF3A	10q26.11	48.86929	0.0399046	48.66280	0.0429001
ESF1	20p12.1	48.82351	0.0406442	40.26756	1.0000000
TCEAL8	Xq22.1	48.76924	0.0415389	30.32699	1.0000000
RNF168	3q29	47.89156	0.0590766	40.39014	1.0000000
NUCB2	11p15.1	47.52251	0.0684739	46.57846	0.0981743
CCP110	16p12.3	47.21328	0.0774723	30.63996	1.0000000
ZNF569	19q13.12	47.18319	0.0784042	35.01402	1.0000000
CHP1	15q15.1	47.17381	0.0786939	46.71912	0.0928712
ZC3H7B	22q13.2	47.11959	0.0804103	31.75604	1.0000000
GNPDA1	5q31.3	46.86648	0.0889439	39.70348	1.0000000
ECI2	6p25.2	46.83204	0.0901676	54.27612	0.0044030
VCP	9p13.3	46.73363	0.0937667	33.68338	1.0000000
ARHGAP8	22q13.31	46.70772	0.0947338	50.13461	0.0237714
PGM1	1p31.3	46.58133	0.0996154	36.39139	1.0000000
ZNF286A	17p12	46.57586	0.0998268	31.41283	1.0000000

Table 5: This table shows genes with the most sex-independent dysregulation (and their chromosomal position), sorted by Fisher score and adjusted p-value. In addition, this table shows the Fisher score and adjusted p-value calculated for an analysis repeated without the adjustment of latent batch effects.