Urinary sodium is positively associated with urinary free cortisol and total cortisol metabolites in a cross-sectional sample of Australian schoolchildren aged 5-12 years and their mothers

Susan J Torres¹, Carley Grimes¹, Caryl A Nowson¹, Sisitha U Jayasinghe^{1,2}, Clinton R Bruce¹, Shaun A Mason¹, Feng J He³, Anne I Turner¹

¹Deakin University, Geelong, Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences, Victoria, Australia

²University of Lausanne, Department of Physiology, Rue du Bugnon 7 1005, Lausanne, Switzerland

³Centre for Environmental and Preventative Medicine, Wolfson Institute of Preventative Medicine, Queen Mary University of London, London EC1M 6BQ, United Kingdom

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI. 10.1017/S0007114518003136

10.1017/S0007114518003148

Abbreviated title: Sodium intake and urinary cortisol

Key words: sodium, potassium, dietary, urinary, cortisol, children, mothers, Australia

Address for Correspondence:

Dr Anne Turner

Institute for Physical Activity and Nutrition

School of Exercise and Nutrition Sciences

Deakin University

Locked Bag 2000

Geelong, Victoria, 3220

Australia

Phone +61-3-9244 6950

E-mail: anne.turner@deakin.edu.au

Abstract

High sodium intake and chronically elevated cortisol levels are independently associated with the development of chronic diseases. In adults, high sodium intake is associated with high levels of urinary cortisol. We aimed to determine the association between urinary sodium and potassium and urinary cortisol in a cross-sectional sample of Australian schoolchildren and their mothers. Participants were a sample of Australian children $(n \ 120)$ and their mothers $(n \ 100)$ recruited through primary schools. We assessed sodium, potassium, free cortisol and cortisol metabolites in one 24-h urine collection. Associations between 24-h urinary electrolytes and 24-h urinary cortisol were assessed using multilevel mixed-effects linear regression models. In children, urinary sodium was positively associated with urinary free cortisol (B: 0.31 (95% CI 0.19, 0.44)) and urinary cortisol metabolites (B: 0.006, (95% CI 0.002, 0.010)). Positive associations were also observed between urinary potassium and urinary free cortisol (\beta: 0.65, 95% CI 0.23, 1.07)) and urinary cortisol metabolites (\beta: 0.02, 95% CI 0.03, 0.031)). In mothers, urinary sodium was positively associated with urinary free cortisol (β :0.23, 95% CI 0.01, 0.50)) and urinary cortisol metabolites (β :0.008, 95% CI 0.0007, 0.016)). Our findings show that daily sodium and potassium intake were positively associated with cortisol production in children and their mothers. Investigation of the mechanisms involved and the potential impact of sodium reduction on cortisol levels in these populations is warranted.

Introduction

There is evidence that excessive sodium intake is associated with the development of hypertension and cardiovascular disease in adults⁽¹⁾, and that modest reductions in sodium intake can lead to decreases in blood pressure in children⁽²⁾. Recent studies have also shown that sodium intake is positively related to obesity in adults⁽³⁾ and children⁽⁴⁾, independent of energy intake. In most developed countries, adults consume too much sodium and this has also been observed in children, with intakes far exceeding dietary recommendations⁽⁵⁾. Potassium is a nutrient known to have beneficial effects on health in adults including reducing blood pressure and risk of stroke and cardiovascular disease, reducing age-related bone loss and reduction of kidney stones⁽⁶⁾.

It has been suggested that a high sodium diet may lead to the development of chronic diseases via increased production of cortisol as measured in urine⁽⁷⁾. Cortisol, the main glucocorticoid in humans, is an important steroid hormone in the regulation of metabolism and stress, which is produced by activation of the hypothalamo-pituitary adrenal axis (HPA axis)⁽⁸⁾. Cortisol can also be produced in tissues by the conversion of inactive cortisone to active cortisol where it can have direct biological effects at a local level⁽⁹⁾. Elevated cortisol has been linked to the development of numerous diseases in adults including cardiovascular disease, type 2 diabetes and metabolic syndrome⁽¹⁰⁾. In a large cross-sectional study of adult men and women, hair cortisol, as a marker of long term cortisol exposure, was positively associated with weight, BMI and waist circumference, and in a retrospective longitudinal analysis was found to be associated with obesity over 4 years⁽¹¹⁾. The potential detrimental effects of excess cortisol levels have also been observed in paediatric populations. For example, greater increases in salivary cortisol in response to mild psychological stress in children was associated with adverse effects on memory function⁽¹²⁾. Recent evidence shows that cortisol is elevated in children and adolescents with depression⁽¹³⁾ and in obese prepubertal girls⁽¹⁴⁾.

There is evidence in adults to suggest that sodium intake is positively associated with urinary cortisol concentrations. For example, in experimental studies conducted in adults, urinary free cortisol increases after consuming high sodium diets of 320 mEq/day⁽¹⁵⁾ and 200 mmol/day⁽¹⁶⁾ and decreases after restricting sodium intake to 20 mmol/day⁽¹⁷⁾ and 10 mmol/day⁽¹⁸⁾. While one study found no evidence of an association between 24-h urinary sodium and cortisol levels, this study measured cortisol levels only in serum rather than

urine⁽¹⁹⁾. These findings are consistent with those of Baudrand and colleagues who measured cortisol in both plasma and urine⁽⁷⁾. These investigators showed that urinary free cortisol and total cortisol metabolite levels in urine but not plasma were significantly higher in participants with high sodium intake compared with those with an adequate sodium intake⁽⁷⁾. These latter two studies suggest that it is not sufficient to measure cortisol in plasma/serum as these levels remain well controlled within a stable homeostatic range. Instead, it appears necessary to measure cortisol levels in urine. While the levels of cortisol in plasma would intuitively seem to be the most important, as these are what the tissues are exposed to, when cortisol produced occurs within tissues, it can have detrimental effects at a local level before entering the systemic circulation and being cleared rapidly by the kidney. Measures of both urinary free cortisol and total cortisol metabolites are necessary to provide a complete picture of urine cortisol excretion, which is thought to be indicative of overall daily cortisol production⁽⁷⁾. Although there is evidence of an association between sodium intake and urinary cortisol in adults, no studies have examined this association in children.

Thus, the primary objective of this study was to test the hypothesis that 24-h urinary sodium is positively associated with 24-h urinary free cortisol and total cortisol metabolites in a sample of Australian schoolchildren aged 5-12 years and their mothers. Given the known health benefits of higher potassium intake, and the lack of data available on potassium and cortisol in adults and children, we further hypothesised that children and their mothers would show a negative association between 24-h urinary potassium and urinary free cortisol and total cortisol metabolites. This was part of a larger cross-sectional study in which we have previously reported a positive association between sodium intake and obesity risk in Australian schoolchildren ⁽⁴⁾.

Methods

Study design and participants

In this study, we measured urinary free cortisol and cortisol metabolites in 24-h urine samples in a sub-group of primary schoolchildren and their mothers who had participated in the Salt and Other Nutrient Intakes in Children (SONIC) study, which included children and their parents sampled from primary schools located in Victoria, Australia from 2009-2013. A detailed description of the study protocol and the primary results have been reported previously^(20,21). The initial sample size for this study was determined by the number of

mothers with available sodium data, which was 107 mothers. If two or more children from one family had participated in the SONIC study, we randomly selected one child per mother to be included. Five mothers were subsequently excluded because cortisol data could not be obtained, but we chose to keep their child in the present study. As we wished to look at the association between urinary sodium and urinary cortisol across the widest possible range of sodium intakes in children, we included an additional 7 children with sodium levels (249-310 mmol/24-h) higher than the range of the already selected children (31-245 mmol/24-h) and 6 children with sodium levels lower than this selected range (13-30 mmol/24-h). Therefore, our sample included 102 mother-child pairs and a total of 120 children.

The study was conducted according to the guidelines laid down in the Declaration of Helsinki. Mothers and children provided written informed consent before commencing the study, which was approved by the Deakin University Human Ethics Committee (Project No: EC 62-2009).

Measures

Socio-Economic status (SES) was based on school postcode, as previously defined⁽⁴⁾.

24-hour urine collection

The 24-h urine collection protocol has been described elsewhere^(4,21). In brief, at the start of the 24-h urine collection, children and mothers were instructed to empty their bladder, discard this urine and note this as the start time of their collection. Following this, all urine voided was collected until the 24-h finish time. Urine samples were considered incomplete and not included in the final analysis if collection time was <20 h or >28 h, total volume was <300ml, the participant reported missing >1 collection, or urinary creatinine for children were <0.1 mmol/kg body weight per day⁽²¹⁾. Mothers with suspected inaccurate urine collections (i.e. urinary creatinine <4 mmol/day for women, or a 24-h urine collection of <500 mL) were excluded.

Aliquots of urine (2ml) that were stored at -80°C were thawed under room temperature and utilised for free cortisol and cortisol metabolite analyses.

Urinary free cortisol analysis

Concentrations of urinary free cortisol were measured using an enzyme-linked immunosorbent assay (Diametra, Milano, Italy). Seven assays were conducted with the lowest detection sensitivity of 2.95 ng/mL at the 95% confidence limit. The intra-assay coefficient of variation was 4.8 % at 47 ng/mL and 8.0 % at 287 ng/mL. The inter-assay coefficient of variation was 10.8 % at 47 ng/mL and 7.9 % at 287 ng/mL.

Extraction of steroids and measurement of cortisol metabolites by gas chromatography–mass spectrometry (GC-MS)

Steroids were extracted from urine samples and measured using GC-MS based on previously published methods⁽²²⁾ with some modifications. Briefly, deuterated internal standards, tetrahydrocortisone-D₅ (THE-D₅) and tetrahydrocortisol-D₅ (THF-D₅; Sigma Aldrich, Castle Hill, NSW, Australia), were added to samples followed by hydrolysis, extraction, and derivitazation steps described previously⁽²²⁾. Nitrogen-dried samples were reconstituted in cyclohexane/pyridine/hexamethyldisilazane (98:1:1) and transferred to vials for GC-MS.

The GC-MS system consisted of Agilent components: a 6890 GC, a 7683 automated sampler and a 5975C mass selective detector. A 30 m x 0.25 mm (I.D.) x 0.25 μ m capillary column (VF-Xms; Agilent Technologies) was used with helium as the carrier gas (flow rate 1.1 ml/min). The GC oven program consisted of an initial hold temperature of 70°C for 1.5 minutes, a 25°C/min ramp to 220°C, a 2°C/min ramp to 276°C, a 25°C/min ramp to 320°C, and a final hold time for 3 min at 320°C. Data recording and analysis occurred using MSD Chemstation software (Agilent Technologies). Endogenous cortisol metabolites, α -cortolone, β -cortolone, α -cortol, β -cortol, tetrahydrocortisone (THE), tetrahydrocortisol (THF) and allotetrahydrocortisol (α -THF) and deuterated internal standards were identified using selective ion monitoring. External standard curves were run daily for each of the target metabolites for purposes of relative quantification. Each metabolite was quantitatively corrected for its own or its closest related deuterated internal standard. The total cortisol metabolites in each sample were calculated as the sum of each individual cortisol metabolite concentration, corrected for the total urine collection volume.

Urinary sodium and potassium analysis

Returned 24-h urine samples were transported to an accredited commercial pathology laboratory (Dorevitch Pathology) for analysis. Urinary sodium (Na) and potassium (K) concentrations were assessed using indirect ion selective electrodes, and urinary creatinine

concentration was assessed using the Jaffe reaction⁽²³⁾ on the Siemens Advia 2400 analyser (Siemens Healthcare). Approximately 90-95% of ingested sodium is excreted in urine⁽²⁴⁾. Because of the high recovery of sodium in urine, 24-hour urine is considered the "gold standard" method to determine dietary sodium intake⁽²⁵⁾. The coefficient of variation for Na and K was <1% and for creatinine it was 3.25%. If the duration of the collection was not exactly 24-h but within 20–28 h, urinary electrolytes, creatinine, urinary free cortisol, urinary cortisol metabolites and total volume were standardised to a 24-h period (i.e. (24-h/urine duration (h))×urinary measure).

Anthropometric measures

Height and weight of children were measured by trained research staff following standard protocols^(4,21). BMI was calculated as weight (kilograms) divided by height (metres) squared. BMI values were converted to age- and sex-adjusted BMI z scores using the 2000 US Centers for Disease Control and Prevention growth charts^(26,27). No anthropometry data were available for the mothers.

Statistical analysis

Descriptive statistics (mean values and standard deviations or numbers and percentages) were calculated to describe the participant characteristics. Differences between children and mothers 24-h urinary electrolytes and 24-h urinary cortisol were assessed using Mann-Whitney U Test. Multilevel mixed-effects linear regression was used to determine the association between urinary Na or urinary K and urinary free cortisol or total cortisol metabolites. All models included a random intercept for school. In children, unadjusted models and models adjusting for age, gender, urinary sodium or urinary potassium are presented. In an additional model, we also adjusted for BMI z-score because obesity may influence the association between urinary electrolytes and urinary cortisol. In mothers, unadjusted models and models adjusting for age, urinary sodium or urinary potassium were fitted. Additionally, the associations were tested between mothers and children for urinary electrolytes and urinary cortisol. Model residuals were assessed for normality and heteroscedasticity using Q-Q plots and plots of residuals against fitted values.

Based on prior research⁽⁷⁾, we estimated an effect size of $r^2=0.17$ for the relationship between urinary sodium and urinary free cortisol and $r^2=0.10$ for the relationship between urinary sodium and urinary cortisol metabolites. Power analyses conducted separately for children (n

= 120) and mothers (n = 100) showed that we had over 90% power to detect these effects, for alpha=0.05. All statistical analysis, including power analysis, was conducted using SPSS 24.0 for Windows (SPSS. Inc.) and STATA SE, version 15 (1985-2017; StataCorp).

Results

Participants

Table 1 shows the demographic characteristics and 24-h urinary electrolytes and cortisol for children and mothers. There was one mother excluded based on an extremely low urinary sodium concentration (10 mmol/24-h, >3 SD below the mean) and one mother excluded with an extremely high urinary free cortisol concentration (529 mmol/24-h, >3 SD above the mean). The final sample consisted of 120 children and 100 mothers. There was no difference in age, sex, SES or BMI z-score between those children in this sub-study compared with those in the original SONIC study (data not shown, P>0.05 for all)⁽⁴⁾. Overall, 53% of children were girls and had an average age of 9.2 years. In total, 13% of children were either overweight or obese. With respect to sodium intake in the children, urinary sodium ranged from 13 to 310 mmol/day with a mean of 104 mmol/day. There were 45 (38%) children who were at or below the recommended dietary sodium upper level and 75 (63%) children who exceeded the upper level (upper level: 4-8 years 60 mmol/day, 9-13 years 86 mmol/day)⁽²⁸⁾. The average age of mothers was 41.7 years. In mothers, urinary sodium ranged from 47 to 237 mmol/day with a mean of 121 mmol/day. There were 36 (36%) mothers who were at or below the recommended sodium upper level for adults of 100 mmol/day)⁽²⁸⁾ and 64 (64%) mothers who exceeded the upper level. Compared to children, the mothers had significantly higher levels of urinary electrolytes, urinary free cortisol and total cortisol metabolites (Table 1).

Associations between urinary electrolytes and urinary cortisol

Fig. 1 shows scatterplots for children's 24-h urinary electrolytes and urinary cortisol and cortisol metabolites. Table 2 shows the multilevel mixed-effects linear regression models for children with urinary electrolytes as the independent variables and urinary free cortisol and urinary cortisol metabolites as the dependent variables. In children, urinary sodium and potassium were both positively associated with urinary free cortisol and urinary cortisol metabolites, and this relationship persisted after adjusting for relevant confounders (age, sex, urinary sodium, urinary potassium and BMI z-score). Results from the fully adjusted models

show that a 1 mmol/24-h increase in urinary sodium was related with a 0.31 nmol/24-h (95% CI 0.19, 0.44) higher urinary free cortisol, and a 1 mmol/24-h increase in urinary potassium was related with a 0.65 nmol/24-h (95% CI 0.23, 1.07) higher urinary free cortisol (Fig. 1). There was no evidence of an association between sodium to potassium ratio and either urinary free cortisol or urinary cortisol metabolites (data not shown).

In mothers, urinary sodium was positively associated with urinary free cortisol and cortisol metabolites in both the unadjusted and adjusted models (Table 2 and Fig. 2). Urinary potassium was positively associated with urinary cortisol metabolites in the unadjusted model but not in the adjusted model and urinary potassium was not associated with urinary free cortisol in unadjusted or adjusted models. In mothers, urinary sodium was positively associated with urinary sodium was positively associated with urinary potassium (r=0.266, P<0.05).

Within mother-child dyads, urinary sodium, free cortisol and cortisol metabolites were positively associated (P<0.05) between mothers and children's, but urinary potassium was not (Table 3).

Discussion

Among a sample of Australian schoolchildren, we have reported novel data for the first time that higher 24-h urinary sodium and potassium levels were associated with higher 24-h urinary free cortisol and urinary cortisol metabolites. In mothers, urinary sodium was positively associated with both urinary free cortisol and cortisol metabolites, while urinary potassium was not associated with urinary free cortisol or cortisol metabolites. Associations in children remained after adjusting for age, gender, BMI zscore, and urinary sodium or urinary potassium. Overall, the magnitude of the associations was greatest in the children. We also found that urinary sodium, free cortisol and cortisol metabolites were positively associated between children and mothers. It should be noted that causation cannot be established due to the cross-sectional nature of this study.

The level of urinary free cortisol in the children in the present study was similar to that in a population of obese children⁽²⁹⁾, but higher than that in a population of healthy children⁽³⁰⁾. Urinary free cortisol levels in the mothers in the present study were similar to those reported by Baudrand et al⁽⁷⁾, however, cortisol metabolite levels were lower. Differences between studies may reflect inter-laboratory differences in methodologies, which has been reported in the analysis of hair cortisol concentrations.⁽³¹⁾

In the present study, our findings suggest for children that a high sodium diet is associated with an increase in daily production of cortisol. As previously reported, elevated cortisol levels have been linked to the development of chronic diseases in paediatric populations $^{(12,13,14)}$. Our findings show that dietary sodium appears to be a factor modifying cortisol production in both paediatric and adult populations. Our findings in mothers are consistent with evidence from studies in adults where high sodium intake was also found to be positively associated with free urinary cortisol and urinary cortisol metabolites⁽⁷⁾. Likewise, in two studies in men that went from a low sodium (10 mmol/day) to high sodium diet (200 mmol/day), urinary cortisol increased by 65 nmol/day⁽¹⁶⁾ and 20 mcg/day⁽¹⁸⁾. Although there is evidence to suggest sodium intake is linked to cortisol production, the precise mechanisms remain to be elucidated. The main pathways of cortisol metabolism include two isoforms of the 11β-hydroxysteroid dehydrogenase (11βHSD), 11βHSD1 and 11βHSD2, which are found in many human tissues⁽⁹⁾. 11βHSD1 converts inactive cortisone to cortisol and 11βHSD2

increased in adipose tissue, resulting in greater production of the active hormone corticosterone (the main glucocorticoid found in rodents), providing support for how sodium might influence cortisol secretion⁽³²⁾. Currently, there is no evidence for an equivalent mechanism in a human model.

Our findings that urinary potassium was positively associated with urinary free cortisol and cortisol metabolites in children are novel as this has not been reported before in this population group. However, these findings were contrary to our hypothesis where we predicted a negative association between urinary potassium and urinary free cortisol and total cortisol metabolites based on the reported health benefits of potassium. Nevertheless, feeding studies in rats have investigated a potential mechanism for how dietary potassium intake might influence corticosterone levels. Feeding rats a diet high in potassium chloride resulted in increased production of 11BHSD2 protein in the renal distal tubules, catalysing the conversion of the active corticosterone to the inactive 11-dehydrocorticosterone ⁽³³⁾. This results in greater specificity for renal distal tubule mineralocorticoid receptors to aldosterone thereby promoting aldosterone-induced potassium excretion. In contrast, Thompson et al found that a high potassium diet was associated with reduced renal 11BHSD2 activity in rats⁽³⁴⁾, thus, enhancing the access of glucocorticoids to the mineralocorticoid receptor to promote urinary potassium excretion. While these two studies show that potassium may be involved in 11BHSD2 regulation in the renal distal tubule, and therefore glucocorticoid excretion, the discordant findings suggest that further research is required in this area. Further studies are needed to determine the relevant mechanisms present in humans and any potential physiological or health implications.

It is important to consider the clinical relevance of our findings. We found that 1 mmol/24-h of urinary sodium was related to a 0.31 nmol/24-h higher urinary free cortisol and that 1 mmol/24-h of urinary potassium was related to a 0.65 nmol/24-h higher urinary free cortisol. While reference ranges for cortisol are available for the diagnosis of clinical conditions such as Cushing's syndrome,⁽³⁵⁾ clinically meaningful levels for cortisol in terms of biological outcomes for non-clinical populations are not well understood. Studies in adults have found that decreasing sodium by 180 mmol/24-h reduced urinary free cortisol by 40 nmol/24-h⁽¹⁷⁾ and 61 nmol/24-h⁽¹⁸⁾. It is not likely, however, that such large reductions in sodium would be achievable in the general population. In the present study, the average sodium intake in the children was 104 mmol/day (compared to an upper level of 86 mmol/day, 9-13 years) and the

average sodium intake in the mothers was 121 mmol/day (compared to an upper level of 100 mmol/day)⁽²⁸⁾. Overall, clinically meaningful reductions in urinary free cortisol are not well understood in paediatric or adult populations, therefore it is difficult to recommend a reduction in urinary sodium and potassium; further research in this area is warranted.

In the present study, we found that urinary free cortisol and cortisol metabolites were positively associated between children and mothers. This is consistent with results from a study that found high maternal hair cortisol was strongly associated with their infant's evening salivary cortisol levels⁽³⁶⁾. In a sample of mothers and their children living in socioeconomically disadvantaged neighbourhoods, hair cortisol levels were also strongly positively associated between mothers and their children⁽³⁷⁾. Similarly, Pratt et al reported concordance for diurnal cortisol secretion between mothers and their 6-year-old children⁽³⁸⁾. Tarullo et al have proposed that this physiological concordance between mother and child has an evolutionary basis, such that having a similar response is adaptive when facing the same external risk⁽³⁶⁾.

There are limitations associated with this study, which must be considered when interpreting the findings. The cross-sectional design of the study prevents the detection of causal relationships between dietary electrolyte intake and cortisol levels. We did not obtain any anthropometric measures in the mothers, so we were unable to adjust for these in the regression analyses. While the association between sodium or potassium and cortisol may be influenced by the types of foods and nutrients consumed by the children and mothers, we were unable to assess this in our study. The convenience sample of schoolchildren potentially limits the generalisability of the findings to a wider population. We only assessed mothers in this study as the relationship between urinary electrolytes and urinary cortisol and cortisol metabolites between mother and child may differ to father and child. A strength of this study was that both urinary cortisol and cortisol metabolites were measured to fully characterise cortisol production.

In conclusion, in a population of schoolchildren and their mothers, this study has shown that high sodium and potassium intake is associated with increased production of cortisol. The negative health implications of elevated cortisol exposure are well documented. Future research should include metabolic studies to investigate mechanisms that might be involved and the effect of sodium reduction on cortisol levels in these target populations.

Acknowledgements

We thank Dr Catherine Huggins for her early work in securing project funding. We acknowledge the Victorian Department of Early Childhood and Development for their support in allowing this study to be conducted within the government school sector. We thank all schools, children and mothers who participated in the study.

C. A. G., C. A. N. and F. J. H. designed and conducted the original SONIC study. A. I. T., S. J. T. and C. A. N. designed and conducted this salt and cortisol sub-study including securing project funding. S. U. J. was responsible for the free urinary cortisol analysis. C. R. B. and S.A.M. were responsible for the urinary cortisol metabolite analysis. S. J. T. analysed the data and drafted the manuscript. All authors critically reviewed the manuscript and read and approved the final version. None of the authors had a conflict of interest.

Financial disclosure

This study was funded by a National Heart Foundation of Australia Grant-in-Aid (G 10M 5021), a Helen MacPherson Smith Trust Fund Project Grant (6002), a Deakin University Faculty of Health Research Development Grant and by seed funding from Deakin University's Institute for Physical Activity and Nutrition. C.R.B. is supported by an Australian Research Council Future Fellowship (FT160100017).

Table 1. Descriptive characteristics of participants (mean values and standard deviations; or

percentages)

	Child	ren	Mothers		
	(<i>n</i> 12	20)	(<i>n</i> 100)		
Characteristics	Mean	SD	Mean	SD	
Age (years)	9.2	2.0	41.7	5.1	
Age range (years)	(5.0-12.8)		(23.3-52.4)		
Female	63		100		
n					
%	53		100		
BMI (kg/m ²)	17.3	3.0	NA		
BMI (z score)	0.1	1.0	NA		
Weight category					
Underweight (%)	10.	0	NA		
Healthy weight (%)	76.	7	NA		
Overweight (%)	11.	7	NA		
Obese (%)	1.7	7	NA		
Urinary sodium (mmol/24-h)	104.2	61.8	121.4	40.1*	
Urinary potassium (mmol/24-h)	46.8	18.5	67.7	18.3*	
Urinary creatinine (mmol/24-h)	5.4	2.2	10.1	2.2*	
Urinary free cortisol (nmol/24-h)	96.2	45.8	114.8	49.0*	
Total cortisol metabolites (mg/24-h)	3.0	1.5	4.3	1.6*	

NA, Not available; BMI, Body Mass Index

*P < 0.05 Mann-Whitney U Test

Table 2. The association between urinary electrolytes and urinary cortisol in schoolchildren and mothers

(Regression coefficients and 95% confidence intervals)

		Urinary free cortisol			Urinary cortisol metabolites			
Children (n 120)								
		β	95% CI	Р	β	95% CI	Р	
Urinary sodium	Model 1*	0.43	0.32, 0.54	< 0.001	0.01	0.008, 0.015	< 0.001	
	Model 2†	0.29	0.17, 0.40	< 0.001	0.006	0.003, 0.010	< 0.001	
	Model 3¶	0.31	0.19, 0.44	< 0.001	0.006	0.002, 0.010	< 0.01	
Urinary potassiu	m							
	Model 1*	1.36	0.99, 1.73	< 0.001	0.04	0.03, 0.05	< 0.001	
	Model 2 [‡]	0.60	0.18, 1.02	< 0.01	0.02	0.004, 0.003	0.01	
	Model 3**	0.65	0.23, 1.07	< 0.01	0.02	0.003, 0.031	0.015	
Mothers (n 100)								
Urinary sodium								
	Model 1*	0.26	0.03, 0.50	0.026	0.009	0.002, 0.017	0.012	
	Model 2§	0.23	0.01, 0.50	0.042	0.008	0.0007, 0.016	0.032	
Urinary potassiu	m							
	Model 1*	0.32	-0.20, 0.83	0.233	0.017	-0.0001, 0.033	0.047	
	Model 2	0.14	-0.41, 0.70	0.618	0.015	-0.002, 0.032	0.087	

*Model 1: unadjusted

†Model 2: adjusted for age, sex, and urinary potassium

[‡]Model 2: adjusted for age, sex and urinary sodium

§Model 2: adjusted for age and urinary potassium (N=94 included in analysis)

Model 2: adjusted for age and urinary sodium (N=94 included in analysis)

Model 3: adjusted for age, sex, urinary potassium and BMI z score

**Model 3: adjusted for ages, sex, urinary sodium and BMI z score

Table 3 Pairwise associations between urinary electrolytes and urinary cortisol in mother-child dyads (N=100)(Regression coefficients and 95% confidence intervals)

Outcome	Predictor		Unadjusted model			
Chil	d Mother	β	95% CI	Р		
Sodiur	n Sodium	0.25	0.06, 0.43	< 0.001		
Potassiu	n Potassium	0.06	-0.11, 0.22	0.514		
Urinary free cortise	l Urinary free cortisol	0.26	0.11, 0.41	< 0.001		
Urinary cortisol metabolite	s Urinary cortisol metabolites	0.19	0.03, 0.36	0.023		

References

1. Kotchen TA, Cowley AW, Jr., Frohlich ED (2013) Salt in health and disease--a delicate balance. *N Engl J Med* **368**, 1229-1237.

2. He FJ, MacGregor GA (2006) Importance of salt in determining blood pressure in children: meta-analysis of controlled trials. *Hypertension* **48**, 861-869.

3. Ma Y, He FJ, MacGregor GA (2015) High salt intake: independent risk factor for obesity? *Hypertension* **66**, 843-849.

4. Grimes CA, Riddell LJ, Campbell KJ *et al.* (2016) 24-h urinary sodium excretion is associated with obesity in a cross-sectional sample of Australian schoolchildren. *Br J Nutr* **115**, 1071-1079.

5. Brown IJ, Tzoulaki I, Candeias V *et al.* (2009) Salt intakes around the world: implications for public health. *Int J Epidemiol* **38**, 791-813.

6. Weaver CM (2013) Potassium and health. Advances In Nutrition 4, 368S-377S.

7. Baudrand R, Campino C, Carvajal CA *et al.* (2014) High sodium intake is associated with increased glucocorticoid production, insulin resistance and metabolic syndrome. *Clin Endocrinol* **80**, 677-684.

8. Folkow B (1997) Physiological aspects of the "defence" and "defeat" reactions. *Acta Physiol Scand Suppl* 640, 34-37.

9. Tomlinson JW, Stewart PM (2001) Cortisol metabolism and the role of 11beta-hydroxysteroid dehydrogenase. *Best Pract Res Clin Endocrinol Metab* **15**, 61-78.

10. Chrousos GP (2009) Stress and disorders of the stress system. Nat Rev Endocrinol 5, 374-381.

11. Jackson SE, Kirschbaum C, Steptoe A (2017) Hair cortisol and adiposity in a population-based sample of 2,527 men and women aged 54 to 87 years. *Obesity* **25**, 539-544.

12. Heffelfinger AK, Newcomer JW (2001) Glucocorticoid effects on memory function over the human life span. *Dev Psychopathol* **13**, 491-513.

13. Stetler C, Miller GE (2011) Depression and hypothalamic-pituitary-adrenal activation: a quantitative summary of four decades of research. *Psychosom Med* **73**, 114-126.

14. Papafotiou C, Christaki E, van den Akker EL *et al.* (2017) Hair cortisol concentrations exhibit a positive association with salivary cortisol profiles and are increased in obese prepubertal girls. *Stress* **20**, 217-222. 15. Wambach G, Bleienheuft C, Bonner G (1986) Sodium loading raises urinary cortisol in man. *J Endocrinol Invest* **9**, 257-259.

16. Litchfield WR, Hunt SC, Jeunemaitre X *et al.* (1998) Increased urinary free cortisol: a potential intermediate phenotype of essential hypertension. *Hypertension* **31**, 569-574.

17. Lewicka S, Nowicki M, Vecsei P (1998) Effect of sodium restriction on urinary excretion of cortisol and its metabolites in humans. *Steroids* **63**, 401-405.

18. Chamarthi B, Kolatkar NS, Hunt SC *et al.* (2007) Urinary free cortisol: an intermediate phenotype and a potential genetic marker for a salt-resistant subset of essential hypertension. *J Clin Endocrinol Metab* **92**, 1340-1346.

19. Afsar B, Ay M (2014) The relationships between cortisol levels, insulin levels, and thyroid hormones with 24-h urinary sodium excretion in never treated essential hypertensive patients. *ARYA Atheroscler* **10**, 159-163.

20. Grimes CA, Riddell LJ, Campbell KJ *et al.* (2017) Dietary intake and sources of sodium and potassium among Australian schoolchildren: results from the cross-sectional Salt and Other Nutrients in Children (SONIC) study. *BMJ Open* **7**, e016639.

21. Grimes CA, Baxter JR, Campbell KJ *et al.* (2015) Cross-Sectional Study of 24-Hour Urinary Electrolyte Excretion and Associated Health Outcomes in a Convenience Sample of Australian Primary Schoolchildren: The Salt and Other Nutrients in Children (SONIC) Study Protocol. *JMIR Res Protoc* **4**, e7.

22. Honour JW (1997) Steroid profiling. Ann Clin Biochem 34 (Pt 1), 32-44.

23. Jaffe M (1886) Ueber den Niederschlag, welchen Pikrinsäure in normalem Harn erzeugt und über eine neue Reaction des Kreatinins. *Z Physiol Chem* **10**, 391-400.

24. Bates CJ TD, Bingham SA, Margetts BM, Nelson M (1997) Biochemical markers of nutrient intake. In *Design Concepts in Nutritional Epidemiology*, 2nd ed., pp. 170-240 [Margetts BM and Nelson M, editors]. Oxford, UK: Oxford University Press, Inc.

25. World Health Organization (2010) Strategies to monitor and evaluate population sodium consumption and sources of sodium in the diet. Geneva, Switzerland: World Health Organization.

26. Cole TJ, Flegal KM, Nicholls D *et al.* (2007) Body mass index cut offs to define thinness in children and adolescents: international survey. *BMJ* **335**, 194.

27. Cole TJ, Bellizzi MC, Flegal KM *et al.* (2000) Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* **320**, 1240-1243.

28. Department of Health and Ageing, National Health and Medical Research Council (2006) *Nutrient Reference Values for Australia and New Zealand. Including Recommended Dietary Intakes.* Canberra: Commonwealth of Australia.

29. Reinehr T, Kulle A, Wolters B *et al.* (2014) Relationships between 24-hour urinary free cortisol concentrations and metabolic syndrome in obese children. *J Clin Endocrinol Metab* **99**, 2391-2399.

30. Shi L, Berkemeyer S, Buyken AE *et al.* (2010) Glucocorticoids and body fat associated with renal uric acid and oxalate, but not calcium excretion, in healthy children. *Metabolism* **59**, 134-139.

31. Stalder T, Steudte-Schmiedgen S, Alexander N *et al.* (2017) Stress-related and basic determinants of hair cortisol in humans: A meta-analysis. *Psychoneuroendocrinology* **77**, 261-274.

32. Usukura M, Zhu A, Yoneda T *et al.* (2009) Effects of a high-salt diet on adipocyte glucocorticoid receptor and 11-beta hydroxysteroid dehydrogenase 1 in salt-sensitive hypertensive rats. *Steroids* **74**, 978-982.

33. Hermans JJ, Fischer MA, Schiffers PM *et al.* (1999) High dietary potassium chloride intake augments rat renal mineralocorticoid receptor selectivity via 11beta-hydroxysteroid dehydrogenase. *Biochim Biophys Acta* **1472**, 537-549.

34. Thompson A, Bailey MA, Michael AE *et al.* (2000) Effects of changes in dietary intake of sodium and potassium and of metabolic acidosis on 11beta-hydroxysteroid dehydrogenase activities in rat kidney. *Exp Nephrol* **8**, 44-51.

35. Pappachan JM, Hariman C, Edavalath M *et al.* (2017) Cushing's syndrome: a practical approach to diagnosis and differential diagnoses. *J Clin Pathol* **70**, 350-359.

36. Tarullo AR, St John AM, Meyer JS (2017) Chronic stress in the mother-infant dyad: Maternal hair cortisol, infant salivary cortisol and interactional synchrony. *Infant Behav Dev* **47**, 92-102.

37. Olstad DL, Ball K, Wright C *et al.* (2016) Hair cortisol levels, perceived stress and body mass index in women and children living in socioeconomically disadvantaged neighborhoods: the READI study. *Stress* **19**, 158-167.

38. Pratt M, Apter-Levi Y, Vakart A *et al.* (2017) Mother-child adrenocortical synchrony; Moderation by dyadic relational behavior. *Horm Behav* **89**, 167-175.

(a)

(b)




Figure 1. Scatterplots for children's 24-h urinary electrolytes and 24-h urinary cortisol (n=120). Scatterplots for (a) urinary sodium and urinary free cortisol, (b) urinary sodium and urinary cortisol metabolites, (c) urinary potassium and urinary free cortisol, (d) urinary potassium and urinary cortisol metabolites. Data were analysed with multilevel mixed-effects linear regressions adjusted for age, sex and BMI z score. Additionally adjusted for urinary potassium (a) and (b), and urinary sodium (c) and (d).




Figure 2. Scatterplots for mother's 24-h urinary electrolytes and 24-h urinary cortisol (n=100). Scatterplots for (a) urinary sodium and urinary free cortisol, (b) urinary sodium and urinary cortisol metabolites, (c) urinary potassium and urinary free cortisol, (d) urinary potassium and urinary cortisol metabolites. Data were analysed with multilevel mixed-effects linear regressions adjusted for age. Additionally adjusted for urinary potassium (a) and (b), and urinary sodium (c) and (d).