2018 INAUGURAL REVIEW PRIZE WINNER



Mitochondrial regulation in skeletal muscle: A role for non-coding RNAs?



Institute for Physical Activity and Nutrition (IPAN), Deakin University, Geelong, Victoria, Australia

Correspondence

Séverine Lamon, Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences, 221 Burwood Highway, Burwood 3125, Australia

Email: severine.lamon@deakin.edu.au

Funding

S.L. was supported by a Discovery Early Career Research Award (DECRA) from the Australian Research Council (ARC) (DE150100538). J.S. was supported by a writing scholarship from the School of Exercise and Nutrition Sciences.

Edited by: Jeremy Ward

Abstract

Skeletal muscle is a highly metabolic tissue characterized by high mitochondrial abundance. As such, skeletal muscle homeostasis relies on the tight control of mitochondrial gene expression to ensure efficient mitochondrial function. Mitochondria retain a conserved genome from prokaryotic ancestors, and mitochondrial gene regulation relies on communication between mitochondrial- and nuclear-encoded transcripts. Small and long non-coding RNAs (ncRNAs) have regulatory roles in the modulation of gene expression. Emerging evidence demonstrates that regulatory ncRNAs, particularly microRNAs (miRNAs) and long ncRNAs (IncRNAs), localize within the mitochondria in diverse physiological and pathological states. These molecules present intriguing possibilities for the regulation of mitochondrial gene expression. Current research suggests that all known miRNAs are encoded by the nuclear genome but can target mitochondrial genes. Initial investigations demonstrate direct interactions between the muscleenriched miR-1 and miR-181c and mitochondrial transcripts, suggesting advanced roles of miRNAs in mitochondrial gene regulation. This review draws evidence from the current literature to discuss the hypothesis that a level of ncRNA-mediated gene regulation, in particular miRNAmediated gene regulation, takes place in the mitochondria. Although ncRNA-mediated regulation of the mitochondrial genome is a relatively unexplored field, it presents exciting possibilities to further our understanding of mitochondrial metabolism and human muscle physiology.

KEYWORDS

mitochondria, non-coding RNA, skeletal muscle

1 | INTRODUCTION

Mitochondria are the primary regulators of energy metabolism in the cell. As such, they play a central role in cellular homeostasis (Herst, Rowe, Carson, & Berridge, 2017; Russell, Foletta, Snow, & Wadley, 2014). In mammalian cells, mitochondria form a dynamic system that relies on the balance between mitochondrial biogenesis and mitochondrial degradation, termed mitophagy (Herst et al., 2017; Yan, Lira, & Greene, 2012). Mitochondria constantly join and separate this reticular network through processes referred to as mitochondrial fusion and mitochondrial fission (Herst et al., 2017). Most eukaryotic tissues contain mitochondria, and differences in mitochondrial abundance reflect tissue-specific energy demands (Little, Safdar, Benton, & Wright, 2011).

Skeletal muscle is a highly metabolic tissue (Zurlo, Larson, Bogardus, & Ravussin, 1990) that is characterized by high-energy needs and high mitochondrial content (Fernández-Vizarra, Enríquez, Pérez-Martos, Montoya, & Fernández-Silva, 2011; Ogata & Yamasaki, 1997). Skeletal muscle mitochondrial content and activity are key determinants of whole-body metabolism and exercise performance (Groennebaek & Vissing, 2017; Nunnari & Suomalainen, 2012). Numerous diseases and conditions, including myopathies, amyotrophic lateral sclerosis, type 2 diabetes and ageing, are associated with impaired mitochondrial content and function in skeletal muscle (Russell et al., 2014).

Mitochondria retain a unique conserved genome from prokaryotic ancestors, referred to as the mitogenome or mitochondrial DNA (mtDNA; Anderson et al., 1981; Groennebaek & Vissing, 2017). The human mitogenome is organized as a circular, covalently closed, double-stranded DNA molecule made of 16,569 bp (Anderson et al., 1981; Andrews et al., 1999). It contains 37 genes, of which 13 encode subunits of enzyme complexes of the oxidative phosphorylation system, 22 encode transfer RNAs (tRNAs) and two encode ribosomal RNAs (rRNAs; Mercer et al., 2011; Taanman, 1999). However, ~1500 polypeptides are located in the mitochondria, and the vast majority

© 2018 The Authors. Experimental Physiology © 2018 The Physiological Society

• What is the topic of this review?

This article draws evidence from the current literature to support the hypothesis that non-coding RNA-mediated gene regulation takes place in the mitochondria. An emphasis is put on skeletal muscle.

• What advances does it highlight?

This review highlights the potential role of micro-RNAs and long non-coding RNAs in mitochondrial gene regulation. The discovery of a new level of skeletal muscle mitochondria-controlled gene regulation presents exciting perspectives for our understanding of skeletal muscle physiology in health and disease.

of them, including 85% of the respiratory complex subunits, are encoded in the nuclear genome (Mercer et al., 2011; Taanman, 1999). Mitochondrial regulation therefore relies on the complex coordination of mitochondrial- and nuclear-encoded transcripts.

Non-coding RNAs (ncRNAs) are RNA molecules that are not translated into a protein. The ncRNAs include the long-known tRNAs and rRNAs, which are key to production of functional proteins in the cytoplasm and the mitochondria (Hombach & Kretz, 2016). In addition, several classes of small [<200 nucleotides (nt)], regulatory ncRNAs have been described during recent decades, including small interfering RNAs (siRNAs), piwi-associated RNAs (piRNAs), microRNAs (miRNAs), small nucleolar RNAs (snoRNAs) and small nuclear RNAs (snRNAs). These short RNA species play a role in the modulation of gene expression (C. Zhang, 2009) or in the processing of other RNA species (Matera, Terns, & Terns, 2007). In contrast, long noncoding RNAs (IncRNAs; > 200 nt) can have multiple functions in the cell, including the promotion or repression of gene transcription (Djebali et al., 2012). The tRNA and rRNA are encoded by both the nuclear and the mitogenome, but the idea of subcellular localization of other ncRNAs, particularly within mammalian tissue, is a more recent concept. In 2006, an early study generated complementary DNA libraries from size-selected murine mitochondrial RNA and cloned a series of putative mitochondrial-encoded small ncRNA species. Presumably owing to low expression levels, their presence could not be confirmed by Northern blot analysis (Lung et al., 2006). RNA sequencing performed on mouse testis, mouse small intestine and human small intestine tissue later confirmed that the human and murine mitogenomes potentially encode thousands of small ncRNAs (Ro et al., 2013). In addition, we now know that IncRNAs can also be encoded by the mitochondrial genome (Gao et al., 2018; Landerer et al., 2011; Rackham et al., 2011) and selectively localize in different cell compartments, including the nucleus, the cytoplasm (Djebali et al., 2012) and the mitochondria.

The localization of regulatory ncRNA species within the mitochondria presents exciting possibilities for mitochondrial gene regulation and opens new avenues to expand our understanding of



FIGURE 1 Possible origin and function of mitochondrial microRNAs (miRNAs) and long non-coding RNAs (IncRNAs). Dashed lines depict relationships that have been hypothesized. Green and red arrows indicate 'enhancement' and 'inhibition', respectively. miRNA, micro-RNA; pri-miRNA, primary microRNA; pre-miRNA, pre microRNA; mt, mitochondrial; IncRNA, long non-coding RNA; PNPase, Polynucleotide Phosphorylase; AGO2, Argonaute2; VDAC, voltage-dependent anion channel; p-bodies, processing-bodies.

human muscle physiology (Figure 1). This review draws evidence from the recent literature to discuss the hypothesis that a level of ncRNA-mediated gene regulation takes place in the mitochondria, with a unique focus on the possible implications for skeletal muscle physiology.

2 | microRNAs IN THE MITOCHONDRIA

Of particular interest is the widely studied class of miRNAs that modulate gene expression (Guo, Ingolia, Weissman, & Bartel, 2010; Humphreys, Westman, Martin, & Preiss, 2005; Olsen & Ambros, 1999). MicroRNA-mediated control of gene expression was first suggested in 1993 (R. C. Lee, Feinbaum, & Ambros, 1993; Reinhart et al., 2000) and has added an extra level of complexity to our understanding of transcriptional and translational regulation (Bartel, 2004). Micro-RNAs are 20-22 nt single-stranded RNA molecules (R. C. Lee & Ambros, 2001; R. C. Lee et al., 1993) that act by binding to specific target mRNA molecules. Their primary role is to silence protein expression, which can be achieved by either of two post-transcriptional mechanisms; degrading target mRNA molecules or preventing the translation of the corresponding protein (Guo et al., 2010; Humphreys et al., 2005; Olsen & Ambros, 1999). In some cases, miRNAs can also upregulate protein translation by stabilizing their target mRNA molecule (Vasudevan, Tong, & Steitz, 2007; X. Zhang et al., 2014). The principles underlying recognition of miRNAs/mRNAs are not fully understood. It was first established that miRNAs bind to specific sequences of the 3' untranslated region (3'UTR) or the coding region of their target transcript (Pasquinelli, 2012) thanks to a 6-8 nt conserved seed sequence localized at their 5' end. This mode of binding can involve perfect or imperfect Watson-Crick recognition (Brennecke, Stark, Russell, & Cohen, 2005) and was long considered the only

1133

one. Experimental evidence, however, demonstrates that ~60% of binding of miRNAs/mRNAs cannot be explained by canonical rules (Chi, Hannon, & Darnell, 2012; Helwak, Kudla, Dudnakova, & Tollervey, 2013), where non-seed motifs or mismatching seed-like motifs are also involved in functional binding. It follows that each miRNA has the potential to target multiple mRNAs through canonical and non-canonical interactions. Approximately 60% of protein-coding genes are regulated by miRNAs (Friedman, Farh, Burge, & Bartel, 2009).

The miRBase database v. 22, a searchable database of published miRNA sequences (Griffiths-Jones, Saini, van Dongen, & Enright, 2008; Kozomara & Griffiths-Jones, 2014), retrieves > 4000 entries corresponding to annotated human miRNAs. MicroRNAs are encoded in both coding and non-coding parts of the nuclear genome (Bartel, 2004). They are initially transcribed into a primary miRNA molecule (pri-miR; Bartel, 2004), which is cleaved by the RNase III-type endonuclease Drosha-associated-with-Pasha to become a pre-miRNA molecule (pre-miR; Y. Lee et al., 2003). The pre-miR molecule is then exported from the nucleus to the cytoplasm by Exportin-5 (Lund, Güttinger, Calado, Dahlberg, & Kutay, 2004), where the endoribonuclease Dicer-1 splices it into a single-stranded mature miRNA molecule (Y. Lee et al., 2003). In the cytoplasm, the mature miRNA strand associates with the Argonaute (AGO) proteins to form RNA-induced silencing complexes (RISCs; Bartel, 2004) and direct them to downregulate gene expression. In mammals, the AGO2 protein is necessary to catalyse the cleavage of the target transcript (Chendrimada et al., 2005; Liu et al., 2004; Meister et al., 2004).

MicroRNAs typically regulate protein expression within the cytoplasm (Bandiera, Mategot, Girard, Demongeot, & Henrion-Caude, 2013). Extracytoplasmic miRNA localization was first demonstrated in the nucleolus of rat skeletal muscle cells using locked nucleic acid (LNA) probes targeting the mature form of the muscle-enriched miR-206 (Politz, Zhang, & Pederson, 2006). Since then, evidence suggesting further subcellular localization of miRNAs has emerged (C. Zhang, 2009), including in the nucleus (Salmanidis, Pillman, Goodall, & Bracken, 2014), processing bodies (Gibbings, Ciaudo, Erhardt, & Voinnet, 2009) and the mitochondria (Barrey et al., 2011; Lung et al., 2006), where miRNAs could regulate the expression of mitochondrial transcripts and subsequent protein translation (Bassi, Heads, Marber, & Clark, 2008; X. Zhang et al., 2014). Complementary DNA libraries generated from size-selected murine liver and kidney mitochondrial RNA (Lung et al., 2006) identified four clones corresponding to nuclear-encoded miRNAs, let-7f-1, let-7 g, miR-122a,b and miR-101b. Further observations confirmed the existence of a range of nuclearencoded miRNAs in highly purified mitochondria isolated from rat liver (Kren et al., 2009) and HeLa cells (Bandiera et al., 2011). It was proposed that the mitochondria might act as a reservoir of miRNAs for the cell, because none of the discovered miRNAs was predicted to target the mitogenome or nuclear transcripts encoding mitochondrial proteins (Kren et al., 2009). A later study, however, revealed that other mitochondrial miRNAs had several putative nuclear and mitochondrial-encoded targets (Bandiera et al., 2011), suggesting that the newly named 'mitomiRs' might reflect another level of crosstalk in the processes underlying mitochondrial regulation.

Current knowledge suggests that all miRNAs are encoded by the nuclear genome (Geiger & Dalgaard, 2017; Ha & Kim, 2014). However, the mapping of a series of pre-miRNA and mature miRNA sequences on the human mitogenome (Bandiera et al., 2011; Barrey et al., 2011; Shinde & Bhadra, 2015) suggests that some miRNAs might originate from the mitochondrial DNA. Whether these miRNA sequences are transcribed from the mitochondrial genome remains speculative.

3 | OVERVIEW OF microRNA EXPRESSION WITHIN RODENT AND HUMAN MITOCHONDRIA

More than 100 miRNA species have been identified in mitochondria isolated from rodent tissue, such as rat liver (Kren et al., 2009), rat hippocampus (W. X. Wang et al., 2015), rat heart (Das et al., 2012), mouse liver (Bian et al., 2010) and mouse heart (Bian et al., 2010; Jagannathan et al., 2015; X. Zhang et al., 2014). At least 20 miRNA species were observed to be more highly expressed in the mouse liver mitochondria than in whole-liver extracts, with several highly expressed whole-liver miRNAs having low expression in the mitochondria (Bian et al., 2010). The enrichment of miRNAs in mitochondria might therefore be independent of their total cellular abundance.

To date, although only a few studies have investigated miRNA expression within human mitochondria, they have identified at least 400 known and 200 purportedly new miRNAs in mitochondria from muscle, kidney and cancer cells (Bandiera et al., 2011; Barrey et al., 2011; Dasgupta et al., 2015; Sripada et al., 2012). Initially, Barrey et al. (2011)) used in situ hybridization and isolation of the mitochondria from human primary cultured myotubes to quantify the expression of 204 miRNAs localized within mitochondria via RTqPCR. Around the same time, 13 nuclear-encoded miRNA species enriched in mitochondria from human cervical cancer (HeLa) cells were observed (Bandiera et al., 2011). Additionally, deep sequencing identified an association of 327 known and 13 putative new miRNAs to mitochondria in HeLa cells, in addition to 428 known and 196 putative new miRNAs to mitochondria in human embryonic kidney (HEK293) cells (Sripada et al., 2012). Other studies have also found miRNA species in isolated human mitochondria from 143B (osteosarcoma) and 206 p0 (derived from 143B) cell lines (Dasgupta et al., 2015; Mercer et al., 2011). Collectively, these studies have established that there are a large number of miRNA species expressed in mitochondria of multiple human cell types.

4 | MITO-myomiRs: POTENTIAL ROLES OF miRNAs IN SKELETAL MUSCLE MITOCHONDRIA

Skeletal muscle makes up 40% of human body mass (Rasmussen & Phillips, 2003). As such, skeletal muscle accounts for 20–30% of whole-body energy expenditure at rest (Zurlo et al., 1990), for up to

1135

90% during exercise (Zurlo et al., 1990) and for up to 50% of mealderived glucose disposal (Radziuk & Inculet, 1983), making it one of the tissues displaying the highest mitochondrial abundance (Fernández-Vizarra et al., 2011). A series of life-threatening and economy-draining conditions, including myopathy, neuromuscular, neurodegenerative and metabolic diseases, are positively associated with impairments in skeletal muscle mitochondrial function (Russell et al., 2014). Expanding our understanding of mitochondrial regulation in muscle health and disease conditions, including regulation by ncRNAs, therefore has the potential to contribute to the identification of new therapeutic targets.

MicroRNAs regulate skeletal muscle mass, function and metabolism (reviewed by Zacharewicz, Lamon, & Russell, 2013). Of particular importance, skeletal muscle displays a series of muscle-enriched miRNAs, termed myomiRs. Canonical myomiRs comprise the miR-1, miR-133 and miR-206 families of miRNAs and originate from three cistrons: (miR-1-2, miR-133a-1), (miR-1-1, miR-133a-2) and (miR-133b, miR-206) (McCarthy, 2011). With time, the myomiR list has expanded to include miR-208a, miR-208b, miR-499 and miR-486 (reviewed by Kirby & McCarthy, 2013). MyomiRs undergo high levels of regulation in skeletal muscle development and adaptation in health and disease (Lamon et al., 2017; Zacharewicz et al., 2013). However, muscle-specific deletions of individual myomiRs lead to only minor phenotypic effects in mouse models (Kirby & McCarthy, 2013), suggesting possible redundancy and fostering further research in this domain. In addition, and without any formal classification, we and others have identified other muscle-enriched miRNAs, including the miR-23 (Russell et al., 2013; Russell et al., 2013), miR-29 (Greco et al., 2009; Russell et al., 2013; Russell et al., 2013; H. Wang et al., 2008), miR-31 (Greco et al., 2009; Russell et al., 2013; Russell et al., 2013) and miR-181 (Naguibneva et al., 2006; Russell et al., 2013; Russell et al., 2013) families of miRNAs. The ubiquitously expressed let-7 family of miRNAs plays an antiproliferative role (Johnson et al., 2007) and can be added to the picture of the miRNAs that are highly expressed in skeletal muscle.

Exercise and disease alter skeletal muscle miRNA expression. For example, in humans, expression levels of the muscle-enriched miR-1, miR-133a, miR-133b and miR-181a are increased (Nielsen et al., 2010; Russell, Lamon et al., 2013), whereas miR-9, miR-23a, miR-23b and miR-31 are decreased (Russell, Lamon et al., 2013), after a single bout of submaximal endurance exercise. Furthermore, the type of exercise can have a direct impact on the nature of the miRNA response (Drummond, McCarthy, Fry, Esser, & Rasmussen, 2008; Ringholm et al., 2011). When compared with healthy muscle, the same miRNAs are dysegulated in several human myopathies and dystrophies (Eisenberg, Alexander, & Kunkel, 2009; Gambardella et al., 2010; Greco et al., 2009), with unloading (Ringholm et al., 2011), and in neurodegenerative diseases that are specifically associated with impaired mitochondrial function, such as amyotrophic lateral sclerosis (Russell et al., 2018; Russell et al., 2013; Williams et al., 2009).

All known myomiRs are encoded in the nucleus, and their preand mature sequences do not map the human mitogenome (Barrey et al., 2011). Barrey et al. (2011)) first demonstrated the presence of mature and immature miRNA forms in the mitochondria isolated from human primary myotubes, and RT-qPCR analyses established the presence of multiple miRNA species, amongst which miR-133a, miR-133b, miR-23a and the ubiquitously expressed let-7 family were within the most abundant. Less abundant but still significantly detected in the mitochondria were the myomiRs miR-206 and miR-1. The presence of these miRNAs in muscle mitochondria prompted the question of their potential gene targets. *In silico* analysis identified putative, canonical target sites on several mitochondrial transcripts for miR-133a and members of the let-7 family (Barrey et al., 2011), suggesting that miRNA-mediated regulation of gene expression might occur within the muscle mitochondria.

Although few studies have identified a functional role for mitochondrial miRNAs, many authors have speculated on their potential gene and protein targets using target prediction tools. These predictions should, however, be considered with extreme caution. Firstly, target prediction algorithms do not predict the majority of miRNA/mRNA interactions (Chi et al., 2012; Helwak et al., 2013). Secondly, even when physical binding is confirmed experimentally, for example by means of a reporter assay, we have frequently observed an absence of regulation of the corresponding gene and/or protein targets in vitro, in vivo or both (Pinto et al., 2017; E. Zacharewicz and S. Lamon, unpublished observations), presumably owning to specific thermodynamic and steric binding conditions. Overall, this indicates that our very partial understanding of miRNA/mRNA binding rules constitutes a major limitation of target predictions. Therefore, if not accompanied by functional validation. in silico analysis of the biological pathways that may or may not be influenced by specific miRNAs should be treated as purely speculative. Emerging experimental approaches will probably help to overcome these limitations in the coming years. For example, cross-linked immunoprecipitation (CLIP) is a recent technique that identifies the direct RNA binding sites of RNA-binding proteins (Licatalosi et al., 2008). When applied to AGO2, it can be used to identify miRNA binding sites and validate their physical association to RISCs in a physiological environment. AGO2 CLIP can be coupled to high-throughput sequencing of RNA (HITS-CLIP), to high-throughput sequencing of RNA isolated from photoactivable ribonucleoside (PAR-CLIP) and to individual nucleotide resolution (iCLIP) (Bottini et al., 2017), providing a comprehensive map of AGO2 binding targets in specific biological conditions.

In a very elegant study, X. Zhang et al. (2014) confirmed that miR-1 was localized in the mitochondria of C_2C_{12} myocytes and that the quantity of miR-1 increased with myotube differentiation. More importantly, the functional complex miR-1/AGO2 bound to the mitochondrial transcripts Cox1, Nd1, Cytb, Cox3 and Atp8 and enhanced their translation, providing the first available evidence for miRNA-mediated mitochondrial gene regulation in skeletal muscle cells (X. Zhang et al., 2014). Reports of miRNA-mediated activation of translation rather than repression are rare but not without precedent (Vasudevan et al., 2007). Although the underlying mechanisms can only be speculative, the prokaryotic origin of mitochondria may offer a possible explanation. Firstly, the AGO2 complex plays different roles in specific biological contexts, and prokaryotic Argonaute proteins possess unique mechanistic features (Hutvagner & Simard, 2008; Shabalina & Koonin, 2008). Secondly, in bacteria, small regulatory RNAs can activate a target mRNA by direct base pairing. A variety

of mechanisms, including the 'anti-antisense mechanisms' and the 3'-processing-mediated transcript stabilization, act respectively to alleviate the inhibition on a ribosome binding site or to increase the stability of a target transcript (Fröhlich & Vogel, 2009). Finally, mitochondrial-encoded mRNAs present polyadenylation patterns inherited from their bacterial ancestors that are different from those found on nuclear-encoded mRNAs (Slomovic, Laufer, Geiger, & Schuster, 2005), including internal polyadenylation. As such, they may present a possible target for positive regulatory mechanisms mediated by miRNAs in the mitochondria (Nouws & Shadel, 2014; Vasudevan et al., 2007).

Interestingly, miR-1 can simultaneously repress the expression of cytoplasmic gene targets and promote the expression of mitochondrial gene targets (X. Zhang et al., 2014). As such, mitomiRs might coordinate the concurrent translation of nuclear- and mitochondrialencoded mitochondrial proteins (Nouws & Shadel, 2014); for example, to control the formation of the respiratory complex. This is especially relevant to skeletal muscle, a tissue that requires constant and rapid adaptation to environmental conditions and changing energy demands. In this regard, the idea of partitioning certain miRNAs between the cytoplasm and the mitochondria to enhance the efficiency of mitochondrial activities in the muscle warrants further investigation. Another example comes from the observation of a functional form of a muscle-enriched miRNA, miR-181, in the mitochondria of cardiac muscle cells (Das et al., 2012). Similar to skeletal muscle, cardiac muscle metabolism primarily relies on substrate oxidation, and cardiac muscle has a high mitochondrial content (Fernández-Vizarra et al., 2011). MiR-181c co-immunoprecipitated with AGO2 complexes extracted from neonatal rat ventricular myocytes. Not only did miR-181c bind to the mitochondrial transcript Cox1 but also it significantly decreased COX1 protein expression, demonstrating that miR-181c directly regulates COX1 translation in the mitochondria (Das et al., 2012). In addition, overexpressing miR-181c in rat cardiac myocytes impaired mitochondrial function by dysregulating the formation of complex IV. Overall, these results suggest that, in cardiac muscle, miR-181c might be partitioned in different cellular compartments to fine-tune the formation of the respiratory complexes.

Some miRNA species may also regulate mitochondrial metabolism and function by acting at multiple subcellular locations. A series of nuclear-encoded miRNAs located in the cytoplasm specifically modulates mitochondrial metabolism and function in skeletal muscle [reviewed by our group (Russell et al., 2014; Tsitkanou, Della Gatta, & Russell, 2016) and others (Lima et al., 2017; Massart, Katayama, & Krook, 2016)]. Of particular interest are miR-494 (Yamamoto et al., 2012), miR-23a (Russell et al., 2013; Wada et al., 2015), miR-696 (Aoi et al., 2010) and miR-542 (Garros et al., 2017). These miRNAs indirectly or directly target key, nuclear-encoded regulators of the mitochondrial function, including mitochondrial transcription factor A (Tfam; Yamamoto et al., 2012), peroxisome proliferatoractivated receptor gamma coactivator 1-alpha (Pgc-1 α ; Aoi et al., 2010; Russell et al., 2013), cytochrome b and cytochrome c oxidase complex IV (Cox4; Wada et al., 2015) and mitochondrial ribosomal protein MRPS10 (Garros et al., 2017). Of these miRNAs, miR-23a and

miR-542 are also expressed in human myotube mitochondria (Barrey et al., 2011), although no putative canonical target site for miR-23a or miR-542 was found in any mitochondrial transcript (Barrey et al., 2011). It is however, premature to dismiss a regulatory role for these miRNAs within the mitochondria. Although the role of miR-23a in the mitochondria is currently unknown, a study demonstrated that miR-23a co-immunoprecipitated with mitochondrial AGO2 complexes isolated from rat brain tissue (W. X. Wang et al., 2015). A direct association between miR-23a and AGO2 might therefore be indicative of functional RISCs within the mitochondria. Bearing in mind the role of miR-23a in the regulation of mitochondrial metabolism in the cytoplasm, the presence of a functional form of miR-23a in the mitochondria warrants further investigation into its potential gene targets and biological effects. A list of miRNAs that are present in muscle mitochondria and have known regulatory functions pertaining to mitochondrial homeostasis is presented in Table 1.

5 | IMMATURE FORMS OF miRNA AND miRNA-ASSOCIATED PROTEINS ARE EXPRESSED WITHIN THE MITOCHONDRIA

Emerging evidence suggests that some level of miRNA biogenesis and processing might occur within the mitochondria themselves. This is based on findings that immature forms of miRNA (Barrey et al., 2011: Shinde & Bhadra, 2015), in addition to members of the miRNA biogenesis machinery (W. X. Wang et al., 2015) and components of the RISC, localize in the mitochondria (Das et al., 2012; Dasgupta et al., 2015; Jagannathan et al., 2015; W. X. Wang et al., 2015). Two pre-miR sequences that aligned with the human mitogenome, pre-mir302a and pre-let-7b, were detected within the perinuclear mitochondria of human myoblasts using in situ hybridization (Barrey et al., 2011). Further investigation found the enrichment of the mature form of miR-let-7b, but not miR-302a, in the mitochondrial fraction. This suggests that the biogenesis of pre-miRNAs might occur within the mitochondria. Likewise, pre-miR-338 was observed in the mitochondrial fraction of axons of rat superior cervical ganglion neurons (Vargas et al., 2016). The authors proposed that a pool of pre-miR-338 in the mitochondria might act as a reservoir of miRNAs to regulate the translation of nuclear-encoded mitochondrial mRNAs. However, further studies are required to examine the extent to which pre-miRs also undergo processing within the mitochondria. Potential transport mechanisms of mature and immature miRNAs into the mitochondria are discussed in other reviews (Bandiera et al., 2013; Borralho, Rodrigues, & Steer, 2015; Duarte, Palmeira, & Rolo, 2014; Macgregor-Das & Das, 2018; Sripada, Tomar & Singh, 2012) and suggest a role for AGO2 (Maniataki & Mourelatos, 2005), polynucleotide phosphorylase (PNPASE; Shepherd et al., 2017; G. Wang et al., 2010), P-bodies (Huang et al., 2011) and voltage-gated ion channels (Salinas et al., 2006).

The presence of AGO2 and dicer in the mitochondria, along with immature forms of miRNAs, reinforces the idea that some level of miRNA-mediated post-transcriptional regulation occurs in **TABLE 1** Skeletal muscle-related mitochondrial microRNAs

	MicroRNA identity	Observed in muscle mitochondria	Experimental model	Gene targets	Reference
Canonical myomiRs					
	miR-1	Barrey et al. (2011); R. Jagannathan et al. (2015); X. Zhang et al. (2014)	C_2C_{12} myoblasts	mtCox1, mtNd1, mtCytb, mtCox3, mtAtp8 (transcriptional activation)	X. Zhang et al. (2014)
	miR-133a	Barrey et al. (2011); R. Jagannathan et al. (2015)	C57BI/6 mice; miR-133a ^{-/-} KO	$Pgc-1\alpha$, $Pgc-1\beta$, $Tfam$ (transcriptional repression)	Nie et al. (2016)
	miR-206	Barrey et al. (2011)	C57Bl/6 mice; miR-206 ^{-/-} KO COS1 cells	HDAC4 (translational repression) Hdac4 (3'UTR, transcriptional repression)	Williams et al. (2009)
Muscle-enriched miRNAs					
	miR-23a	Barrey et al. (2011)	Human skeletal muscle HeLa cells miR-23a Tg mice C57Bl/6 mice	PGC-1 α (transcriptional/translational repression) Pgc-1 α (3'UTR, transcriptional repression) PGC-1 α , COXIV, Cyt-c (translational repression) PGC-1 α (transcriptional/translational repression), Cyt-c (transcriptional repression)	Russell, Wada, et al. (2013) Wada et al. (2015) Safdar et al. (2009)
	miR-31	Barrey et al. (2011)	Human skeletal muscle C_2C_{12} myoblasts	HDAC4, NRF1 (transcriptional/translational repression) <i>Hdac4</i> , <i>Nrf</i> 1 (3'UTR, transcriptional repression)	Russell, Lamon, et al. (2013)
Cardiac miRNAs					
	miR-181c	Barrey et al. (2011); Das et al. (2012)	Neonatal rat primary cardiomyocytes	mtCOX1 (translational repression), mtCOX2 (transcriptional/translational repression), mtCox1 (3'UTR, transcriptional repression)	Das et al. (2012)
	miR-378	Barrey et al. (2011); R. Jagannathan et al. (2015)	HL-1 HEK293	mtATP6 (translational repression) mtATP6 (3'UTR, transcriptional repression)	R. Jagannathan et al. (2015)

Muscle-related mitochondrial miRNAs were classified as canonical myomiRs, skeletal muscle-enriched miRNAs and cardiac muscle-enriched miRNAs. Their validated gene targets are indicated, in addition to the level of miRNA-mediated regulation. miRNA, MicroRNA; Tg, transgenic; UTR, untranslated region.

the mitochondria. An early seminal study used Western blotting to identify the AGO2 protein in purified mouse liver mitochondria (Bian et al., 2010). This observation was confirmed in rat brain tissue, where co-immunoprecipitation of the AGO protein complexes followed by Western blotting revealed the presence of AGO2 in both the mitochondrial matrix and the outer mitochondrial membrane (W. X. Wang et al., 2015). Studies confirmed the presence of AGO2 in human mitochondria. AGO2 protein expression was detected in a purified mitochondria fraction of HeLa cells using Western blotting, immunofluorescence confocal microscopy and co-immunoprecipitation with the mitochondrial transcript COX3 (Bandiera et al., 2011), and in a purified mitochondrial fraction of 143B human bone osteosarcoma-derived cells using Western blotting (Dasgupta et al., 2015). With respect to skeletal muscle, tandem massspectrometry analysis confirmed the presence of AGO2, but not AGO1 and AGO3, in highly purified mitochondrial and mitoplast fractions extracted from C_2C_{12} cells. The number of AGO2 molecules present in the mitochondria increased with C_2C_{12} cell differentiation, suggesting that more AGO2 might be imported into the mitochondria upon myotube differentiation (X. Zhang et al., 2014). These observations were extended to the DICER protein, which was detected in the mitochondrial matrix and outer mitochondrial membrane of rat brain by Western blotting (W. X. Wang et al., 2015). Using the same method, DICER was observed in the mitochondria of different fractions of axons of rat superior cervical ganglion neurons (Vargas et al., 2016), a result confirmed by co-immunostaining with the mitochondrial marker VDAC1.

The presence of AGO2 and DICER in the mitochondria suggests that immature forms of miRNAs might originate from the mitogenome. However, AGO2 and DICER are versatile proteins that are involved in several biological processes that can be miRNA dependent or independent (Janowski et al., 2006; R. C. Lee et al., 1993). AGO2 localizes in different cellular compartments, including the nucleus (Ohrt et al., 2008), the Golgi apparatus (Kim, Maizel, & Chen, 2014), the endoplasmic reticulum and processing bodies (Leung, Calabrese, & Sharp, 2006). Depending on its localization, AGO2 regulates different transcriptional and post-transcriptional processes (Janowski et al., 2006). Likewise, DICER is an endoribonuclease that

1137

cleaves both double-stranded RNA and pre-miRs into mature siRNAs (Carthew & Sontheimer, 2009; Tomari & Zamore, 2005) and miRNAs (Bernstein, Caudy, Hammond, & Hannon, 2001), respectively (R. C. Lee et al., 1993). A few studies started to provide insights into whether DICER and AGO2, respectively, process pre-miRs and associate with mature miRNAs in the mitochondria, X. Zhang et al. (2014) provided several lines of evidence not only that AGO2 was present in the mitochondria, but also that the miRNA/AGO2 complex interacted with mitochondrial transcripts to enhance mitochondrial translation. The underlying mechanisms, however, remain speculative, partly because GW182, the functional binding partner of AGO2, was not present in the mitochondria (X. Zhang et al., 2014). Interestingly, an earlier study by Ro et al. (2013) showed that DICER inactivation impaired, but did not suppress, the production of mitochondrial-encoded small ncRNAs in somatic testis cells. The authors made the intriguing suggestion that mitochondrial small ncRNAs might be the product of other, currently undescribed, mitochondrial ribonucleases. These early studies demonstrate that a functional form of AGO2 is present in the mitochondria and suggest that DICER might partly process premiRNAs in the mitochondria.

6 | LONG NON-CODING RNAS IN THE MITOCHONDRIA

Long non-coding RNAs are a class of ncRNA that has received less attention than their shorter counterparts, maybe owing to the diversity and complexity of their roles in the cell. The IncRNAs can be arbitrarily defined as all ncRNAs > 200 nt in length. They are usually shorter than 10,000 nucleotides and can be classified by abundance ('high' versus 'low') or by function (comprehensively reviewed by Djebali et al., 2012; X. Wang, Song, Glass, & Rosenfeld, 2011). A conservative estimate is that the human nuclear genome might encode ~17,000 IncRNAs (X. Wang et al., 2011). Long ncRNAs can be single or double stranded, and although the function of a doublestranded RNA molecule is unclear, authors have suggested that these molecules might act as regulators of their single-stranded equivalents (Rackham et al., 2011; X. Wang et al., 2011). These molecules may or may not be polyadenylated, determining their stability (Wilusz et al., 2012). Long ncRNAs regulate gene and protein expression in the cell through a variety of molecular processes (Djebali et al., 2012), including promoting or repressing gene transcription, providing a scaffold for regulatory proteins, facilitating histone methylation and regulating protein activities through allosteric modifications. In addition, IncRNAs are also are involved in the regulation of other ncRNA species (Cesana et al., 2011; Djebali et al., 2012; Jarroux, Morillon, & Pinskaya, 2017; Kung et al., 2013; Ponting, Oliver, & Reik, 2009). Human IncRNAs are mostly transcribed by RNA polymerase II, with some rare IncRNAs being dependent on RNA polymerase III (Jarroux et al., 2017). As miRNAs, IncRNAs regulate mitochondrial function and are dysregulated in pathological states. For example, the IncRNA Tug1 binds to and positively regulates Pgc-1 α , increasing mitochondrial content and respiration (Long et al.,

2016), whereas aberrant *Tug1* expression reduces $Pgc-1\alpha$ transcripts and contributes to the mitochondrial dysregulation characteristic of diabetic nephropathy in mouse kidney cells.

Long non-coding RNAs are also encoded by the mitochondrial genome (Gao et al., 2018; Landerer et al., 2011; Rackham et al., 2011), but our knowledge and understanding of their putative role in the mitochondria is still in its infancy. In 2011, a study used deepsequencing, Northern blotting and strand-specific gRT-PCR in the mitochondria from HeLa cells to identify three IncRNAs generated from the mitogenome: IncND5, IncND6 and IncCytb RNA (Rackham et al., 2011). The knockdown of five nuclear-encoded proteins involved in RNA processing independently decreased the abundance of these IncRNAs, suggesting a role for these proteins in the regulation of mitochondrial ncRNA expression (Rackham et al., 2011). Of particular interest, all three lncRNAs were found in relatively high abundance in skeletal muscle mitochondria, when compared with other tissues (Rackham et al., 2011). Whether these IncRNAs act in the mitochondria themselves or are exported from the mitochondria to other cellular compartments (Cesana et al., 2011; Landerer et al., 2011) using and/or contributing to the mitochondrial-nuclear communication pathway (Butow & Avadhani, 2004), and what their roles are, is still to be determined. Recently, two new IncRNAs encoded by the human mitogenome, MDL1 and MDL1AS, were discovered in the MCF-7 breast cancer cell line (Gao et al., 2018). These IncRNAs might be involved in positive and negative feedback systems that increase or stabilize the expression of mitochondrial transcripts (Gao et al., 2018). A set of chimeric IncRNAs also exists in mouse and human mitochondria (Dong, Yoshitomi, Hu, & Cui, 2017), but with no evidence of expression in muscle tissue in either case. Future research on IncRNAs and their regulatory role in the mitochondria will help to expand our understanding of the regulation of skeletal muscle physiology and of mitochondrial biology in general.

7 | PERSPECTIVE AND CONCLUSION

The efficient regulation of mitochondrial metabolism is essential to whole-cell homeostasis. The ncRNAs, specifically miRNAs, play an increasingly important role in the regulation of gene expression across a variety of tissues. Emerging evidence highlights the subcellular localization of miRNAs within the mitochondria of the brain, heart, liver and, more recently, skeletal muscle. The idea of miRNA-mediated regulation of the mitochondrial genome therefore presents exciting possibilities for future research.

Initial forays into mitochondrial miRNAs suggested that the mitochondrion was simply a reservoir for cytosolic miRNAs (Duarte, Palmeira, & Rolo, 2015). However, direct observations of interactions between mitochondrial miRNAs and mitochondrial transcripts indicate that miRNAs have more advanced roles in influencing mitochondrial metabolism and morphology. The presence of premiRNAs within human myoblast mitochondria (Barrey et al., 2011; Shinde & Bhadra, 2015) suggests that the mitochondrial genome might also harbour sequences for a number of miRNAs. The prospect of extranuclear, non-canonical miRNA biogenesis pathways is an intriguing avenue for future investigations.

The bulk of current knowledge about miRNA profiles in pathological, stress and exercise conditions stems from wholecell investigations. The response of miRNAs at the subcellular level is, however, largely unknown and is limited to a handful of studies. Differential miRNA expression exists in spatially distinct subsarcolemmal and intermyofibrillar mitochondrial subpopulations within rodent cardiac muscle (Jagannathan et al., 2015). Although expression of miR-378 in whole cardiac muscle was unchanged after diabetic insult, miR-378 expression increased in intermyofibrillar mitochondria, suggesting that miRNAs might be redistributed between mitochondrial subpopulations in pathological states (Jagannathan et al., 2015). Of functional significance, miR-378 was also found to bind to mitochondrial-encoded Atp6, along with an associated reduction in ATP6 protein abundance and enzyme activity in intermyofibrillar mitochondria (Jagannathan et al., 2015). These findings suggest that, independent of wholecell miRNA populations, mitochondrial miRNAs might have advanced regulatory roles in both mitochondrial and whole-cell metabolism.

As yet, the extent of miRNA-mediated regulation of mitochondrial metabolism in healthy skeletal muscle is an underdeveloped field. However, it is reasonable to speculate that miRNAs within the mitochondria might play a role in the increased mitochondrial biogenesis observed after exercise. Firstly, observational studies of whole muscle report the differential expression of miRNAs after acute exercise and exercise training (Nielsen et al., 2010; Russell et al., 2013; Zacharewicz et al., 2014). For example, miR-1 and miR-133a expression increased, whereas miR-206 expression remained unchanged, in whole skeletal muscle after an intense bout of endurance exercise (Nielsen et al., 2010; Russell et al., 2013). MiR-1/AGO2 binds to the mitochondrial transcripts Cox1, Nd1, Cytb, Cox3 and Atp8 in murine skeletal muscle cells (X. Zhang et al., 2014), and further in sillico analysis suggests that miR-133a and miR-206 might have more putative binding sites within the mitochondrial genome (Das et al., 2012). Secondly, PGC-1a, NRF-1 and TFAM mRNA, all components of the mitochondrial biogenesis pathway, are under miRNA-mediated control in human myoblasts (Barrey et al., 2011) and rodent skeletal muscle cells (Nie et al., 2016; Safdar, Abadi, Akhtar, Hettinga, & Tarnopolsky, 2009). Thirdly, some miRNAs that respond to exercise are found in the mitochondria and play a role in mitochondrial regulation. Examples from our own research are miR-23a and miR-31. MiR-23a is reduced in human skeletal muscle after a single bout of endurance exercise (Russell et al., 2013) and binds to Pgc-1α 3'UTR in vitro (Russell et al., 2013). In addition, overexpressing miR-23a in mice downregulates the skeletal muscle protein abundance of PGC-1 α , cytochrome *b* and cytochrome *c* oxidase subunit IV (Russell et al., 2013). Likewise, miR-31 is reduced in human skeletal muscle after a single bout of endurance exercise and targets HDAC4 and NRF1, genes regulating muscle growth and mitochondrial biogenesis, respectively (Russell et al., 2013). Finally, the upregulation in mitochondrial biogenesis after exercise is influenced by factors at the subcellular level. For example, the subcellular localization of

PGC-1 α , albeit between the cytosol and nucleus, is essential to the regulation of mitochondrial biogenesis (Wright et al., 2007). For these reasons, studies investigating how miRNAs respond to endurance exercise within skeletal muscle mitochondria are required to examine how miRNAs mediate adaptive responses in healthy skeletal muscle. The miRNA-mediated regulation of the mitochondrial genome will provide greater depth to our understanding of the adaptive responses targeting mitochondrial metabolism after endurance exercise.

In conclusion, the regulatory role of ncRNAs, particularly miRNAs, in mitochondrial metabolism suggests that their expression might be altered within the mitochondria in pathological and physiological states. Although research in this area is currently in its infancy, preliminary studies observe interactions between muscle-enriched miRNAs and mitochondrial mRNAs that might directly influence mitochondrial metabolism within skeletal and cardiac muscle tissue in vitro. As yet, the physiological significance of miRNAs within human skeletal muscle mitochondria can only be speculative. Coordination between subcellular compartments, for example between the nucleus, cytosol and mitochondria, is essential for efficient mitochondrial function. The expression of the components of certain mitochondrial pathways may therefore be mediated by miRNAs at the subcellular level. Such regulation might even extend to simultaneous changes in miRNA expression in multiple subcellular compartments. Our understanding of mitochondrial miRNA biogenesis, physiology and function will only increase as interest in this field continues to grow.

COMPETING INTERESTS

None declared.

AUTHOR CONTRIBUTIONS

G.W. and S.L. provided a conceptual framework for the review. J.S. and S.L. searched and interpreted the literature. All authors wrote and revised the manuscript. All authors approved the final version of the manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

ORCID

Séverine Lamon (i) http://orcid.org/0000-0002-3271-6551

REFERENCES

Anderson, S., Bankier, A. T., Barrell, B. G., de Bruijn, M. H., Coulson, A. R., Drouin, J., ... Young, I. G. (1981). Sequence and organization of the human mitochondrial genome. *Nature*, 290, 457– 465.

- Andrews, R. M., Kubacka, I., Chinnery, P. F., Lightowlers, R. N., Turnbull, D. M., & Howell, N. (1999). Reanalysis and revision of the Cambridge reference sequence for human mitochondrial DNA. *Nature Genetics*, 23, 147. https://doi.org/10.1038/13779
- Aoi, W., Naito, Y., Mizushima, K., Takanami, Y., Kawai, Y., Ichikawa, H., & Yoshikawa, T. (2010). The microRNA miR-696 regulates PGC-1α in mouse skeletal muscle in response to physical activity. *American Journal of Physiology-Endocrinology and Metabolism*, 298, E799–E806. https://doi.org/10.1152/ajpendo.00448.2009
- Bandiera, S., Mategot, R., Girard, M., Demongeot, J., & Henrion-Caude, A. (2013). MitomiRs delineating the intracellular localization of micro-RNAs at mitochondria. *Free Radical Biology & Medicine*, 64, 12–19. https://doi.org/10.1016/j.freeradbiomed.2013.06.013
- Bandiera, S., Ruberg, S., Girard, M., Cagnard, N., Hanein, S., Chretien, D., ... Henrion-Caude, A. (2011). Nuclear outsourcing of RNA interference components to human mitochondria. *PLoS One*, *6*, e20746. https://doi.org/10.1371/journal.pone.0020746
- Barrey, E., Saint-Auret, G., Bonnamy, B., Damas, D., Boyer, O., & Gidrol, X. (2011). Pre-microRNA and mature microRNA in human mitochondria. *PLoS One*, 6, e20220. https://doi.org/10.1371/journal.pone.0020220
- Bartel, D. P. (2004). MicroRNAs: Genomics, biogenesis, mechanism, and function. Cell, 116, 281–297.
- Bassi, R., Heads, R., Marber, M. S., & Clark, J. E. (2008). Targeting p38-MAPK in the ischaemic heart: Kill or cure? *Current Opinion in Pharmacology*, *8*, 141–146. https://doi.org/10.1016/j.coph.2008.01.002
- Bernstein, E., Caudy, A. A., Hammond, S. M., & Hannon, G. J. (2001). Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature*, 409, 363–366. https://doi.org/10.1038/35053110
- Bian, Z., Li, L. M., Tang, R., Hou, D. X., Chen, X., Zhang, C. Y., & Zen, K. (2010). Identification of mouse liver mitochondria-associated miRNAs and their potential biological functions. *Cell Research*, 20, 1076–1078. https://doi.org/10.1038/cr.2010.119
- Borralho, P. M., Rodrigues, C. M., & Steer, C. J. (2015). Micro-RNAs in mitochondria: An unexplored niche. Advances in Experimental Medicine and Biology, 887, 31–51. https://doi.org/ 10.1007/978-3-319-22380-3_3
- Bottini, S., Hamouda-Tekaya, N., Tanasa, B., Zaragosi, L. E., Grandjean, V., Repetto, E., & Trabucchi, M. (2017). From benchmarking HITS-CLIP peak detection programs to a new method for identification of miRNAbinding sites from Ago2-CLIP data. *Nucleic Acids Research*, 45, e71. https://doi.org/10.1093/nar/gkx007
- Brennecke, J., Stark, A., Russell, R. B., & Cohen, S. M. (2005). Principles of microRNA-target recognition. *PLoS Biology*, *3*, e85. https://doi.org/10.1371/journal.pbio.0030085
- Butow, R. A., & Avadhani, N. G. (2004). Mitochondrial signaling: The retrograde response. *Molecular Cell*, 14, 1–15.
- Carthew, R. W., & Sontheimer, E. J. (2009). Origins and mechanisms of miRNAs and siRNAs. *Cell*, 136, 642–655. https://doi.org/ 10.1016/j.cell.2009.01.035
- Cesana, M., Cacchiarelli, D., Legnini, I., Santini, T., Sthandier, O., Chinappi, M., ... Bozzoni, I. (2011). A long noncoding RNA controls muscle differentiation by functioning as a competing endogenous RNA. *Cell*, 147, 358–369. https://doi.org/10.1016/j.cell.2011.09.028
- Chendrimada, T. P., Gregory, R. I., Kumaraswamy, E., Norman, J., Cooch, N., Nishikura, K., & Shiekhattar, R. (2005). TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing. *Nature*, 436, 740– 744. https://doi.org/10.1038/nature03868
- Chi, S. W., Hannon, G. J., & Darnell, R. B. (2012). An alternative mode of microRNA target recognition. *Nature Structural & Molecular Biology*, 19, 321–327. https://doi.org/10.1038/nsmb.2230

- Das, S., Ferlito, M., Kent, O. A., Fox-Talbot, K., Wang, R., Liu, D., ... Steenbergen, C. (2012). Nuclear miRNA regulates the mitochondrial genome in the heart. *Circulation Research*, 110, 1596–1603. https://doi.org/10.1161/CIRCRESAHA.112.267732
- Dasgupta, N., Peng, Y., Tan, Z., Ciraolo, G., Wang, D., & Li, R. (2015). miRNAs in mtDNA-less cell mitochondria. *Cell Death Discovery*, 1, 15004. https://doi.org/10.1038/cddiscovery.2015.4
- Djebali, S., Davis, C. A., Merkel, A., Dobin, A., Lassmann, T., Mortazavi, A., ... Gingeras, T. R. (2012). Landscape of transcription in human cells. *Nature*, 489, 101–108. https://doi.org/10.1038/nature11233
- Dong, Y., Yoshitomi, T., Hu, J. F., & Cui, J. (2017). Long noncoding RNAs coordinate functions between mitochondria and the nucleus. *Epigenetics & Chromatin*, 10, 41. https://doi.org/ 10.1186/s13072-017-0149-x
- Drummond, M. J., McCarthy, J. J., Fry, C. S., Esser, K. A., & Rasmussen, B. B. (2008). Aging differentially affects human skeletal muscle microRNA expression at rest and after an anabolic stimulus of resistance exercise and essential amino acids. *American Journal* of Physiology-Endocrinology and Metabolism, 295, E1333–E1340. https://doi.org/10.1152/ajpendo.90562.2008
- Duarte, F. V., Palmeira, C. M., & Rolo, A. P. (2014). The role of microRNAs in mitochondria: Small players acting wide. *Genes (Basel)*, *5*, 865–886. https://doi.org/10.3390/genes5040865
- Duarte, F. V., Palmeira, C. M., & Rolo, A. P. (2015). The emerging role of mitomiRs in the pathophysiology of human disease. *Advances in Experimental Medicine and Biology*, 888, 123–154. https://doi.org/10.1007/978-3-319-22671-2_8
- Eisenberg, I., Alexander, M. S., & Kunkel, L. M. (2009). miRNAS in normal and diseased skeletal muscle. *Journal of Cellular and Molecular Medicine*, 13, 2–11. https://doi.org/10.1111/j.1582-4934.2008.00524.x
- Fernández-Vizarra, E., Enríquez, J. A., Pérez-Martos, A., Montoya, J., & Fernández-Silva, P. (2011). Tissue-specific differences in mitochondrial activity and biogenesis. *Mitochondrion*, 11, 207–213. https://doi.org/10.1016/j.mito.2010.09.011
- Friedman, R. C., Farh, K. K., Burge, C. B., & Bartel, D. P. (2009). Most mammalian mRNAs are conserved targets of microRNAs. *Genome Research*, 19, 92–105. https://doi.org/10.1101/gr.082701.108
- Fröhlich, K. S., & Vogel, J. (2009). Activation of gene expression by small RNA. Current Opinion in Microbiology, 12, 674–682. https://doi.org/10.1016/j.mib.2009.09.009
- Gambardella, S., Rinaldi, F., Lepore, S. M., Viola, A., Loro, E., Angelini, C., ... Botta, A. (2010). Overexpression of microRNA-206 in the skeletal muscle from myotonic dystrophy type 1 patients. *Journal of Translational Medicine*, *8*, 48. https://doi.org/10.1186/1479-5876-8-48
- Gao, S., Tian, X., Chang, H., Sun, Y., Wu, Z., Cheng, Z., ... Bu, W. (2018). Two novel IncRNAs discovered in human mitochondrial DNA using PacBio full-length transcriptome data. *Mitochondrion*, 38, 41–47. https://doi.org/10.1016/j.mito.2017.08.002
- Garros, R. F., Paul, R., Connolly, M., Lewis, A., Garfield, B. E., Natanek, S. A., ... Kemp, P. R. (2017). MicroRNA-542 promotes mitochondrial dysfunction and smad activity and is elevated in intensive care unit-acquired weakness. American Journal of Respiratory and Critical Care Medicine, 196, 1422–1433. https://doi.org/10.1164/rccm.201701-01010C
- Geiger, J., & Dalgaard, L. T. (2017). Interplay of mitochondrial metabolism and microRNAs. *Cellular and Molecular Life Sciences*, 74, 631–646. https://doi.org/10.1007/s00018-016-2342-7
- Gibbings, D. J., Ciaudo, C., Erhardt, M., & Voinnet, O. (2009). Multivesicular bodies associate with components of miRNA effector complexes and modulate miRNA activity. *Nature Cell Biology*, 11, 1143–1149. https://doi.org/10.1038/ncb1929

- Greco, S., De Simone, M., Colussi, C., Zaccagnini, G., Fasanaro, P., Pescatori, M., ... Martelli, F. (2009). Common micro-RNA signature in skeletal muscle damage and regeneration induced by Duchenne muscular dystrophy and acute ischemia. *FASEB Journal*, 23, 3335–3346. https://doi.org/10.1096/fj.08-128579
- Griffiths-Jones, S., Saini, H. K., van Dongen, S., & Enright, A. J. (2008). miRBase: Tools for microRNA genomics. *Nucleic Acids Research*, 36, Database issue, D154–D158. https://doi.org/10.1093/nar/gkm952
- & Vissing, K. (2017). Impact of resistance Groennebaek. T., training mitochondrial muscle biogenesis, on skeletal Physiology, and function. Frontiers in content. 8, 713. https://doi.org/10.3389/fphys.2017.00713
- Guo, H., Ingolia, N. T., Weissman, J. S., & Bartel, D. P. (2010). Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature*, 466, 835–840. https://doi.org/10.1038/nature09267
- Ha, M., & Kim, V. N. (2014). Regulation of microRNA biogenesis. Nature Reviews Molecular Cell Biology, 15, 509–524. https://doi.org/10.1038/nrm3838
- Helwak, A., Kudla, G., Dudnakova, T., & Tollervey, D. (2013). Mapping the human miRNA interactome by CLASH reveals frequent noncanonical binding. *Cell*, 153, 654–665. https://doi.org/10.1016/j.cell.2013.03.043
- Herst, P. M., Rowe, M. R., Carson, G. M., & Berridge, M. V. (2017). Functional mitochondria in health and disease. *Frontiers in Endocrinology (Lausanne)*, 8, 296. https://doi.org/10.3389/fendo.2017.00296
- Hombach, S., & Kretz, M. (2016). Non-coding RNAs: Classification, biology and functioning. Advances in Experimental Medicine and Biology, 937, 3– 17. https://doi.org/10.1007/978-3-319-42059-2_1
- Huang, L., Mollet, S., Souquere, S., Le Roy, F., Ernoult-Lange, M., Pierron, G., ... Weil, D. (2011). Mitochondria associate with P-bodies and modulate microRNA-mediated RNA interference. *The Journal of Biological Chemistry*, 286, 24219–24230. https://doi.org/10.1074/jbc.M111.240259
- Humphreys, D. T., Westman, B. J., Martin, D. I., & Preiss, T. (2005). Micro-RNAs control translation initiation by inhibiting eukaryotic initiation factor 4E/cap and poly(A) tail function. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 16961–16966. https://doi.org/10.1073/pnas.0506482102
- Hutvagner, G., & Simard, M. J. (2008). Argonaute proteins: Key players in RNA silencing. Nature Reviews Molecular Cell Biology, 9, 22–32. https://doi.org/10.1038/nrm2321
- Jagannathan, R., Thapa, D., Nichols, C. E., Shepherd, D. L., Stricker, J. C., Croston, T. L., ... Hollander, J. M. (2015). Translational regulation of the mitochondrial genome following redistribution of mitochondrial micro-RNA in the diabetic heart. *Circulation. Cardiovascular genetics*, *8*, 785– 802. https://doi.org/10.1161/CIRCGENETICS.115.001067
- Janowski, B. A., Huffman, K. E., Schwartz, J. C., Ram, R., Nordsell, R., Shames, D. S., ... Corey, D. R. (2006). Involvement of AGO1 and AGO2 in mammalian transcriptional silencing. *Nature Structural & Molecular Biology*, 13, 787–792. https://doi.org/10.1038/nsmb1140
- Jarroux, J., Morillon, A., & Pinskaya, M. (2017). History, discovery, and classification of IncRNAs. Advances in Experimental Medicine and Biology, 1008, 1–46. https://doi.org/10.1007/978-981-10-5203-3_1
- Johnson, C. D., Esquela-Kerscher, A., Stefani, G., Byrom, M., Kelnar, K., Ovcharenko, D., ... Slack, F. J. (2007). The *let-7* microRNA represses cell proliferation pathways in human cells. *Cancer Research*, *67*, 7713–7722. https://doi.org/10.1158/0008-5472.CAN-07-1083
- Kim, Y. J., Maizel, A., & Chen, X. (2014). Traffic into silence: Endomembranes and post-transcriptional RNA silencing. *The EMBO Journal*, 33, 968–980. https://doi.org/10.1002/embj.201387262

- Kirby, T. J., & McCarthy, J. J. (2013). MicroRNAs in skeletal muscle biology and exercise adaptation. *Free Radical Biology and Medicine*, 64, 95–105. https://doi.org/10.1016/j.freeradbiomed.2013.07.004
- Kozomara, A., & Griffiths-Jones, S. (2014). miRBase: Annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Research*, 42, Database issue, D68–D73. https://doi.org/10.1093/nar/gkt1181
- Kren, B. T., Wong, P. Y., Sarver, A., Zhang, X., Zeng, Y., & Steer, C. J. (2009). MicroRNAs identified in highly purified liver-derived mitochondria may play a role in apoptosis. *RNA Biology*, *6*, 65–72.
- Kung, J. T., Colognori, D., & Lee, J. T. (2013). Long noncoding RNAs: Past, present, and future. *Genetics*, 193, 651–669. https://doi.org/10.1534/genetics.112.146704
- Lamon, S., Zacharewicz, E., Butchart, L. C., Orellana, L., Mikovic, J., Grounds, M. D., & Russell, A. P. (2017). MicroRNA expression patterns in postnatal mouse skeletal muscle development. *BMC Genomics*, 18, 52. https://doi.org/10.1186/s12864-016-3399-2
- Landerer, E., Villegas, J., Burzio, V. A., Oliveira, L., Villota, C., Lopez, C., ... Burzio, L. O. (2011). Nuclear localization of the mitochondrial ncRNAs in normal and cancer cells. *Cell Oncolology (Dordr)*, 34, 297–305. https://doi.org/10.1007/s13402-011-0018-8
- Lee, R. C., & Ambros, V. (2001). An extensive class of small RNAs in *Caenorhabditis elegans. Science*, 294, 862.
- Lee, R. C., Feinbaum, R. L., & Ambros, V. (1993). The C. elegans heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. Cell, 75, 843–854.
- Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., ... Kim, V. N. (2003). The nuclear RNase III Drosha initiates microRNA processing. *Nature*, 425, 415–419. https://doi.org/10.1038/nature01957
- Leung, A. K., Calabrese, J. M., & Sharp, P. A. (2006). Quantitative analysis of Argonaute protein reveals microRNA-dependent localization to stress granules. *Proceedings of the National Academy* of Sciences of the United States of America, 103, 18125–18130. https://doi.org/10.1073/pnas.0608845103
- Licatalosi, D. D., Mele, A., Fak, J. J., Ule, J., Kayikci, M., Chi, S. W., ... Darnell, R. B. (2008). HITS-CLIP yields genome-wide insights into brain alternative RNA processing. *Nature*, 456, 464–469. https://doi.org/10.1038/nature07488
- Lima, T. I., Araujo, H. N., Menezes, E. S., Sponton, C. H., Araujo, M. B., Bomfim, L. H., ... Silveira, L. R. (2017). Role of microRNAs on the regulation of mitochondrial biogenesis and insulin signaling in skeletal muscle. *Journal of Cell Physiology*, 232, 958–966. https://doi.org/10.1002/jcp.25-645
- Little, J. P., Safdar, A., Benton, C. R., & Wright, D. C. (2011). Skeletal muscle and beyond: The role of exercise as a mediator of systemic mitochondrial biogenesis. *Applied Physiology, Nutrition, and Metabolism*, 36, 598–607. https://doi.org/10.1139/h11-076
- Liu, J., Carmell, M. A., Rivas, F. V., Marsden, C. G., Thomson, J. M., Song, J. J., ... Hannon, G. J. (2004). Argonaute2 is the catalytic engine of mammalian RNAi. *Science*, 305, 1437–1441. https://doi.org/10.1126/science.1102513
- Long, J., Badal, S. S., Ye, Z., Wang, Y., Ayanga, B. A., Galvan, D. L., ... Danesh, F. R. (2016). Long noncoding RNA *Tug1* regulates mitochondrial bioenergetics in diabetic nephropathy. *The Journal of Clinical Investigation*, 126, 4205–4218. https://doi.org/10.1172/JCI87927
- Lund, E., Güttinger, S., Calado, A., Dahlberg, J. E., & Kutay, U. (2004). Nuclear export of microRNA precursors. *Science*, 303, 95–98. https://doi.org/10.1126/science.1090599
- Lung, B., Zemann, A., Madej, M. J., Schuelke, M., Techritz, S., Ruf, S., ... Huttenhofer, A. (2006). Identification of small non-coding RNAs from

mitochondria and chloroplasts. Nucleic Acids Research, 34, 3842–3852. https://doi.org/10.1093/nar/gkl448

- Macgregor-Das, A. M., & Das, S. (2018). A microRNA's journey to the center of the mitochondria. American Journal of Physiology-Heart and Circulatory Physiology, in press. https://doi.org/10.1152/ajpheart.00714.2017
- Maniataki, E., & Mourelatos, Z. (2005). Human mitochondrial tRNAMet is exported to the cytoplasm and associates with the Argonaute 2 protein. *RNA*, 11, 849–852. https://doi.org/10.1261/rna.2210805
- Massart, J., Katayama, M., & Krook, A. (2016). microManaging glucose and lipid metabolism in skeletal muscle: Role of microRNAs. *Biochimica et Biophysica Acta*, 1861, 2130–2138. https://doi.org/10.1016/j.bbalip.2016.05.006
- Matera, A. G., Terns, R. M., & Terns, M. P. (2007). Non-coding RNAs: Lessons from the small nuclear and small nucleolar RNAs. *Nature Reviews Molecular Cell Biology*, 8, 209–220. https://doi.org/10.1038/nrm2124
- McCarthy, J. J. (2011). The MyomiR network in skeletal muscle plasticity. Exercise and Sport Sciences Reviews, 39, 150–154. https://doi.org/10.1097/JES.0b013e31821c01e1
- Meister, G., Landthaler, M., Patkaniowska, A., Dorsett, Y., Teng, G., & Tuschl, T. (2004). Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Molecular Cell*, 15, 185–197. https://doi.org/10.1016/j.molcel.2004.07.007
- Mercer, T. R., Neph, S., Dinger, M. E., Crawford, J., Smith, M. A., Shearwood, A. M., ... Mattick, J. S. (2011). The human mitochondrial transcriptome. *Cell*, 146, 645–658. https://doi.org/10.1016/j.cell.2011.06.051
- Naguibneva, I., Ameyar-Zazoua, M., Polesskaya, A., Ait-Si-Ali, S., Groisman, R., Souidi, M., ... Harel-Bellan, A. (2006). The micro-RNA *miR-181* targets the homeobox protein Hox-A11 during mammalian myoblast differentiation. *Nature Cell Biology*, *8*, 278. https://doi.org/10.1038/ncb1373
- Nie, Y., Sato, Y., Wang, C., Yue, F., Kuang, S., & Gavin, T. P. (2016). Impaired exercise tolerance, mitochondrial biogenesis, and muscle fiber maintenance in miR-133a-deficient mice. FASEB Journal, 30, 3745– 3758. https://doi.org/10.1096/fj.201600529R
- Nielsen, S., Scheele, C., Yfanti, C., Akerstrom, T., Nielsen, A. R., Pedersen, B. K., & Laye, M. J. (2010). Muscle specific microRNAs are regulated by endurance exercise in human skeletal muscle. *The Journal of Physiology*, 588, 4029–4037. https://doi.org/10.1113/jphysiol.2010.189860
- Nouws, J., & Shadel, G. S. (2014). microManaging mitochondrial translation. *Cell*, 158, 477–478. https://doi.org/10.1016/j.cell.2014.07.010
- Nunnari, J., & Suomalainen, A. (2012). Mitochondria: In sickness and in health. Cell, 148, 1145–1159. https://doi.org/ 10.1016/j.cell.2012.02.035
- Ogata, T., & Yamasaki, Y. (1997). Ultra-high-resolution scanning electron microscopy of mitochondria and sarcoplasmic reticulum arrangement in human red, white, and intermediate muscle fibers. *The Anatomical Record*, 248, 214–223.
- Ohrt, T., Mütze, J., Staroske, W., Weinmann, L., Höck, J., Crell, K., ... Schwille, P. (2008). Fluorescence correlation spectroscopy and fluorescence cross-correlation spectroscopy reveal the cytoplasmic origination of loaded nuclear RISC *in vivo* in human cells. *Nucleic Acids Research, 36*, 6439–6449. https://doi.org/10.1093/nar/gkn693
- Olsen, P. H., & Ambros, V. (1999). The lin-4 regulatory RNA controls developmental timing in *Caenorhabditis elegans* by blocking LIN-14 protein synthesis after the initiation of translation. *Developmental Biology*, 216, 671-680. https://doi.org/10.1006/dbio.1999.9523
- Pasquinelli, A. E. (2012). MicroRNAs and their targets: Recognition, regulation and an emerging reciprocal relationship. *Nature Reviews Genetics*, 13, 271–282. https://doi.org/10.1038/nrg3162

- Pinto, S. K., Lamon, S., Stephenson, E. J., Kalanon, M., Mikovic, J., Koch, L. G., ... Camera, D. M. (2017). Expression of microRNAs and target proteins in skeletal muscle of rats selectively bred for high and low running capacity. American Journal of Physiology-Endocrinology and Metabolism, 313, E335–E343. https://doi.org/10.1152/ajpendo.00043.2017
- Politz, J. C., Zhang, F., & Pederson, T. (2006). MicroRNA-206 colocalizes with ribosome-rich regions in both the nucleolus and cytoplasm of rat myogenic cells. Proceedings of the National Academy of Sciences of the United States of America, 103, 18957–18962. https://doi.org/10.1073/pnas.0609466103
- Ponting, C. P., Oliver, P. L., & Reik, W. (2009). Evolution and functions of long noncoding RNAs. *Cell*, 136, 629–641. https://doi.org/10.1016/j.cell.2009.02.006
- Rackham, O., Shearwood, A. M., Mercer, T. R., Davies, S. M., Mattick, J. S., & Filipovska, A. (2011). Long noncoding RNAs are generated from the mitochondrial genome and regulated by nuclear-encoded proteins. *RNA*, 17, 2085–2093. https://doi.org/10.1261/rna.029405.111
- Radziuk, J., & Inculet, R. (1983). The effects of ingested and intravenous glucose on forearm uptake of glucose and glucogenic substrate in normal man. *Diabetes*, 32, 977–981.
- Rasmussen, B. B., & Phillips, S. M. (2003). Contractile and nutritional regulation of human muscle growth. *Exercise and Sport Sciences Reviews*, 31, 127–131.
- Reinhart, B. J., Slack, F. J., Basson, M., Pasquinelli, A. E., Bettinger, J. C., Rougvie, A. E., ... Ruvkun, G. (2000). The 21-nucleotide *let-7* RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature*, 403, 901–906. https://doi.org/10.1038/35002607
- Ringholm, S., Bienso, R. S., Kiilerich, K., Guadalupe-Grau, A., Aachmann-Andersen, N. J., Saltin, B., ... Pilegaard, H. (2011). Bed rest reduces metabolic protein content and abolishes exercise-induced mRNA responses in human skeletal muscle. *American Journal* of *Physiology-Endocrinology and Metabolism*, 301, E649–E658. https://doi.org/10.1152/ajpendo.00230.2011
- Ro, S., Ma, H. Y., Park, C., Ortogero, N., Song, R., Hennig, G. W., ... Yan, W. (2013). The mitochondrial genome encodes abundant small noncoding RNAs. *Cell Research*, 23, 759–774. https://doi.org/10.1038/cr.2013.37
- Russell, A. P., Foletta, V. C., Snow, R. J., & Wadley, G. D. (2014). Skeletal muscle mitochondria: A major player in exercise, health and disease. *Biochimica et Biophysica Acta*, 1840, 1276–1284. https://doi.org/10.1016/j.bbagen.2013.11.016
- Russell, A. P., Ghobrial, L., Ngo, S., Yerbury, J., Zacharewicz, E., Chung, R., & Lamon, S. (2018). Dysregulation of microRNA biogenesis machinery and microRNA/RNA ratio in skeletal muscle of amyotrophic lateral sclerosis mice. *Muscle Nerve*, 57, 838–847. https://doi.org/10.1002/mus.26039
- Russell, A. P., Lamon, S., Boon, H., Wada, S., Guller, I., Brown, E. L., ... Akimoto, T. (2013). Regulation of miRNAs in human skeletal muscle following acute endurance exercise and short-term endurance training. *The Journal of Physiology*, 591, 4637–4653. https://doi.org/10.1113/jphysiol.2013.255695
- Russell, A. P., Wada, S., Vergani, L., Hock, M. B., Lamon, S., Leger, B., ... Akimoto, T. (2013). Disruption of skeletal muscle mitochondrial network genes and miRNAs in amyotrophic lateral sclerosis. *Neurobiology of Disease*, 49, 107–117. https://doi.org/10.1016/j.nbd.2012.08.015
- Safdar, A., Abadi, A., Akhtar, M., Hettinga, B. P., & Tarnopolsky, M. A. (2009). miRNA in the regulation of skeletal muscle adaptation to acute endurance exercise in C57BI/6J male mice. *PLoS One*, *4*, e5610. https://doi.org/10.1371/journal.pone.0005610
- Salinas, T., Duchene, A. M., Delage, L., Nilsson, S., Glaser, E., Zaepfel, M., & Marechal-Drouard, L. (2006). The voltage-dependent anion channel, a major component of the tRNA import machinery

in plant mitochondria. Proceedings of the National Academy of Sciences of the United States of America, 103, 18362–18367. https://doi.org/10.1073/pnas.0606449103

- Salmanidis, M., Pillman, K., Goodall, G., & Bracken, C. (2014). Direct transcriptional regulation by nuclear microRNAs. *The International Journal of Biochemistry & Cell Biology*, 54, 304–311. https://doi.org/10.1016/j.biocel.2014.03.010
- Shabalina, S. A., & Koonin, E. V. (2008). Origins and evolution of eukaryotic RNA interference. *Trends in Ecology & Evolution*, 23, 578–587. https://doi.org/10.1016/j.tree.2008.06.005
- Shepherd, D. L., Hathaway, Q. A., Pinti, M. V., Nichols, C. E., Durr, A. J., Sreekumar, S., ... Hollander, J. M. (2017). Exploring the mitochondrial microRNA import pathway through polynucleotide phosphorylase (PNPase). Journal of Molecular and Cellular Cardiology, 110, 15–25. https://doi.org/10.1016/j.yjmcc.2017.06.012
- Shinde, S., & Bhadra, U. (2015). A complex genome-microRNA interplay in human mitochondria. *BioMed Research International*, 2015, 206382. https://doi.org/10.1155/2015/206382
- Slomovic, S., Laufer, D., Geiger, D., & Schuster, G. (2005). Polyadenylation and degradation of human mitochondrial RNA: The prokaryotic past leaves its mark. *Molecular and Cellular Biology*, 25, 6427–6435. https://doi.org/10.1128/mcb.25.15.6427-6435.2005
- Sripada, L., Tomar, D., Prajapati, P., Singh, R., Singh, A. K., & Singh, R. (2012). Systematic analysis of small RNAs associated with human mitochondria by deep sequencing: Detailed analysis of mitochondrial associated miRNA. *PLoS One*, 7, e44873. https://doi.org/10.1371/journal.pone.0044873
- Sripada, L., Tomar, D., & Singh, R. (2012). Mitochondria: One of the destinations of miRNAs. *Mitochondrion*, 12, 593–599. https://doi.org/10.1016/j.mito.2012.10.009
- Taanman, J. W. (1999). The mitochondrial genome: Structure, transcription, translation and replication. *Biochimica et Biophysica Acta*, 1410, 103– 123.
- Tomari, Y., & Zamore, P. D. (2005). Perspective: Machines for RNAi. Genes & Development, 19, 517–529. https://doi.org/10.1101/gad.1284105
- Tsitkanou, S., Della Gatta, P. A., & Russell, A. P. (2016). Skeletal muscle satellite cells, mitochondria, and microRNAs: Their involvement in the pathogenesis of ALS. *Frontiers in Physiology*, 7, 403. https://doi.org/10.3389/fphys.2016.00403
- Vargas, J. N., Kar, A. N., Kowalak, J. A., Gale, J. R., Aschrafi, A., Chen, C. Y., ... Kaplan, B. B. (2016). Axonal localization and mitochondrial association of precursor microRNA 338. *Cellular and Molecular Life Sciences*, 73, 4327–4340. https://doi.org/10.1007/s00018-016-2270-6
- Vasudevan, S., Tong, Y., & Steitz, J. A. (2007). Switching from repression to activation: MicroRNAs can up-regulate translation. *Science*, 318, 1931– 1934. https://doi.org/10.1126/science.1149460
- Wada, S., Kato, Y., Sawada, S., Aizawa, K., Park, J. H., Russell, A. P., ... Akimoto, T. (2015). MicroRNA-23a has minimal effect on endurance exerciseinduced adaptation of mouse skeletal muscle. *Pflugers Archiv*, 467, 389– 398. https://doi.org/10.1007/s00424-014-1517-z
- Wang, G., Chen, H. W., Oktay, Y., Zhang, J., Allen, E. L., Smith, G. M., ... Teitell, M. A. (2010). PNPASE regulates RNA import into mitochondria. *Cell*, 142, 456–467. https://doi.org/10.1016/j.cell.2010.06.035
- Wang, H., Garzon, R., Sun, H., Ladner, K. J., Singh, R., Dahlman, J., ... Guttridge, D. C. (2008). NF-κB-YY1-miR-29 regulatory circuitry in skeletal myogenesis and rhabdomyosarcoma. *Cancer Cell*, 14, 369–381. https://doi.org/10.1016/j.ccr.2008.10.006
- Wang, W. X., Visavadiya, N. P., Pandya, J. D., Nelson, P. T., Sullivan, P. G., & Springer, J. E. (2015). Mitochondria-associated microRNAs in rat hippocampus following traumatic brain injury. *Experimental Neurology*, 265, 84–93. https://doi.org/10.1016/j.expneurol.2014.12.018

- Wang, X., Song, X., Glass, C. K., & Rosenfeld, M. G. (2011). The long arm of long noncoding RNAs: Roles as sensors regulating gene transcriptional programs. *Cold Spring Harbor Perspectives in Biology*, *3*, a003756. https://doi.org/10.1101/cshperspect.a003756
- Williams, A. H., Valdez, G., Moresi, V., Qi, X., McAnally, J., Elliott, J. L., ... Olson, E. N. (2009). MicroRNA-206 delays ALS progression and promotes regeneration of neuromuscular synapses in mice. *Science*, 326, 1549–1554. https://doi.org/10.1126/science.1181046
- Wilusz, J. E., JnBaptiste, C. K., Lu, L. Y., Kuhn, C. D., Joshua-Tor, L., & Sharp, P. A. (2012). A triple helix stabilizes the 3' ends of long noncoding RNAs that lack poly(A) tails. *Genes & Development*, 26, 2392–2407. https://doi.org/10.1101/gad.204438.112
- Wright, D. C., Han, D.-H., Garcia-Roves, P. M., Geiger, P. C., Jones, T. E., & Holloszy, J. O. (2007). Exercise-induced mitochondrial biogenesis begins before the increase in muscle PGC-1α expression. Journal of Biological Chemistry, 282, 194–199.
- Yamamoto, H., Morino, K., Nishio, Y., Ugi, S., Yoshizaki, T., Kashiwagi, A., & Maegawa, H. (2012). MicroRNA-494 regulates mitochondrial biogenesis in skeletal muscle through mitochondrial transcription factor A and Forkhead box j3. American Journal of Physiology-Endocrinology and Metabolism, 303, E1419–E1427. https://doi.org/10.1152/ajpendo.00097.2012
- Yan, Z., Lira, V. A., & Greene, N. P. (2012). Exercise training-induced regulation of mitochondrial quality. *Exercise and Sport Sciences Reviews*, 40, 159–164. https://doi.org/10.1097/JES.0b013e3182575599
- Zacharewicz, E., Della Gatta, P., Reynolds, J., Garnham, A., Crowley, T., Russell, A. P., & Lamon, S. (2014). Identification of micro-RNAs linked to regulators of muscle protein synthesis and regeneration in young and old skeletal muscle. *PLoS One*, 9, e114009. https://doi.org/10.1371/journal.pone.0114009
- Zacharewicz, E., Lamon, S., & Russell, A. P. (2013). MicroRNAs in skeletal muscle and their regulation with exercise, ageing, and disease. *Frontiers in Physiology*, *4*, 266. https://doi.org/10.3389/fphys.2013.00266
- Zhang, C. (2009). Novel functions for small RNA molecules. *Current Opinion* in Molecular Therapeutics, 11, 641–651.
- Zhang, X., Zuo, X., Yang, B., Li, Z., Xue, Y., Zhou, Y., ... Fu, X. D. (2014). MicroRNA directly enhances mitochondrial translation during muscle differentiation. *Cell*, 158, 607–619. https://doi.org/10.1016/j.cell.2014.05.047
- Zurlo, F., Larson, K., Bogardus, C., & Ravussin, E. (1990). Skeletal muscle metabolism is a major determinant of resting energy expenditure. *Journal of Clinical Investigation*, 86, 1423– 1427.https://doi.org/10.1172/JCI114857

AUTHOR'S BIOGRAPHY



Séverine Lamon graduated from the University of Lausanne, Switzerland, in 2010. She is a former Swiss National Science Foundation (SNSF) and Australian Research Council (ARC) early career fellowship recipient, and currently a senior lecturer at Deakin University, Australia. Her research focuses on understanding the factors

that underlie the regulation of skeletal muscle mass in health and disease conditions, with a specific interest in the role of non-coding RNAs in the molecular regulation of skeletal muscle growth. Her research program spans cell, rodent and human models, with the aim to ultimately develop applied interventions to maintain skeletal muscle mass in human subjects. **Glenn Wadley** is a muscle physiologist and

the associate Head of School (Research) of Deakin University's School of Exercise and Nutrition Sciences (SENS), ranked as the world's top sports science research school for 2016 and 2017 (ARWU University ranking). **Jessica Silver** is a PhD student supervised by Dr Lamon and A/Prof. Wadley.

ILEY

How to cite this article: Silver J, Wadley G, Lamon S. Mitochondrial regulation in skeletal muscle: A role for non-coding RNAs? *Exp Physiol.* 2018;103:1132-1144. https://doi.org/10.1113/EP086846