

## FLOWERING AND FRUITING RESPONSES OF STRAWBERRY TO GROWTH HORMONE AND CHILLING GROWN UNDER TUNNEL CONDITIONS

Samia Ikram<sup>1,2\*</sup>, Khalid Mahmood Qureshi<sup>1</sup> and Nauman Khalid<sup>3,4</sup>

<sup>1</sup>Department of Horticulture, Pir Mehar Ali Shah Arid Agriculture University Rawalpindi 43000, Pakistan;

<sup>2</sup>Horticultural Research Institute for Floriculture and Landscaping, Orchard Scheme Area Murree Road, Islamabad, Pakistan; <sup>3</sup>Algae Biomass and Energy System R&D Center, University of Tsukuba, 1-1-1 Tennoudai, Tsukuba, Ibaraki 305-8577 Japan; <sup>4</sup>School of Agriculture, University of Management and Technology, Lahore 54000, Pakistan.

\*Corresponding author's e-mail: samiaikram@hotmail.com

Strawberry (*Fragaria ananasa*) is a berry fruit belonging to the family Rosaceae and requires certain amount of cold temperature to initiate flowering. Present study was conducted with the objective to study flowering and fruiting responses of strawberry to growth hormone and chilling under tunnel conditions. The study comprised of six different treatments as; control, GA<sub>3</sub> (gibberellic acid) 200 ppm, GA<sub>3</sub> 400 ppm, chilling at 4°C, chilling at 4°C + GA<sub>3</sub> 200 ppm and chilling at 4°C + GA<sub>3</sub> 400 ppm. Treatments were applied at vegetative stage before flowering. It was inferred from the study that strawberry plants responded much better to application of GA<sub>3</sub> at 400 ppm for their growth, flower stimulation, their number, quality features while chilling at 4°C proved better for maximum fruit production and better quality under tunnel condition.

**Key words:** Strawberry, gibberellic acid, chilling, flowering, fruiting, tunnel conditions

### INTRODUCTION

Strawberry (*Fragaria ananasa*) a soft fruit that belongs to family Rosaceae is stimulated by cold weather. Strawberry is native to North America. Its name has been derived from farmers mulching called "Straw", earlier the strawberries were just planted for medicinal purpose but later on Carolus Linnaeus for the first time introduced strawberry as edible fruit (Lieten *et al.*, 1995). Strawberry is herbaceous perennial which is propagated by means of seed, short rhizomes and leafless stolons. The beautiful red colored fleshy fruit is full of achenes on its surface (Anderson, 2006). Strawberry is rich source of vitamins mainly vitamin C, fiber, potassium and phytochemical especially ellagic acid and anthocyanins which act as anti-inflammatory and antioxidant agents (Hannum, 2004).

Chilling is required for breaking the dormancy of strawberry (Lieten *et al.*, 1995), and to accelerate flowering ability (Sung and Amasino, 2004). There are several models that have been developed for measuring winter chill, e.g. the Chilling Hours Model (Luedeling *et al.*, 2010), the Utah Model and the Dynamic Model (Fishman *et al.*, 1987). These models differ from each other greatly in their sensitivity to climate change (Luedeling *et al.*, 2009), which makes the choice of the model a crucial determinant of forecasting the extent of climate change and effects on winter chill. A certain quantity of chilling has been shown to improve flower bud formation of the short-day strawberry plant (Taylor, 2002).

Vegetative and reproductive developments of strawberry under the control of genes are extremely vulnerable to environmental factors (Battey *et al.*, 1998). GA<sub>3</sub> appliance improved length of petiole and area of leaf in strawberry plants when used in different concentrations (Paroussi *et al.*, 2002). The application of GA<sub>3</sub> (50 ppm) to the latent buds could achieve 97% break of their onset of dormancy (Ahmed and Ragab, 2005). A treatment of 10 ppm GA<sub>3</sub> applied to fruit frost protected plants during the 2<sup>nd</sup> week of January (Ozguven and Yilmaz, 2002). Ozguven and Yilmaz (2002) also reported that GA<sub>3</sub> applied at 5, 10 and 20 ppm had no effect on the average fruit weight and total soluble solid content in cv. 'Camarosa'.

Many studies have shown that runner quality is a major factor that affects both productivity and fruit quality (Durner *et al.*, 2000). High vitamin C quantity has been observed in strawberry fruit plant treated with GA<sub>3</sub> during pre-flowering stage (Sharma and Singh, 2009). Parallel results have been reported when GA<sub>3</sub> was sprayed at pre-flowering stage, caused high vitamin C contents and total soluble solids and also enhanced firmness in capsicum (Ouzounidou *et al.*, 2010) and sweet cherry (Kappel and Macdonald, 2007). Gibberellic acid application also reported to boost the fruit firmness (Usenik *et al.*, 2005). Hence present study was designed to investigate the effect of chilling and growth hormone on the reproductive growth of strawberry plants and the role of GA<sub>3</sub> in fulfilling chilling requirement to stimulate flowering and fruiting.

### MATERIAL AND METHODS

**Chemicals:** Gibberellic acid, sodium hydroxide, hydrochloric acid and L-ascorbic acid were purchased from Sigma-Aldrich, Germany Milli-Q water with a resistivity of 18 MΩ cm was used to prepare all solutions. All of these chemicals were of analytical grade.

**Growth conditions and plant materials:** Present study was conducted in climatic conditions of Rawalpindi, Pakistan under plastic tunnel conditions. Runners of strawberry cv. “Chandler” were acquired from Swat, Pakistan. The runners were kept in cold chamber (Mack Pharmatech PVT Limited) at 4°C for about fifteen days to meet the chilling needs of strawberry plants. The runners were planted under field conditions in tunnels (well drained sandy loam soil, 22-25°C temperature with 65-75% humidity. Plantation was done on ridges by keeping plant to plant distance of 8 cm and row to row distance of 30 cm, irrigation was given soon after planting and then applied on weekly basis and as per plant requirement.

A foliar spray of GA<sub>3</sub> was applied before flowering after 45 days of plantation the concentration of GA<sub>3</sub> in combination with different chilling temperatures used were as follows: T<sub>1</sub>: control, T<sub>2</sub>: GA<sub>3</sub> 200 ppm, T<sub>3</sub>: GA<sub>3</sub> 400 ppm, T<sub>4</sub>: chilling 4°C, T<sub>5</sub>: chilling 4°C + GA<sub>3</sub> 200 ppm and T<sub>6</sub>: chilling 4°C + GA<sub>3</sub> 400 ppm.

**Parameters evaluated:** Following reproductive and quality parameters were studied; days to flower initiation, numbers of flowers, number of fruit set/plant, percentage fruit set, fruit weight (g), fruit size (cm) and firmness (pounds). Total soluble solids (TSS °Brix), L-ascorbic acid contents (mg/ml) and titratable acidity was determined by standard procedures (AOAC, 1990).

**Statistical analysis:** The experiment was laid down by using linear model randomized complete block design (RCBD). The collected observations were analyzed using Statistix 8.1. The means were compared by using Least Significant Difference (LSD) at 5% probability level (Steel *et al.*, 1997).

## RESULTS AND DISCUSSION

The data relating to the role of chilling and growth hormone affecting vegetative and reproductive growth of strawberry are presented in Table 1. The data was collected in the morning.

**Number of days required to open first flower:** It was observed (Table 1) that the lowest days (72.66) were taken by the plants treated with T<sub>5</sub> to start flower production followed by T<sub>3</sub> (74.33), T<sub>4</sub> (75.00), T<sub>6</sub> (75.33) the other treatments (T<sub>2</sub>) stimulated flower after 80.66 days and maximum days (90.33) were taken by T<sub>1</sub> to flower induction. Gibberellic acid functions to induce early flowering in all horticultural plants including strawberry. Sharma and Singh (2009) suggested that GA<sub>3</sub> positively influenced initiation, duration, number of flowers, harvesting season and yield of strawberry plant. Early flowering has also been reported in other horticulture crops when sprayed with GA<sub>3</sub> before flowering. Moreover, number of fruits per plants and fruit size was also increased (Ouzounidou *et al.*, 2010). Paroussi *et al.* (2002) reported that flowering was boosted by 200 mg L<sup>-1</sup> GA<sub>3</sub> treatment. GA<sub>3</sub> application caused noticeable inflorescence abnormalities especially in cv. Seascape, irrespective of photoperiod (Paroussi *et al.*, 2002) and the possible reason of inducing early flowering is due to the stimulation of Floragin hormone by gibberellic acid.

**Number of flowers:** The highest number of flowers per plant (25.33) was counted in treatment T<sub>3</sub> (400 ppm GA<sub>3</sub>). Non-significant ( $p>0.05$ ) data was shown by the rest of the treatments with the contribution of T<sub>4</sub> (24.88) T<sub>1</sub> (23.66), T<sub>5</sub> (21.33), T<sub>2</sub> (20.11) and minimum number of flowers (18.77) were counted for T<sub>6</sub>. Our results revealed that gibberellic acid and chilling help to produce more number of flowers as compared to control, similar results were stated by the previous work of many researchers that GA<sub>3</sub> application improved buds and open flowers thus, promoting earliness without effecting marketable yield (El-Shabasi *et al.*, 2009; Paroussi *et al.*, 2002). GA<sub>3</sub> at 10 ppm or ethrel at 250 ppm improved the number of flowers and total monthly yield (El-Shabasi *et al.*, 2009) (Table 1).

**Flower diameter:** The data related to flower diameter (cm) is presented in Table 1 which shows that maximum flower

**Table 1. Effect of chilling and growth hormone of strawberry on reproductive parameters of strawberry.**

| Treatment                                               | Number of days required to open first flower | Average number of flowers | Flower diameter         | No. of fruit set         | Fruit set (%)            | Total No. of fruits       |
|---------------------------------------------------------|----------------------------------------------|---------------------------|-------------------------|--------------------------|--------------------------|---------------------------|
| T <sub>1</sub> : Control                                | 90.33±0.66 <sup>A</sup>                      | 23.6±1.71 <sup>AB</sup>   | 3.32±0.10 <sup>AB</sup> | 12.50±1.07 <sup>AB</sup> | 90.70±6.68 <sup>AB</sup> | 26.20±1.25 <sup>ABC</sup> |
| T <sub>2</sub> : GA <sub>3</sub> 200 ppm                | 80.66±2.40 <sup>B</sup>                      | 20.10±0.40 <sup>BC</sup>  | 2.64±0.20 <sup>C</sup>  | 7.40±1.06 <sup>B</sup>   | 83.70±0.84 <sup>AB</sup> | 29.16±1.06 <sup>AB</sup>  |
| T <sub>3</sub> : GA <sub>3</sub> 400 ppm                | 74.33±0.28 <sup>C</sup>                      | 25.30±0.88 <sup>A</sup>   | 2.81±0.15 <sup>BC</sup> | 9.70±0.48 <sup>B</sup>   | 81.40±1.89 <sup>B</sup>  | 23.70±0.21 <sup>BC</sup>  |
| T <sub>4</sub> : Chilling 4°C                           | 75.00±0.57 <sup>C</sup>                      | 24.80±0.44 <sup>AC</sup>  | 3.34±0.42 <sup>A</sup>  | 15.40±0.78 <sup>A</sup>  | 93.70±2.99 <sup>A</sup>  | 31.50±0.76 <sup>A</sup>   |
| T <sub>5</sub> : Chilling 4°C (GA <sub>3</sub> 200 ppm) | 72.60±4.45 <sup>C</sup>                      | 21.30±2.41 <sup>ABC</sup> | 2.63±0.20 <sup>C</sup>  | 8.80±1.87 <sup>B</sup>   | 86.10±2.99 <sup>AB</sup> | 20.80±1.89 <sup>C</sup>   |
| T <sub>6</sub> : Chilling 4°C (GA <sub>3</sub> 400 ppm) | 75.30±1.76 <sup>C</sup>                      | 18.70±0.70 <sup>C</sup>   | 2.39±0.32 <sup>C</sup>  | 9.30±0.29 <sup>B</sup>   | 67.20±2.80 <sup>C</sup>  | 14.20±0.19 <sup>D</sup>   |

Means not sharing a letter differ significantly at  $P<0.05$ , ± = standard error.

diameter (3.34 cm) was observed in T<sub>4</sub> followed by T<sub>1</sub> (3.22 cm). The rest of the treatments helped to produce flower diameter of 2.81 cm at T<sub>3</sub>, 2.64 cm at T<sub>2</sub>, 2.63 cm at T<sub>5</sub> and the lowest 2.39 cm was produced at T<sub>6</sub>. Chilling temperatures were just as effective on reducing the time to first flower as previously reported (Risser and Robert, 1993). Chilling initiates the production of gibberellic acid and is also involved in breakdown of carbohydrates and other food reserves and their translocation to production sites that is why it help to produce larger fruit diameter in comparison with the rest of treatments and control.

**Number of fruit set:** Number of fruit set effected by chilling and exogenous application of gibberellic acid (Table 1) explained that maximum fruit set was found in T<sub>4</sub> (15.4 fruits per plant) chilling treatment while fruit number of 12.55, 9.77, 9.33, 8.88 and 7.44 were produced by T<sub>1</sub>, T<sub>3</sub>, T<sub>6</sub>, T<sub>5</sub> and T<sub>2</sub>, respectively. Chilling produced the highest fruit set in strawberry as it stimulates the production of endogenous gibberellins production which help to boost fruit set number as mentioned in previous literature. Gibberellins enhanced number of flower and flower bud per plant and also the total marketable yield (Kasim *et al.*, 2007; Sharma and Singh, 2009). Gibberellic acid significantly influences the initiation and duration of flowering and harvesting season, numbers of flowers per plant, fruit set and numbers of fruits per plant. It also increased the total yield and decreased the average fruit size (Sharma and Singh, 2009). Pre harvest foliar application of GA<sub>3</sub> at 50 ppm showed major increment in early yield (as compared to other treatments) of Globe artichoke whereas, no such increase was observed in late yield, this increase in early yield was due to early flowering thus no positive results were observed in total yield (Kasim *et al.*, 2007).

**Fruit set percentage:** The data pertaining to percentage fruit set is presented in Table 1. It describes that the highest 93.78 fruit percentage was noted in T<sub>4</sub> followed by T<sub>1</sub>(control) with the fruit set of 90.73%. The rest of the treatments contributed to 81.41% at T<sub>3</sub>, 86.18% at T<sub>5</sub>, 83.78% at T<sub>2</sub> and lowest of 67.24% at T<sub>6</sub>, respectively. The research findings proved that chilling contributed to the highest fruit set, the fruits produced were fresh and of good size and weight. GA<sub>3</sub> also contributed to good fruit set percentage but most of the fruits were albino and were not of marketable quality. The reason was that most of the biosynthate produced were used for runner production which resulted in poor fruit quality. These findings were in line with the work of Sharma and Singh (2009), who stated that gibberellic acid significantly increased the total yield and decreased the average fruit size.

**Total number of fruits:** The results obtained on total number of fruits presented in Table 1 proved that maximum number of total fruits (31.55) was obtained from treatment T<sub>4</sub> followed by three T<sub>2</sub> (29.16), T<sub>1</sub> (26.22), T<sub>3</sub> (23.78), T<sub>5</sub> (20.83) and T<sub>6</sub> (14.22) fruits, respectively. Chilling produced maximum number of fruit (put in fruit number) alone by stimulating the production of endogenous GA<sub>3</sub> as compared


to other treatments. Similar results were found by Sharma and Singh (2009) who reported that gibberellic acid increased the number of fruits comparable results in other fruits like capsicum. Foliar application of GA<sub>3</sub> notably accelerated the time of flowering by 8.3% as compared to control in *Capsicum annum* and also increased number of fruits per plant (Ouzounidou *et al.*, 2010). Single spray of gibberellic acid also induce fruit setting in sweet oranges, whereas did not affect final yield. More numbers of fruits were also reported while fruit size remains unaltered (Agusti *et al.*, 1982).

**Fruit size (cm):** The fruit size is presented in Figure 1a explains that maximum fruit size (20.66 cm) was observed in T<sub>4</sub> followed by T<sub>1</sub> (18.77 cm) T<sub>5</sub> (18.46 cm), T<sub>6</sub> (16.40), T<sub>3</sub> (15.78) and minimum of 15.40 cm by T<sub>2</sub> respectively. Chilling contributed to the maximum fruit size by transferring all the biosynthates to the fruits while the treatments with gibberellic acid used them for runner production as a result fruit quality decreased in term of fruit size and weight. Our findings are in accordance with the work of (Paroussi *et al.*, 2002) who reported that gibberellic acid enhanced the percentage of aborted flowers and malformed fruits, resulting in a major reduction in total marketable yield (Agusti *et al.*, 1982; Sharma and Singh, 2009). Gibberellic acid in late season on kiwi fruit canes increase in pedicel length but reduced berry weight and fruit diameter (Burge *et al.*, 1990). This might be due to sink of GA<sub>3</sub> treated plant's reserves towards early flower production and fruit setting and runner production.

**Fruit weight (g):** Fresh Fruit weight as affected by different treatments with chilling and exogenous application of gibberellic acid is presented in Figure 1b. It revealed that heaviest fruits (12.67 g) were obtained with T<sub>4</sub> chilling treatment alone followed by T<sub>1</sub> (control) with fruit weight of 10.66 which was then followed by T<sub>2</sub> (9.24 g), T<sub>5</sub> (7.38 g), T<sub>3</sub> (7.28 g) and the lightest (4.41 g) by T<sub>6</sub>, respectively. Our findings proved the positive increasing effect of chilling on increasing fruit weight as compared to the rest of the treatments and the reason was the efficient supply of food reserves from source to sink. Our results further found the negative effect of gibberellins on fruit weight as described by other scientists (Sharma and Singh, 2009), tomatoes, kiwifruit (Burge *et al.*, 1990) and globe artichoke (Kasim *et al.*, 2007). GA<sub>3</sub> application reduced the time needed for inflorescence emergence, induced flowering and maximized the number of flower buds and open flowers in most growing conditions. GA<sub>3</sub> at (50 mg/l) did not affect total marketable yield, while at 200 mg/l, combined with long photoperiod, enhanced the percentage of aborted flowers plus malformed fruits, resulting in a significant reduction in total marketable yield (Paroussi *et al.*, 2002).


**Fruit Firmness (pound):** The data related to fruit firmness is presented in Figure 2a which explains that the firmest fruits (0.82 pounds) were produced by T<sub>3</sub> followed by T<sub>6</sub> (0.79

pounds) and the other treatments contributed to the fruit firmness were as T<sub>5</sub> (0.75 pounds), T<sub>4</sub> (0.62 pounds), T<sub>2</sub> (0.60 pounds) and the lowest was noted in T<sub>1</sub> (0.56 pounds). Our research study found that gibberellic acid helped to maintain fruit firmness than chilling and control and this is because it regulates the activities of cell wall hydraulic enzymes which are responsible for fruit softening (Kondo and Danjo, 2001). These research findings are in agreement with the work of Canli and Orhan (2009), Cline and Trought (2007), Ozkaya and Dundar (2008). Application of 10, 20 and 30 ppm foliar sprays of gibberellic acid on sweet cherry can reduce the loss of fruit firmness (Ozkaya and Dundar, 2008). Sweet cherry trees were treated with gibberellic acid at the concentrations of 0, 15 and 25 ppm to investigate that which concentration is optimum for fruit quality. The results indicated that when gibberellic acid is sprayed at optimum concentration of 25 ppm at two different locations, caused 38% and 25% more fruit firmness than control (Canli and Orhan, 2009). The applications on sweet cherries revealed that foliar spray of gibberellic acid either applied repeatedly or singly, increased fruit cracking and fruit firmness but delayed the color development of fruits (Cline and Trought, 2007).




**Figure 1. Effect of chilling and growth hormone on strawberry (a) fruit size (b) fruit weight (under tunnel conditions).**

**Total soluble solids (TSS °Brix):** The statistical data presented in Figure 2b describes that maximum total soluble solids were produced by treatment with T<sub>3</sub> (10.80 °Brix) followed by T<sub>2</sub> (9.90 °Brix) statistically at par. The rest of the treatments contributed TSS as T<sub>5</sub> (8.66 °Brix), T<sub>6</sub> (7.96 °Brix), T<sub>1</sub> (6.56 °Brix) control, T<sub>4</sub> (6.50 °Brix), respectively. Gibberellic acid increase TSS in strawberry as compared with chilling and control treatments. These findings are similar with the work of Kappel and McDonald (2007) that GA<sub>3</sub> treated sweet cherries showed more TSS level as compared to control and chilling treatments and the maximum total soluble solids (TSS) was obtained with GA<sub>3</sub> application (Singh and Singh, 2006) and also the fruit were larger, firmer and had more soluble solids as compared to untreated plants (Facteau *et al.*, 1985).



**Figure 2. Effect of chilling and growth hormone on strawberry (a) firmness and (b) total soluble solids (under tunnel conditions).**



**Figure 3. Effect of chilling and growth hormone on strawberry (a) ascorbic acid contents and (b) titratable acidity (under tunnel conditions).**

**L-ascorbic acid (mg/100ml of juice):** L-ascorbic acid of 93.03 mg/100ml was produced by T<sub>3</sub> followed by T<sub>2</sub> (89.90 mg/100ml) and the rest of the treatments contributed as T<sub>6</sub> (88.56 mg/100ml) T<sub>4</sub> (86.90 mg/100ml), T<sub>1</sub> (87.00 mg/100ml) and 85.80 mg/100ml by T<sub>5</sub> (Fig. 3a). Strawberries are the vital source of ascorbic acid contents for the human diet (Food and Nutrition Board, 1989) and they contain more ascorbic acid as compared to oranges (Ayub *et al.*, 2010). Vitamin C is a main quality parameter which is most susceptible to degradation because of oxidation compared to the rest of nutrients during food processing and storage our results revealed the increasing effect of GA<sub>3</sub> application on vitamin C and these findings are in parallel

with that of other scientists including Sharma and Singh (2009) reported high ascorbic acid quantity in strawberry fruit plant treated with GA<sub>3</sub> during pre-flowering stage, Kappel and Macdonald (2007) suggested that GA<sub>3</sub> at 90 ppm contributed to increase the production of quality parameters like ascorbic acid and acidity of strawberry and Ouzounidou *et al.* (2010) also observed similar results by working on capsicum.

**Titrateable acidity:** The sugars and acids are used as vital measure of taste attribute for strawberries (Wozniak *et al.*, 1997) which attracts the consumers. As described in (Fig. 3b) the most significant results were observed in treatment with T<sub>3</sub> (0.69) followed by T<sub>5</sub> (0.55), T<sub>2</sub> (0.00), T<sub>6</sub> (0.50), T<sub>4</sub> (0.20) and the lowest of 0.17 by T<sub>1</sub> (control). Titrateable acidity is associated with the levels of organic acids present in the fruits as these are significant parameters for maintaining fruit quality. In strawberries citric acid is the chief acid adding to 90% of entire organic acid content. Our work revealed that gibberellins had positive effect on elevation of titrateable acidity which is undesirable with respect to consumption. These results coincide with findings of Ouzounidou *et al.* (2010) who worked on effect of gibberellic acid in capsicum plant and found non-significant results regarding titrateable acidity. El-Otmani and Coggins Jr (1991) also confirmed our results regarding effect of gibberellic acid and similar results were reported by Kappel and Macdonald (2007), they intimated that yield was not affected by GA<sub>3</sub> spray whereas early GA<sub>3</sub> spray produces firmer fruits. Total soluble solids in GA<sub>3</sub> treated strawberry plants were higher than control. Moreover, by delaying the time of spray titrateable acidity was increased in gibberellic acid treated fruits.

It is concluded from the above mentioned results that gibberellic acid though induced early flowering but decreased marketable yield by producing malformed fruits either alone or in combination with chilling while chilling alone produced maximum quality yield by increasing fruit size weight and average number of fruits. Gibberellic acid concentration of 400 ppm improved quality parameters in term of TSS, firmness, titrateable acidity and ascorbic acid so are recommended for processed fruit products.

## REFERENCES

- Agusti, M., F., Garcia-Mari and J.L. Guardiola. 1982. Gibberellic acid and fruit set in sweet orange. *Sci. Hortic.* 7:257-264.
- Ahmed, H.F.S. and M.I. Ragab. 2005. Hormone levels and protein patterns in dormant and non-dormant buds of strawberry, and induction of bud break by gibberellic acid. *Egypt. J. Biol.* 5:35-42.
- Anderson, M.K. 2006. Beach strawberry *Fragaria chiloensis* (L.) P. Mill. Spp. Lucida (Vilm) Staudt. Newsletter USDA, NRCS: National Plant Data Center.

- AOAC. 1990. Official Methods of Analysis, 15<sup>th</sup> Ed. Association of Analytical Chemists. Virginia, Arlington, USA.
- Ayub, M., J. Ullah, A. Muhammad and A. Zeb. 2010. Evaluation of strawberry juice preserved with chemical preservatives at refrigeration temperature. *Int. J. Nut. Metabol.* 2:27-32.
- Batley, N.H., P. Le Miere, A. Tehranifar, C. Cekic, S. Taylor, K.J. Shives and M.J. Wilkinson. 1998. Genetic and Environmental Control of Flowering in Strawberry. Wallingford, CAB International Press.
- Burge, G.K., C.B. Spence and N.D. Broadbent. 1990. Effects of gibberellic acid and paclobutrazol on fruit size, shape, locule number and pedicel length of kiwifruit. *Sci. Hortic.* 42:243-249.
- Canli, F.A. and H. Orhan. 2009. Effects of preharvest gibberellic acid applications on fruit quality of '0900 Ziraat' sweet cherry. *HortTechnology* 19:127-129.
- Cline, J.A. and M. Trought. 2007. Effect of gibberellic acid on fruit cracking and quality of Bing and Sam sweet cherries. *Can. J. Plant. Sci.* 87:545-550.
- Durner, E.F., E.B., Poling and L. Maas. 2000. Recent advances in strawberry plug transplant technology. *HortTechnology* 12:545-550.
- El-Otmani, M. and C.W. Coggins Jr. 1991. Growth regulator effects on retention of quality of stored citrus fruits. *Sci. Hortic.* 45:261-272.
- El-Shabasi, M.S.S., M.E. Ragab, I.I., El-Oksh and Y.M.M. Osman. 2009. Response of strawberry plants to some growth regulators. *Acta Hort.* 842:725-728.
- Facteau, T.J., K.E. Rowe and N.E. Chestnut. 1985. Response patterns of gibberellic acid-treated sweet cherry fruit at different soluble solids levels and leaf:fruit ratios. *Sci. Hortic.* 27:257-262.
- Fishman, S., A. Erez and G.A. Couvillon. 1987. The temperature dependence of dormancy breaking in plants: Mathematical analysis of a two-step model involving a cooperative transition. *J. Theor. Biol.* 124:473-483.
- Food and Nutrition Board. 1989. National Research Council. Recommended Dietary Allowances, 10<sup>th</sup> Ed. Washington, National Academy Press.
- Hannum, S.M. 2004. Potential impact of strawberries on human health: A review of the science. *Crit. Rev. Food. Sci. Nutr.* 44:1-17.
- Kappel, F. and R. Macdonald. 2007. Early gibberellic acid spray increase fruiting and fruit size of sweetheart sweet cherry. *J. Am. Pomol. Soc.* 61:38-42.
- Kasim, A.T.M., A.M. Abd-El-Hameid and N.H.M. El-Greadly. 2007. A comparison study on the effect of some treatment on earliness, yield and quality of Globe Artichoke (*Cynara scolymus* L.). *Res. J. Agric. Biol. Sci.* 3:695-700.
- Kondo, S. and C. Danjo. 2001. Cell wall polysaccharide metabolism during fruit development in sweet cherry 'Satoh-nishiki' as affected by gibberellic acid. *J. Jpn. Soc. Hort. Sci.* 70:178-184.
- Lieten, F., J.M. Kinet and G. Bernier. 1995. Effect of prolonged cold storage on the production capacity of strawberry plants. *Sci. Hortic.* 60:213-219.
- Luedeling, E., J. Gebauer and A. Buerkert. 2009. Climate change effects on winter chill for tree crops with chilling requirements on the Arabian Peninsula. *Clim. Chang.* 96:219-237.
- Luedeling, E., M.H. Zhang, V., Luedeling and E.H. Girvetz. 2010. Sensitivity of winter chill models for fruit and nut trees to climatic changes expected in California's Central Valley. *Agric. Ecol. Environ.* 131:23-31.
- Ouzounidou, G., I. Ilias, A. Giannakoula and P. Papadopoulou. 2010. Comparative study on the effects of various plant growth regulators on growth, quality and physiology of *Capsicum annuum*. *Pak. J. Bot.* 42:805-814.
- Ozguven, A.I. and C. Yilmaz. 2002. The effect of gibberellic acid treatments on the yield and fruit quality of strawberry (*Fragaria x ananassa*) cv. Camarosa. *Acta Hort.* 567:277-280.
- Ozkaya, O. and O. Dundar. 2008. Chemical and physical determination of gibberellic acid effects on postharvest quality of sweet cherry. *Asian J. Chem.* 20:751-756.
- Paroussi, G., D.G. Voyiatzis, E. Paroussis and P.D. Drogoudi. 2002. Effect of GA<sub>3</sub> and photoperiod regime on growth and flowering in strawberry. *Acta Hort.* 567:273-276.
- Paroussi, G., D.G. Voyiatzis, E. Paroussis and P.D. Drogoudi. 2002. Growth, flowering and yield responses to GA(3) of strawberry grown under different environmental conditions. *Sci. Hortic.* 96:103-113.
- Risser, G. and F. Robert. 1993. What cold treatments promote growth in strawberry. *Acta Hort.* 48:381-383.
- Sharma, R.R. and R. Singh. 2009. Gibberellic acid influences the production of malformed and button berries, and fruit yield and quality in strawberry (*Fragaria x ananassa* Duch.). *Sci. Hortic.* 119:430-433.
- Singh, A. and J.N. Singh. 2006. Studies on influence of biofertilizers and bioregulators on flowering, yield and fruit quality of strawberry cv. sweet Charlie. *Ann. Agric. Res.* 27:261-264.
- Steel, R.G.D., J.H. Torrie and M.A. Boston. 1997. Principles and procedures of statistics: A biometrical approach, 3<sup>rd</sup> Ed. McGraw Hill Book Company Inc., New York, USA.
- Sung, S. and R.M. Amasino. 2004. Vernalization and epigenetics: how plants remember winter. *Curr. Opin. Plant Biol.* 7:4-10.
- Taylor, D.R. 2002. The physiology of flowering in strawberry. *Acta Hort.* 567:245-251.
- Usenik, V., D. Kastelec and F. Štampar. 2005. Physicochemical changes of sweet cherry fruits related

to application of gibberellic acid. Food Chem. 90:663-671.

Wozniak, W., B. Radajewska, A. Reszelska-Sieciechowicz and I. Dejwor. 1997. Sugars and acid content influence organoleptic evaluation of fruits of ix strawberry cultivars from controlled cultivation. Acta Hort. 439:333-336.