

Mechanisms of Physical Activity Behavior Change for Prostate Cancer Survivors: A Cluster Randomized Controlled Trial

Melinda J. Craike, PhD¹ • Cadeyrn J. Gaskin, PhD² • Mohammadreza Mohebbi, PhD² • Kerry S. Courneya, PhD³ • Patricia M. Livingston, PhD⁴

Published online: 1 February 2018

© The Author(s) 2018. Published by Oxford University Press on behalf of the Society of Behavioral Medicine.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Abstract

Background Exercise is beneficial for prostate cancer survivors. Therefore, understanding the mechanisms of physical activity (PA) behavior change is imperative.

Purpose The ENGAGE study was an exercise intervention for prostate cancer survivors, which improved vigorous physical activity (VPA) at postintervention and follow-up. The purpose of this study was to assess (a) whether the intervention improved social cognitive determinants of behavior and (b) the extent to which social cognitive determinants mediated the effect of the exercise intervention on VPA.

Methods Overall, 147 men consented to be involved in the study (intervention = 54, usual care = 93). Data from baseline, postintervention (12 weeks) and follow-up (6 months) were used in this analysis. Social cognitive determinants were measured using appropriate measures. VPA was measured using an adapted version of the Leisure-Time Exercise Questionnaire.

Results Compared with the control condition, men in the intervention condition had higher *task self-efficacy* postintervention (+16.23; 95% confidence interval [CI] +9.19 to +23.31; effect size [d] = 0.85, $p < .001$) and at follow-up (+12.58; 95% CI = +4.45 to +20.71, $d = 0.50$, $p = .002$). Task self-efficacy partially mediated the effect of the exercise intervention on VPA (indirect effect: $B = 19.90$; 95% CI 1.56 to 38.25, $p = .033$).

Conclusion The intervention improved the belief among prostate cancer survivors that they could perform challenging exercises for longer periods of time, which partially explained the positive effect of the intervention on VPA.

Australia and New Zealand Clinical Trials Registration ACTRN12610000609055.

Keywords Cancer • Oncology • Physical activity • Social cognitive theory

Introduction

With earlier detection and more effective treatments, more men with prostate cancer are living longer postdiagnosis. For example, in Australia, 5 year relative survival rates increased to 93.0% for the years 2007–2011 [1]. Prostate cancer survivors, however, experience significant physical and psychological morbidity following their diagnosis and subsequent treatment [2, 3]. Physical activity (PA) has a role in alleviating this morbidity [4, 5], but engagement in PA is markedly lower than what is recommended [3, 6]. A recent study of 463 prostate cancer survivors, for

✉ Melinda Craike
Melinda.craike@vu.edu.au

¹ Institute of Sport, Exercise and Active Living (ISEAL), Victoria University, Footscray, Australia

² Faculty of Health, Biostatistical Unit, Deakin University, Geelong, Australia

³ Faculty of Physical Education and Recreation, University of Alberta, Edmonton, Canada

⁴ Faculty of Health, School of Nursing & Midwifery, Deakin University, Geelong, Australia

example, revealed that only 12.3% were sufficiently active [6]. To improve participation in PA among this population, theory-based behavior change interventions must be designed [7] to target the most pertinent mechanisms that will facilitate change in behavior [8].

Behavior change interventions that aim to support PA participation in prostate cancer survivors are scarce. In one study, researchers examined the effect of a home-based intervention on PA for prostate cancer survivors, which utilized implementation intentions [9], and reported an increase in PA participation at 1 month but not at the 3 month follow-up [10]. Furthermore, the mediators of behavior change were not examined [10]. To date, most trials among prostate cancer survivors have focused on the efficacy of exercise programs for improving health outcomes during and following treatment [4, 5, 11]. Consequently, there is now a substantial body of evidence that supports the positive effect of exercise for improving incontinence, fitness, fatigue, body composition, muscular endurance, aerobic endurance, and quality of life among prostate cancer survivors [4, 5, 11]. For these men to achieve the health benefits of exercise, research must also focus on interventions to support them to participate in PA and explore the mechanisms of behavior change.

The use of theories to guide behavior change interventions is important because theory provides a generalizable framework, facilitates an understanding why interventions do, or do not, work, and assists the accumulation of knowledge [8]. Social cognitive theory (SCT) [12, 13] is well suited to exercise programs that aim to change behavior [14]. According to SCT, the main determinants of behavior are self-efficacy, outcome expectations, goals, and sociostructural factors [13, 14]. SCT-based exercise interventions have shown promise in increasing PA participation for cancer survivors [15–18]. There are several limitations of existing research in this area, however. Few researchers have conducted mediation analyses or reported changes in SCT determinants, which has resulted in limited understanding how interventions affect behavior change; follow-up PA is rarely reported; thus, there is uncertainty about the longer term impact of exercise interventions; and finally, researchers have focused on home-based interventions and breast cancer survivors, which limit the generalizability of findings of SCT-based PA interventions to other types of intervention delivery modes (such as supervised or combined supervised and home-based interventions) and other cancer populations [18].

In the ENGAGE study, we examined the efficacy of a clinician-referred 12 week exercise intervention for prostate cancer survivors. The primary outcome was self-reported minutes of moderate-vigorous PA per week, and the main outcomes of the study have been previously reported [19]. In terms of PA participation, the intervention, compared with usual care, did not significantly improve minutes of moderate-vigorous PA per week. However, the intervention

significantly increased vigorous physical activity (VPA) and the proportion of men meeting PA guidelines postintervention (12 weeks) [19]. The effect of the intervention on VPA was maintained at follow-up (6 months); men in the intervention condition participated in 56 minutes (95% CI 14.2, 97.5) per week more VPA than those in the control condition. However, the difference in VPA at 12 months was not significant [20]. This paper addresses secondary aims of the ENGAGE study: (a) to assess whether the exercise intervention improved social cognitive determinants of PA behavior (i.e., self-efficacy, outcome expectations, goals, and sociostructural factors) between baseline and postintervention (12 weeks) and to see if these changes were maintained at follow-up (6 months) and (b) to assess the extent to which social cognitive determinants mediated the effect of the intervention on VPA [21]. Because the intervention did not have a significant effect on the primary outcome of moderate-vigorous PA, but an effect on VPA was found, we used minutes of VPA at follow-up as our outcome.

Methods

This trial was registered with the Australia and New Zealand Clinical Trials Register (ACTRN12610000609055).

Design and Procedures

Details of the ENGAGE study methods have been reported elsewhere [21] and are briefly presented here. The study was a two-arm multicenter, cluster randomized controlled trial. The eligibility criteria for inclusion in this study were as follows: completed active treatment for prostate cancer within the previous 3–12 months (patients on hormone treatment were eligible to participate), treated with curative intent, and the ability to complete surveys in the English language. The exclusion criteria for this study included musculoskeletal, cardiovascular, or neurological disorders that could limit exercising. The patients' treating clinicians provided medical clearance prior to involvement in the study.

Patients were identified through urology and radiation oncology outpatient clinics across three public health services and four private clinics located in Melbourne, Australia. Recruitment occurred from October 2011 to June 2013. Patients were approached and provided with information about the study prior to their consultation with their treating clinicians. During the consultation, clinicians randomized to the intervention condition determined the suitability of the patient for the intervention, provided a blue referral slip, and recommended that the patient be involved in the exercise program. Clinicians randomized to the control condition provided usual care regarding PA advice (typically, minimal information) and did not provide a referral to the exercise program. A study researcher later phoned eligible patients to gain informed consent.

Sample size calculations were based on the primary outcome of the main study (self-reported participation in moderate–vigorous PA) [19, 21], rather than the secondary analysis reported here.

Ethical approval was obtained from the university and each of the health services involved in the study. All participants provided written informed consent.

Intervention

The 12-week exercise program was based on PA guidelines for cancer survivors [22, 23], which recommend 150 minutes per week of moderate–vigorous PA. Participants were asked to complete three, 50 minute sessions of PA per week. Two of these sessions were supervised sessions in community gyms and one session was an unsupervised, home-based session. Sessions were predominantly one-on-one sessions, with men occasionally training in pairs.

The social cognitive determinants of behavior informed our design of the intervention [14, 24]. The 12-week exercise program was intended to increase PA through, in part, enhancing exercise-related self-efficacy in prostate cancer survivors and assisting them to develop positive outcome expectations, set beneficial goals, and reduce perceived impediments to exercise participation. Our previous research suggested that the involvement of clinicians in referring men to the program improved men's confidence in performing PA [25], which might increase self-efficacy through increasing men's belief in their ability to undertake exercise. Exercise trainers, who conducted the gym sessions, were trained in the principles of SCT and their application to PA uptake and adherence for prostate cancer survivors through an induction session prior to their commencement in the project. Exercise trainers were postgraduate exercise physiology students and were supervised by an accredited exercise physiologist. In the first exercise session, exercise trainers were instructed to encourage participants to set short-term and longer-term SMART (i.e. specific, measurable, attainable, realistic, timely) exercise goals; incorporate discussions of the participant's beliefs in their ability to exercise, benefits of exercise, outcome expectations from exercise, factors that facilitate their participation and how to maximize these; and identify potential barriers and strategies for overcoming them. Exercise trainers were instructed to revisit these discussions periodically throughout the training sessions and regularly counseled participants throughout the 12-week program. The topics for these discussions were based on the main concepts of the SCT [14, 24]. Exercise trainers were provided with checklists to ensure that they were regularly addressing the key topics, including setting goals, aligning activity with outcome expectations, overcoming barriers, and drawing on facilitators, during exercise sessions over the 12-week period.

For the unsupervised sessions, men were provided with a home-based PA program instruction sheet, instructions on how to perform the home-based exercises, and a Thera-Band to complete the exercises. These home-based sessions were designed to increase adherence to PA following completion of the intervention for those who preferred to exercise at home. To facilitate adherence to PA following the program, participants were also offered an exercise program and discounted gym membership.

The intervention included supervised and home-based PA. The supervised component was important because participants had regular contact and encouragement from exercise trainers and could be directly observed and instructed on correct exercise technique, which reduces the risk of injury and may also improve adherence. Home-based activity might increase adherence to PA because it is cost-effective for participants and PA can be incorporated into daily routines [26]. Men in the control condition were not referred to an exercise program and received usual care, which typically does not include recommendations for PA.

Random Assignment

Clinicians were invited to participate in the study and were randomly allocated using an online random number generator to either the intervention or control condition.

Measures

Demographic and clinical characteristics were assessed at baseline. Data presented in this analysis are based on PA behavior and the social cognitive determinants assessed at baseline (T1), postintervention (12 weeks, T2), and at follow-up, 3 months after the completion of the intervention (6 months, T3).

Demographic and clinical characteristics

Date of birth, level of education, and type of treatment (surgery, radiotherapy and hormone therapy) were collected through self-report. Weeks since treatment completion and cancer stage were collected through medical records.

Primary outcome

An adapted version of the Leisure-Time Exercise Questionnaire [27] was used to measure PA participation. In the Leisure-Time Exercise Questionnaire, participants report the average weekly frequency of strenuous, moderate, and light exercise over the past month. We asked participants to report the average duration of time they spend exercising at each intensity, in addition to the frequency during an average week in the past month [15, 27] and we also removed examples of physical activities that were not common in Australia. The Leisure-Time

Exercise Questionnaire has been used extensively with cancer survivor groups [27, 28]. In the current analyses, we used data on minutes per week of strenuous PA as our outcome (i.e., VPA).

Mediators

Outcome expectations

Outcome expectations were measured using the Multidimensional Outcome Expectations for Exercise Scale (MOEES) [29]. This scale was designed for use in older adults and comprises 15 items that represent three subscales: physical (e.g., exercise will strengthen my bones), social (e.g., exercise will improve my social standing), and self-evaluative (e.g., exercise will give me a sense of personal accomplishment) outcome expectations. Participants respond to each item on a 5-point Likert scale ranging from 1 (strongly disagree) to 5 (strongly agree). The instrument's construct validity has been demonstrated through confirmatory factor analysis and significant correlations with other constructs consistent with the predictions inherent in SCT [29, 30]. We used the total outcome expectations scale as well as the three subscales in our analyses. In the current sample, Cronbach's alpha for the total MOEES scale and subscales were high: total outcome expectations (0.92), physical (0.87), social (0.84), and self-evaluative (0.87).

Self-efficacy

Both task and *barrier self-efficacy* were measured using methods Bandura [24, 31] suggested. Task self-efficacy was assessed using nine items, asking participants to rate how confident they are, ranging from 0% (not at all confident) to 100% (extremely confident), that they can perform three different types of PA (walking fast, running fast, swimming or cycling hard, and doing exercises with weights) for three time durations (10, 20, and 30 minutes). Similar items have been included in previous PA and cancer research in which task self-efficacy was measured [32]. Higher scores indicate higher levels of task self-efficacy. In the current sample, Cronbach's alpha for the task self-efficacy items was high (0.92). Barrier self-efficacy was assessed using 25 items, asking participants to rate how confident they are, ranging from 0% (not at all confident) to 100% (extremely confident), that they can perform PA when faced with specific barriers. Examples of barriers included, "when I am feeling tired" and "when there are other interesting things to do." The items were based on Bandura's [31] self-efficacy to regulate exercise instrument, with supplementary items from research on PA for prostate cancer survivors [25] and cancer patients [32]. Higher scores indicate higher levels of barrier self-efficacy. In the current sample, Cronbach's alpha for the barrier self-efficacy items was high (0.95).

Sociostructural factors

Sociostructural factors facilitating (operationalized as motivation) or impeding (operationalized as barriers) participation in PA were measured using guidelines for the development of scales to assess barriers to PA for cancer patients [33]. The content of these scales was based on previous research on the perceived benefits and barriers to PA for prostate cancer survivors [25]. Participants are asked their perceptions of the frequency with which motivations (10 items) and barriers (12 items) occur using a 5-point Likert scale anchored with 0 (never) and 4 (always). Examples of motivation items include "socializing with other people" and "increasing my energy levels." Examples of barrier items include "injuries (e.g. muscular and joint problems)" and "lacking time to do exercise." In the current sample, Cronbach's alphas for the barrier and motivation items were high (0.83 and 0.92, respectively).

Goals

Goals were measured based on items from previous PA research that assessed SCT constructs [34]. Participants are asked to report the number of days per week they intended to perform at least 30 minutes of exercise at a light, moderate, or vigorous intensity and were asked about current goals and their 12 week exercise goals (2 items).

Statistical Approach

We used STATA version 14 to analyze data. There was minimal missing data for the variables included in this analysis (overall <2% values missing). We calculated the mean score for MOEES, exercise motivation, exercise barriers, task self-efficacy, and barrier self-efficacy if at least 70% of the items were answered. Two items measured exercise goals; a score was calculated if at least one of these was answered. When less than 70% of responses were available for a scale, or where all items were reported as "not applicable," data were treated as missing. For longitudinal analyses, we accounted for missing data using the generalized estimating equations (GEE) approach, and we eliminated missing data from other analyses ([35]).

Our process for testing the possible mediation of the intervention effect on PA was guided by Baron and Kenny's [36] principles. As others have argued [37], however, the structural equation modeling (SEM) approach is more appropriate than Baron and Kenny's [36] approach because it estimates all paths simultaneously instead of assuming that equations 1 to 3 are independent. As such, the SEM approach was used for examining mediation. We only tested social cognitive determinants as possible mediators whether they were affected by the intervention (i.e., condition assignment) and associated with follow-up VPA. We assessed the impact of the intervention on social cognitive determinants between

baseline and postintervention (12 weeks) and baseline and follow-up (6 months) using time-by-treatment interactions in a repeated measures split-plot in time ANOVA model for continuous outcomes. We estimated model parameters using GEE with an exchangeable working correlation matrix to take account of the repeated measures for each participant. We also conducted post hoc contrasts to determine follow-up-by-intervention impact. We calculated effect sizes (follow-up-by-intervention effect) using Cohen's d . We conducted a series of regression analyses using change scores to investigate the extent to which changes in the social cognitive determinants of behavior postintervention (12 weeks) and follow-up (6 months) explained the increased VPA at follow-up (6 months). We adjusted for baseline VPA and the baseline data of the SCT determinants for each analysis. We also reported eta-squared effect sizes for these analyses.

Finally, we used path regression analyses using the SEM approach to test for possible mediation of the intervention effect on VPA. We controlled for both baseline social cognitive determinants and VPA in the path analysis. That is, we considered the baseline data of potential mediators (i.e. social cognitive determinants) effect modifiers for postintervention and follow-up mediator data as well as baseline and follow-up VPA. We also considered baseline VPA as an effect modifier for follow-up VPA. The autoregressive SEM model [38] was used for this purpose. We report standardized coefficients to aid in the interpretation of the mediation effects. Mediation is present when the proposed mediator maintains a significant relationship with the outcome, whereas condition does not. We also reported the controlled direct effect and natural indirect effect of mediators. We did not adjust for multiple comparisons because our mediation analyses were performed by implementing SEMs where all pathways, including the mediation effects, were assessed simultaneously. As this study was a cluster randomized controlled trial, we examined the effect of clustering by clinician for the social cognitive determinants and VPA by calculating the intra-cluster correlation coefficient (ICC) for the mean difference between each social cognitive determinant. Each ICC was less than 0.05, except task self-efficacy (ICC = 0.198) and follow-up VPA (ICC = 0.31). Therefore, we evaluated the clustering effect of clinician in the path analysis in a sensitivity analysis by including clinician as a fixed effect modifier factor.

Results

Study Flow and Participant Characteristics

The flow of participants and a detailed description of the demographic and clinical characteristics of the sample have been reported previously [19]. Overall, 147 (46%; 54

intervention and 93 control) eligible patients consented to participate in the study. Participants ranged in age from 39 to 84 years ($M = 65.6$; $SD = 8.5$); 27.2% had a university degree; the majority had stage I or II disease (82.8%); and 43.5% were treated with surgery only, with 21.1% treated with surgery and radiotherapy.

We obtained postintervention (T2) data from 88.4% of participants ($n = 130$; 47 intervention, 83 control). There was no significant difference in attrition rates between the intervention and control conditions (control, 11%; intervention, 13%; $p = .70$). We obtained 6 month follow-up (T3) data from 121 (83.3%; 43 intervention, 78 control) participants. There was no significant difference in attrition rates between the intervention and control groups at T3 (control, 16.3%; intervention, 20.5%; $p = .23$).

As previously reported [19], 85% of participants adhered to at least 18 of the 24 supervised sessions. For participants who returned their home exercise diary (74%), 81% completed at least 9 of the 12 of the prescribed home-based weekly sessions.

Effect of the Exercise Intervention on Social Cognitive Determinants Postintervention (T2, 12 weeks)

Descriptive statistics and social cognitive determinant change scores immediately postintervention (12 weeks) across the two conditions are presented in Table 1. Compared with the control condition, participants in the intervention condition had significantly higher task self-efficacy with a large effect size ($+16.23$; 95% CI = $+9.19$ to $+23.31$, $d = 0.85$, $p < .001$). There were no significant differences between the intervention and control conditions for any other social cognitive determinant, and effect sizes were small (all less than $d = 0.30$).

Effect of the exercise intervention on social cognitive determinants at follow-up (T3, 6 months)

Descriptive statistics and social cognitive determinant change scores at follow-up (6 months) across the two conditions are presented in Table 2. Compared with the control condition, participants in the intervention condition had higher task self-efficacy at follow-up, with a moderate effect size ($+12.58$; 95% CI = $+4.45$ to $+20.71$, $d = 0.50$; $p = .002$). There were no significant differences between the two conditions for any other social cognitive determinants at follow-up.

Association between Social Cognitive Determinants and Follow-up VPA

Associations between social cognitive determinants at postintervention (T2, 12 weeks) and VPA at follow-up (T3, 6 months) showed that increases in barrier self-efficacy ($B = 1.61$, 95% CI = $+0.57$ to $+2.65$, $\eta^2 = 0.08$,

Table 1 Effects of the Intervention on Social Cognitive Determinants at Postintervention (T2; $N = 127$)

	Baseline	Postintervention	Mean change	Between-group comparison	d	p
	M (SD)	M (SD)	M (95% CI)	M (95% CI)		
Outcome expectations (range 1–5)						
Int	4.02 (0.46)	4.00 (0.45)	−0.02 (−0.17 to +0.08)	+0.01 (−0.14 to +0.16)	0.02	.89
Cont	3.87 (0.60)	3.83 (0.62)	−0.03 (−0.13 to +0.06)			
Outcome expectations—physical (range 1–5)						
Int	4.43 (0.39)	4.45 (0.42)	+0.02 (−0.10 to +0.15)	+0.01 (−0.15 to +0.17)	0.02	.91
Cont	4.33 (0.58)	4.34 (0.55)	+0.02 (−0.09 to +0.12)			
Outcome expectations—social (range 1–5)						
Int	3.27 (0.77)	3.21 (0.70)	−0.06 (−0.21 to +0.09)	+0.06 (−0.16 to +0.30)	0.10	.59
Cont	3.06 (0.87)	2.94 (0.90)	−0.12 (−0.27 to +0.02)			
Outcome expectations—self-evaluative (range 1–5)						
Int	4.13 (0.54)	4.09 (0.535)	−0.05 (−0.18 to +0.08)	−0.03 (−0.23 to +0.18)	−0.06	.78
Cont	3.97 (0.67)	3.94 (0.81)	−0.02 (−0.15 to +0.11)			
Task self-efficacy (range 0–100)						
Int	53.00 (28.07)	72.31 (25.60)	+18.17 (+11.50 to +24.84)	+16.23 (+9.19 to +23.31)	0.85	<.001
Cont	56.81 (25.41)	59.15(26.86)	+1.70 (−2.04 to +5.44)			
Barrier self-efficacy (range 0–100)						
Int	63.81 (19.25)	70.22 (19.41)	+7.09 (+1.29 to +12.90)	+5.33 (−2.17 to +12.82)	0.25	.16
Cont	58.16 (23.84)	58.96 (26.23)	+2.10 (−2.51 to +6.71)			
Barriers (range 0–4)						
Int	0.46 (0.43)	0.44 (0.43)	−0.03 (−0.17 to +0.12)	+0.05 (−0.13 to +0.23)	0.08	.61
Cont	0.63 (0.585)	0.55 (0.57)	−0.07 (−0.18 to +0.03)			
Motivation (range 0–4)						
Int	2.44 (1.19)	2.11 (1.24)	−0.33 (−0.66 to 0.00)	−0.01 (−0.41 to −0.39)	−0.08	.98
Cont	2.06 (1.00)	1.66 (1.09)	−0.32 (−0.56 to −0.09)			
Goals (range 1–9)						
Int	6.21 (1.72)	6.26 (1.83)	+0.04 (−0.54 to +0.61)	+0.38 (−0.34 to +1.10)	0.17	.30
Cont	5.72 (2.48)	5.32 (2.56)	−0.34 (−0.77 to +0.09)			

Baseline and postintervention means and standard deviations are raw data; Higher scores for each variable represent higher levels of that variable. Mean change and between-group comparison are based on model parameters using generalized estimating equations (GEE). At T2, 130 questionnaires were received with baseline data missing for 3 participants. The GEE approach was used for between-group comparisons to account for baseline missing data.

Int intervention condition; Cont control condition; M mean; SD standard deviation; CI confidence interval; d Cohen's d effect size.

$p = .003$) and reductions in motivation ($B = -21.19$, 95% CI = -41.60 to -0.78 , $\eta^2 = 0.04$, $p = .04$) predicted improved participation in follow-up VPA (Table 3).

Associations between social cognitive determinants at follow-up (T3, 6 months) and VPA at follow-up (T3, 6 months) showed that increases in outcome expectations—social ($B = 39.92$, 95% CI = $+8.99$ to $+70.86$, $\eta^2 = 0.06$, $p = .01$), task self-efficacy ($B = 1.48$, 95% CI = $+0.45$ to $+2.50$, $\eta^2 = 0.07$, $p = .005$), barrier self-efficacy ($B = 1.72$, 95% CI = $+0.79$ to $+2.66$, $\eta^2 = 0.11$, $p = <.001$), and goal setting ($B = 11.29$, 95% CI = $+1.45$ to $+21.13$, $\eta^2 = 0.05$, $p = .03$)—predicted improved participation in follow-up VPA.

Mediation Analysis

The intervention had a positive effect on task self-efficacy at follow-up, and task self-efficacy at follow-up was associated with VPA at follow-up. Task self-efficacy was the only social cognitive determinant that met the criteria for the mediation analysis. The path analysis is shown in Fig. 1. The association between the exercise intervention (condition) and follow-up VPA was only slightly reduced from $\beta = 0.25$ ($p = .001$) to $\beta = 0.18$ ($p = .01$) and was still significant after adjusting for follow-up task self-efficacy. The path from follow-up task self-efficacy remained significantly associated with follow-up VPA ($\beta = 0.35$, $p = <.001$). Follow-up

Table 2 Effects of the Intervention on Social Cognitive Determinants at Follow-up (T3; *N* = 118)

	Baseline	Follow-up	Mean change	Between-group comparison	<i>d</i>	<i>p</i>
	<i>M</i> (<i>SD</i>)	<i>M</i> (<i>SD</i>)	<i>M</i> (95% <i>CI</i>)	<i>M</i> (95% <i>CI</i>)		
Outcome expectations (range 1–5)						
Int	4.03 (0.45)	4.08 (0.44)	+0.03 (−0.03 to +0.08)	+0.02 (−0.16 to +0.19)	0.06	.86
Cont	3.85 (0.56)	3.89 (0.61)	+0.02 (−0.04 to +0.08)			
Outcome expectations—physical (range 1–5)						
Int	4.45 (0.40)	4.50 (0.40)	+0.02 (−0.04 to +0.09)	+0.03 (−0.16 to +0.21)	0.04	.78
Cont	4.33 (0.54)	4.36 (0.53)	+0.01 (−0.05 to +0.07)			
Outcome expectations—social (range 1–5)						
Int	3.24 (0.71)	3.33 (0.64)	+0.04 (−0.035 to +0.12)	+0.02 (−0.25 to +0.28)	0.02	.91
Cont	3.01 (0.87)	3.06 (0.95)	+0.03 (−0.06 to +0.12)			
Outcome expectations—self-evaluative (range 1–5)						
Int	4.16 (0.54)	4.20 (0.57)	+0.02 (−0.05 to +0.09)	0.00 (−0.20 to +0.21)	−0.07	.97
Cont	3.95 (0.67)	3.99 (0.75)	+0.02 (−0.05 to +0.09)			
Task self-efficacy (range 0–100)						
Int	53.75 (27.56)	71.27 (28.51)	+8.31 (+4.41 to +12.22)	+12.58 (+4.45 to +20.71)	0.50	.002
Cont	56.95 (27.24)	60.63 (25.36)	+1.95 (−0.16 to +4.06)			
Barrier self-efficacy (range 0–100)						
Int	63.57 (19.16)	71.08 (21.49)	+3.72 (+0.40 to +7.03)	+4.39 (−3.77 to +12.55)	0.13	.29
Cont	57.75 (24.17)	61.29 (26.77)	+1.48 (−0.96 to +3.91)			
Barriers (range 0–4)						
Int	0.44 (0.43)	0.41 (0.55)	−0.02 (−0.10 to +0.07)	+0.03 (−0.20 to +0.25)	0.01	.82
Cont	0.63 (0.60)	0.56 (0.63)	−0.03 (−0.10 to +0.04)			
Motivation (range 0–4)						
Int	2.49 (1.17)	2.26 (1.33)	−0.12 (−0.27 to +0.04)	−0.06 (−0.47 to +0.36)	−0.08	.79
Cont	2.02 (1.00)	1.84 (1.06)	−0.09 (−0.22 to +0.04)			
Goals (range 1–9)						
Int	6.16 (1.70)	6.51 (1.73)	+0.17 (−0.13 to +0.48)	+0.42 (−0.45 to +1.28)	0.17	.35
Cont	5.65 (2.42)	5.56 (2.59)	−0.03 (−0.31 to +0.24)			

Baseline and postintervention means and standard deviations are raw data. Higher scores for each variable represent high levels of that variable; Mean change and between-group comparison are based on model parameters using generalized estimating equations (GEE). At T3, 121 questionnaires were received with baseline data missing for 3 participants. The GEE approach was used for between-group comparisons to account for baseline missing data.

Int intervention condition; *Cont* control condition; *M* mean; *SD* standard deviation; *CI* confidence interval; *d* Cohen's *d* effect size.

task self-efficacy was a partial mediator of the effect of the exercise intervention on follow-up VPA, as demonstrated through a significant indirect effect on VPA (natural indirect effect: $B = 19.90$; 95% *CI* 1.56 to 38.25, $p = .03$), with the intervention continuing to have a controlled direct effect ($B = 54.12$, 95% *CI* 9.64 to 98.60, $p = .02$).

Discussion

Our study is one of the few exercise interventions to focus on PA behavior change for prostate cancer survivors and also addressed several limitations of previous SCT-based exercise interventions for cancer survivors [18]. We examined an exercise intervention that incorporated both supervised and unsupervised exercise,

changes in scores on measures representing theoretical constructs of SCT, and conducted mediation analyses to examine whether changes in social cognitive determinants affected changes in VPA. We found that the exercise intervention was effective in increasing task self-efficacy postintervention, which was maintained at follow-up, and task self-efficacy partially explained the effect of the intervention on longer term VPA.

Self-efficacy is a key determinant of SCT [13], and the intervention may have achieved a positive effect on task self-efficacy in a number of ways. Bandura [13] proposed that expectations of personal efficacy are derived from four sources of information: performance accomplishments, *vicarious experience*, *verbal persuasion*, and *physiological states*. *Performance accomplishments* may

Table 3 Associations between Social Cognitive Determinants and Post Intervention Follow-up Vigorous Physical Activity (VPA)

	<i>B</i>	95% CI	η^2	<i>p</i>
Postintervention (12 weeks)				
Outcome expectations	6.29	−46.07 to +58.65	0.00	.81
Outcome expectations—physical	−20.14	−67.34 to +27.05	0.01	.40
Outcome expectations—social	23.31	−10.69 to +57.31	0.02	.18
Outcome expectations—self-evaluative	−0.94	−39.67 to +37.78	0.00	.96
Task self-efficacy	0.95	−0.21 to +2.11	0.02	.11
Barrier self-efficacy	1.61	+0.57 to +2.65	0.08	.003
Barriers	32.20	−12.67 to +77.07	0.02	.16
Motivation	−21.19	−41.60 to −0.78	0.04	.04
Goals	−9.67	−21.09 to +1.76	0.03	.10
Follow-up (6 months)				
Outcome expectations	25.50	−21.86 to +72.86	0.01	.29
Outcome expectations—physical	−9.50	−53.09 to +34.08	0.00	.67
Outcome expectations—social	39.92	+8.99 to +70.86	0.06	.01
Outcome expectations—self-evaluative	13.01	−26.45 to +52.46	0.00	.52
Task self-efficacy	1.48	+0.45 to +2.50	0.07	.005
Barrier self-efficacy	1.72	+0.79 to +2.66	0.11	<.001
Barriers	−12.16	−49.30 to +24.98	0.00	.52
Motivation	5.20	−15.27 to +25.67	0.00	.62
Goals	11.29	+1.45 to +21.13	0.05	.03

Analyses were adjusted for baseline VPA and baseline of the social cognitive determinant.




Fig. 1. Path model testing follow-up task self-efficacy as a mediator of the effects of the exercise intervention on follow-up vigorous physical activity (VPA). Model adjusted for baseline value of task self-efficacy and baseline VPA. Parentheses indicate the path from the exercise intervention to follow-up VPA without task self-efficacy in the model

have been improved through participants completing the exercises and the feedback and guidance on performing strength and aerobic exercise provided by exercise trainers. During the supervised sessions, success was encouraged through starting the program at a level achievable by the participants and building the intensity throughout the program. Vicarious experience may have been improved through observations of other gym members participating in exercise at the community-based gyms in which exercise sessions were located. Verbal persuasion was likely to be influential through the clinician's referral to the exercise program and the ongoing encouragement by exercise trainers. Finally, through participation in

the exercise program, men may have learned to interpret physiological states that accompany exercise such as increased heart rate, sweating, and muscle fatigue as signs that their body was responding normally to the exercise, rather than as a negative sign of inability to exercise or as a reason to discontinue activity. Future research could generate a better understanding of the relative importance of each of these information sources.

Direct comparisons with other interventions are limited because we could not find any other research that focused only on prostate cancer survivors, which examined change in self-efficacy. Home-based PA interventions that have included prostate cancer survivors (along

with breast cancer survivors) have shown mixed findings on their effectiveness in improving self-efficacy [17, 39]. For example, a home-based diet and PA intervention for older breast and prostate cancer survivors (43% prostate cancer), which included telephone counseling and mailed materials had a positive effect on self-efficacy for PA. However, this intervention did not improve PA participation [39]. Conversely, a mailed print intervention to improve dietary and PA practices among newly diagnosed breast and prostate cancer survivors (44% prostate cancer) had no effect on self-efficacy for PA [17]. Although each of these studies used different measures of self-efficacy, the findings are comparable because each assessed task self-efficacy, rather than barrier self-efficacy. Studies of other cancer population groups, which have assessed the impact of PA interventions on self-efficacy, also resulted in conflicting findings [40, 41]. A recent systematic review found that although PA interventions improve self-efficacy among healthy older adults, the use of self-regulatory techniques was associated with *reduced* self-efficacy [42]. Drawing conclusions about the factors that may lead to an increase of self-efficacy among prostate cancer survivors is hampered by the limited amount of research in this area; our results suggest that a combined supervised and home-based intervention can enhance task self-efficacy for this group.

We found that task self-efficacy was a partial mediator of the effect of the exercise intervention on longer term VPA. Previous researches focusing on men [43] and older adults [44] have shown that self-efficacy is a key determinant of PA participation. Examination of the mediators of PA interventions has usually focused on self-efficacy and, as previously discussed, studies that have included prostate cancer survivors either did not improve self-efficacy [17] or did not increase PA participation [39]. Several researchers have examined mediators of PA behavior change for cancer survivors using the theory of planned behavior [28, 45, 46] and showed that perceived behavioral control [28, 45] and planning and intention [46] were partial mediators of the effect of the intervention on PA behavior. Taken collectively, it appears that theoretical constructs in both SCT and theory of planned behavior only partially explain PA behavior change in cancer survivors, suggesting that other, unmeasured factors may explain the effect of such interventions on PA behavior change.

Contrary to our expectations, apart from a positive effect on task self-efficacy, the exercise intervention did not significantly improve the other social cognitive determinants assessed in our study. This is in contrast with a recent intervention aimed at increasing PA among cancer survivors and their carers that had a positive impact on behavioral goal setting and outcome expectations. This intervention included intensive sessions on goal setting and weekly monitoring of goals using a pedometer and diary [47]. Thus, it

may be that more intensive sessions that are focused on goal setting and monitoring is required to elicit changes.

There are also several other possible explanations for why our intervention did not have an impact on other social cognitive determinants. First, several of the social cognitive determinants in our study demonstrated high (or low) scores at baseline and might have demonstrated ceiling and floor effects. For example, the mean score for outcome expectations was close to 4 (possible range 1–5) and barriers was less than 0.7 (possible range 0–4). Second, response shift theory [48] posits that, as a consequence of an intervention, the participants' awareness and understanding of the variable being measured changes and alters each participant's perspective in his or her self-evaluation. For example, men who participated in the exercise program may have found that the physical outcomes that they expected to achieve from exercise may have changed from what they expected prior to taking up regular exercise. Vallance et al. [46] noted that response shift theory may provide an explanation for why researchers find changes in PA behavior, whereas cognitive variables remain relatively stable.

We found that increased social expectations of PA participation, barrier self-efficacy, and goal setting at follow-up predicted increased participation in VPA, which provides support for SCT. Increased barrier self-efficacy and reduced exercise motivation at postintervention also predicted improved VPA. Interventions that successfully improve perceptions of social outcomes of exercise, increase beliefs that one can overcome barriers to PA participation, and promote goal setting are likely to lead to improvements in VPA. In particular, barrier self-efficacy seems to be an important factor as it was associated with longer term VPA at both postintervention and follow-up, with large effect sizes. Furthermore, in a recent systematic review, goal setting has been identified as a significant predictor of PA behavior across a range of population groups [49] and should also be considered an important component of PA interventions. Future interventions should focus on these factors to improve participation in VPA among prostate cancer survivors.

Our study had a number of strengths. The randomized controlled study design provided the necessary causal relationships for examining mediation, and its prospective design minimized the potential for overestimation of associations that can occur when PA predictors are examined in cross-sectional studies [50]. Our examination of the mediators of follow-up VPA is also important to assess the factors that influence longer term behavior change. The limitations of this study must also be considered when interpreting our results. The intervention did not have a sustained impact on minutes of moderate/vigorous PA or meeting guidelines for PA, so we instead focused only on VPA. We used a self-report measure of PA, which can introduce measurement error.

Finally, although we followed guidelines for the development of measures of SCT determinants, some of our measures have not been validated in previous research.

Summary

Clinician referral to a combined supervised and home-based exercise program improved task self-efficacy, and this increase partially explained longer term engagement in VPA. Future PA behavior change interventions should focus on improving task self-efficacy, as well as social outcome expectations, goal setting, and barrier self-efficacy to improve VPA among men with prostate cancer. Furthermore, future research should examine the mediators of PA behavior for prostate cancer survivors so that the important mechanisms of behavior change can be identified and targeted.

Acknowledgements The authors acknowledge their colleagues who contributed to the main study: Jo Salmon, Steve F. Fraser, Suzanne Broadbent, Mari Botti, Bridie Kent, and the ENGAGE Uro-Oncology Clinicians' Group. This study was funded by the Australian Research Council (LP100200176) and the Prostate Cancer Foundation of Australia, with in-kind support from YMCA Victoria, Eastern Health, Epworth Healthcare, North Eastern Metropolitan Integrated Cancer Service, and Peter MacCallum Cancer Centre. Kerry S. Courneya is supported by the Canada Research Chairs Program.

Compliance with Ethical Standards

Authors' Contributions MJC, CJG, KSC, PML contributed to the intervention development, study design, and developed data collection procedures. MRM designed the data analysis plan, undertook data analysis and interpreted the statistical analysis. All authors contributed to interpreting the data. MJC drafted the manuscript. All authors read, contributed to and approved the final manuscript.

Conflict of Interest The authors declare that they have no conflict of interest.

Ethical Approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed Consent Informed consent was obtained from all individual participants included in the study.

References

1. Australian Institute of Health and Welfare. *Cancer in Australia: An Overview 2014*. Canberra: AIHW; 2014: 1–220.
2. Vanagas G, Mickeviciene A, Ulys A. Does quality of life of prostate cancer patients differ by stage and treatment? *Scand J Public Health*. 2013;41(1):58–64.
3. Chipperfield K, Fletcher J, Millar J, et al. Predictors of depression, anxiety and quality of life in patients with

- prostate cancer receiving androgen deprivation therapy. *Psychooncology*. 2013;22(10):2169–2176.
4. Keogh JW, MacLeod RD. Body composition, physical fitness, functional performance, quality of life, and fatigue benefits of exercise for prostate cancer patients: A systematic review. *J Pain Symptom Manage*. 2012; 43(1):96–110.
5. Baumann FT, Zopf EM, Bloch W. Clinical exercise interventions in prostate cancer patients—a systematic review of randomized controlled trials. *Support Care Cancer*. 2012;20(2):221–233.
6. Galvão DA, Newton RU, Gardiner RA, et al. Compliance to exercise-oncology guidelines in prostate cancer survivors and associations with psychological distress, unmet supportive care needs, and quality of life. *Psychooncology*. 2015;24(10):1241–1249.
7. Pinto BM, Floyd A. Theories underlying health promotion interventions among cancer survivors. *Semin Oncol Nurs*. 2008;24(3):153–163.
8. Michie S. Designing and implementing behaviour change interventions to improve population health. *J Health Serv Res Policy*. 2008;13(Suppl 3):S64–69.
9. Gollwitzer P. Implementation intentions. Strong effects of simple plans. *Am Psychol*. 1999;54(7):493–503.
10. McGowan EL, North S, Courneya KS. Randomized controlled trial of a behavior change intervention to increase physical activity and quality of life in prostate cancer survivors. *Ann Behav Med*. 2013;46(3):382–393.
11. Gardner JR, Livingston PM, Fraser SF. Effects of exercise on treatment-related adverse effects for patients with prostate cancer receiving androgen-deprivation therapy: A systematic review. *J Clin Oncol*. 2014;32(4):335–346.
12. Bandura A. *Social Foundations of Thought and Action: A Social Cognitive Theory*. Englewood Cliffs, NJ: Prentice-Hall; 1986.
13. Bandura A. Self-efficacy: Toward a unifying theory of behavioral change. *Psychol Rev*. 1977;84(2):191–215.
14. Bandura A. Health promotion by social cognitive means. *Health Educ Behav*. 2004;31(2):143–164.
15. Rogers LQ, Courneya KS, Anton PM, et al. Effects of the BEAT Cancer physical activity behavior change intervention on physical activity, aerobic fitness, and quality of life in breast cancer survivors: A multicenter randomized controlled trial. *Breast Cancer Res Treat*. 2015;149(1):109–119.
16. Pinto BM, Papandonatos GD, Goldstein MG, Marcus BH, Farrell N. Home-based physical activity intervention for colorectal cancer survivors. *Psychooncology*. 2013;22(1):54–64.
17. Mosher CE, Fuemmeler BF, Sloane R, et al. Change in self-efficacy partially mediates the effects of the FRESH START intervention on cancer survivors' dietary outcomes. *Psychooncology*. 2008;17(10):1014–1023.
18. Stacey FG, James EL, Chapman K, Courneya KS, Lubans DR. A systematic review and meta-analysis of social cognitive theory-based physical activity and/or nutrition behavior change interventions for cancer survivors. *J Cancer Surviv*. 2015;9(2):305–338.
19. Livingston PM, Craike MJ, Salmon J, et al.; ENGAGE Uro-Oncology Clinicians' Group. Effects of a clinician referral and exercise program for men who have completed active treatment for prostate cancer: A multicenter cluster randomized controlled trial (ENGAGE). *Cancer*. 2015;121(15):2646–2654.
20. Gaskin CJ, Craike M, Mohebbi M, Courneya KS, Livingston PM. A clinician referral and 12-week exercise training program for men with prostate cancer: Outcomes to 12 months

- of the ENGAGE cluster randomized controlled trial. *J Phys Act Health*. 2017;14(5):353–359.
21. Livingston PM, Salmon J, Courneya KS, et al. Efficacy of a referral and physical activity program for survivors of prostate cancer [ENGAGE]: Rationale and design for a cluster randomised controlled trial. *BMC Cancer*. 2011;11:237.
 22. Schmitz KH, Courneya KS, Matthews C, et al.; American College of Sports Medicine. American College of Sports Medicine roundtable on exercise guidelines for cancer survivors. *Med Sci Sports Exerc*. 2010;42(7):1409–1426.
 23. Hayes SC, Spence RR, Galvão DA, Newton RU. Australian Association for Exercise and Sport Science position stand: Optimising cancer outcomes through exercise. *J Sci Med Sport*. 2009;12(4):428–434.
 24. Bandura A. *Self-Efficacy: The Exercise of Control*. New York, NY: W. H. Freeman and Company; 1997.
 25. Craike MJ, Livingston PM, Botti M. An exploratory study of the factors that influence physical activity for prostate cancer survivors. *Support Care Cancer*. 2011;19(7):1019–1028.
 26. King AC, Haskell WL, Taylor CB, Kraemer HC, DeBusk RF. Group- vs home-based exercise training in healthy older men and women. A community-based clinical trial. *JAMA*. 1991;266(11):1535–1542.
 27. Courneya KS, Friedenreich CM, Quinney HA, et al. A longitudinal study of exercise barriers in colorectal cancer survivors participating in a randomized controlled trial. *Ann Behav Med*. 2005;29(2):147–153.
 28. Courneya KS, Stevinson C, McNeely ML, et al. Effects of supervised exercise on motivational outcomes and longer-term behavior. *Med Sci Sports Exerc*. 2012;44(3):542–549.
 29. Wójcicki TR, White SM, McAuley E. Assessing outcome expectations in older adults: The multidimensional outcome expectations for exercise scale. *J Gerontol b Psychol Sci Soc Sci*. 2009;64(1):33–40.
 30. McAuley E, Motl RW, White SM, Wójcicki TR. Validation of the multidimensional outcome expectations for exercise scale in ambulatory, symptom-free persons with multiple sclerosis. *Arch Phys Med Rehabil*. 2010;91(1):100–105.
 31. Bandura A. Guide for constructing self-efficacy scales. In: Pajares F, Urden T, eds. *Self-efficacy Beliefs of Adolescents*. Charlotte, NC: Information Age Publishing; 2006:307–337.
 32. Rogers LQ, Courneya KS, Verhulst S, Markwell S, Lanzotti V, Shah P. Exercise barrier and task self-efficacy in breast cancer patients during treatment. *Support Care Cancer*. 2006;14(1):84–90.
 33. Brawley LR, Culos-Reed SN, Angove J, Hoffman-Goetz L. Understanding the barriers to physical activity for cancer patients. *J Psychosoc Oncol*. 2002;20(4):1–21.
 34. Doerksen SE, Umstadtd MR, McAuley E. Social cognitive determinants of moderate and vigorous physical activity in college freshmen. *J Appl Soc Psychol*. 2009;39(5):1201–1213.
 35. Horton NJ, Kleinman KP. Much ado about nothing. *Am Stat*. 2007;61(1):79–90.
 36. Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *J Pers Soc Psychol*. 1986;51(6):1173–1182.
 37. Zhao X, Lynch JG, Chen Q. Reconsidering Baron and Kenny: Myths and truths about mediation analysis. *J Consum Res*. 2010;37(2):197–206.
 38. Cole DA, Maxwell SE. Testing mediational models with longitudinal data: Questions and tips in the use of structural equation modeling. *J Abnorm Psychol*. 2003;112(4):558–577.
 39. Demark-Wahnefried W, Clipp EC, Morey MC, et al. Lifestyle intervention development study to improve physical function in older adults with cancer: Outcomes from Project LEAD. *J Clin Oncol*. 2006;24(21):3465–3473.
 40. Ligibel JA, Meyerhardt J, Pierce JP, et al. Impact of a telephone-based physical activity intervention upon exercise behaviors and fitness in cancer survivors enrolled in a cooperative group setting. *Breast Cancer Res Treat*. 2012;132(1):205–213.
 41. Wang YJ, Boehmke M, Wu YW, Dickerson SS, Fisher N. Effects of a 6-week walking program on Taiwanese women newly diagnosed with early-stage breast cancer. *Cancer Nurs*. 2011;34(2):E1–13.
 42. French DP, Olander EK, Chisholm A, Mc Sharry J. Which behaviour change techniques are most effective at increasing older adults' self-efficacy and physical activity behaviour? A systematic review. *Ann Behav Med*. 2014;48(2):225–234.
 43. Young MD, Plotnikoff RC, Collins CE, Callister R, Morgan PJ. A test of social cognitive theory to explain men's physical activity during a gender-tailored weight loss program. *Am J Mens Health*. 2016;10(6):NP176–NP187.
 44. Perkins JM, Multhaup KS, Perkins HW, Barton C. Self-efficacy and participation in physical and social activity among older adults in Spain and the United States. *Gerontologist*. 2008;48(1):51–58.
 45. Jones LW, Courneya KS, Fairey AS, Mackey JR. Does the theory of planned behavior mediate the effects of an oncologist's recommendation to exercise in newly diagnosed breast cancer survivors? Results from a randomized controlled trial. *Health Psychol*. 2005;24(2):189–197.
 46. Vallance JK, Courneya KS, Plotnikoff RC, Mackey JR. Analyzing theoretical mechanisms of physical activity behavior change in breast cancer survivors: Results from the activity promotion (ACTION) trial. *Ann Behav Med*. 2008;35(2):150–158.
 47. Stacey FG, James EL, Chapman K, Lubans DR. Social cognitive theory mediators of physical activity in a lifestyle program for cancer survivors and carers: Findings from the ENRICH randomized controlled trial. *Int J Behav Nutr Phys Act*. 2016;13(1):49.
 48. Howard GS, Dailey PR. Response-shift bias: A source of contamination of self-report measures. *J Appl Psychol*. 1979;64(2):144–150.
 49. Young MD, Plotnikoff RC, Collins CE, Callister R, Morgan PJ. Social cognitive theory and physical activity: A systematic review and meta-analysis. *Obes Rev*. 2014;15(12):983–995.
 50. Weinstein ND. Misleading tests of health behavior theories. *Ann Behav Med*. 2007;33(1):1–10.